

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
8 July 2004 (08.07.2004)

PCT

(10) International Publication Number
WO 2004/056347 A2

(51) International Patent Classification⁷: **A61K 31/00**

(21) International Application Number:
PCT/DK2003/000931

(22) International Filing Date:
22 December 2003 (22.12.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PA 2002 01991 20 December 2002 (20.12.2002) DK
60/439,382 10 January 2003 (10.01.2003) US

(71) Applicant (for all designated States except US): **NOVO
NORDISK A/S** [DK/DK]; Nøve Allé, DK-2880 Bagsværd
(DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KAARSHOLM,
Niels, Christian** [DK/DK]; 38, Clausholmvej, DK-2720
Vanløse (DK). **MADSEN, Peter** [DK/DK]; 7 Ulveh-
jerg, DK-2880 Bagsværd (DK). **SCHLEIN, Morten**
[DK/DK]; Under Elmene 16, 4. tv., DK-2300 Copenhagen
S (DK). **OLSEN, Helle, Birk** [DK/DK]; Skoleladden 23,
DK-3450 Allerød (DK). **HAVELUND, Svend** [DK/DK];
Kurvej 24, DK-2880 Bagsværd (DK). **STEENSGAARD,
Dorte, Bjerre** [DK/DK]; Stockholmsgade 3, st. tv.,
DK- 2100_København Ø (DK). **LUDVIGSEN, Svend**
[DK/DK]; 13, Baunedalen, DK-3540 Lyngby (DK).

JAKOBSEN, Palle [DK/DK]; 14 Langkær Vænge,
DK-3500 Værløse (DK). **PETERSEN, Anders, Klarskov**
[DK/DK]; Nærum Hovedgade 64G, DK-2850 Nærum
(DK). **SCHLUCKEBIER, Gerd** [DE/DK]; Vævergade 5,
4th, DK-2200 København N (DK).

(74) Common Representative: **NOVO NORDISK A/S**; Nøve
Allé, DK-2880 Bagsværd (DK).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR,
CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,
MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU,
SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: **STABILISED INSULIN COMPOSITIONS**

(57) Abstract: The present invention provides pharmaceutical compositions comprising insulin and novel ligands for the His^{B10} Zn²⁺ sites of the R-state insulin hexamer. The resulting preparations have improved physical and chemical stability.



WO 2004/056347 A2

STABILISED INSULIN COMPOSITIONS

FIELD OF THE INVENTION

5 This invention relates to insulin compositions stabilised by adding ligands for the His^{B10} Zn²⁺ sites of the R-state insulin hexamer, as well as methods for preparation and use of such preparations.

BACKGROUND OF THE INVENTION

10 Diabetes is a general term for disorders in man having excessive urine excretion as in diabetes mellitus and diabetes insipidus. Diabetes mellitus is a metabolic disorder in which the ability to utilize glucose is partly or completely lost.


Since the discovery of insulin in the 1920's, continuous strides have been made to improve the treatment of diabetes mellitus. To help avoid extreme glycaemia levels, diabetic patients often practice multiple injection therapy, whereby insulin is administered with each meal. Many diabetic patients are treated with multiple daily insulin injections in a regimen 15 comprising one or two daily injections of a protracted insulin composition to cover the basal requirement, supplemented by bolus injections of rapid acting insulin to cover the meal-related requirements.

Insulin compositions having a protracted profile of action are well known in the art. Thus, one 20 main type of such insulin compositions comprises injectable aqueous suspensions of insulin crystals or amorphous insulin. Typically, the insulin in these compositions is provided in the form of protamine insulin, zinc insulin or protamine zinc insulin

Soluble, rapid acting insulin compositions usually comprise insulin, insulin analogue or insulin derivative together with zinc ion, phenolic preservative, isotonicity agent, and a buffer 25 substance. In addition, the preparation may optionally contain some salts and/or surfactants. Such preparations contain insulin in the form of an R-state hexamer.

Insulin Allostery.

30 The insulin hexamer is an allosteric protein that exhibits both positive and negative cooperativity and half-of-the-sites reactivity in ligand binding. This allosteric behaviour consists of two interrelated allosteric transitions designated L^A₀ and L^B₀, three inter-converting allosteric conformation states (eq. 1),



- 5 designated T_6 , T_3R_3 , and R_6 and two classes of allosteric ligand binding sites designated as the phenolic pockets and the His^{B10} anion sites. These allosteric sites are associated only with insulin subunits in the R conformation.

Insulin Hexamer Structures and Ligand Binding.

10

The T- to R-transition of the insulin hexamer involves transformation of the first nine residues of the B chain from an extended conformation in the T-state to an alpha-helical conformation in the R-state. This coil-to-helix transition causes the N-terminal residue, Phe^{B1} , to undergo an $\sim 30 \text{ \AA}$ change in position. This conformational change creates

15 hydrophobic pockets (the phenolic pockets) at the subunit interfaces (three in T_3R_3 , and six in R_6), and the new B-chain helices form 3-helix bundles (one in T_3R_3 and two in R_6) with the bundle axis aligned along the hexamer three-fold symmetry axis. The $\text{His}^{B10} \text{Zn}^{2+}$ in each R_3 unit is forced to change coordination geometry from octahedral to either tetrahedral (monodentate ligands) or pentahedral (bidentate ligands). Formation of the helix bundle

20 creates a narrow hydrophobic tunnel in each R_3 unit that extends from the surface $\sim 12 \text{ \AA}$ down to the His^{B10} metal ion. This tunnel and the $\text{His}^{B10} \text{Zn}^{2+}$ ion form the anion binding site. Ligands for the $\text{His}^{B10} \text{Zn}^{2+}$ sites of the R-state insulin hexamer have been disclosed in US 5830999.

Hexamer Ligand Binding and Stability of Insulin compositions.

The *in vivo* role of the T to R transition is unknown. However, the addition of allosteric ligands (e.g. phenol and chloride ion) to insulin compositions is widely used. Hexamerization is driven by coordination of Zn^{2+} at the His^{B10} sites to give T_6 . Following subcutaneous injection,

30 some dilution of the depot will take place over time and the ligands of soluble hexamers most likely diffuse away from the protein relatively rapidly. This is probably due to one or more phenomena including the binding of Zn^{2+} by surrounding tissue and albumin, the relatively larger space available for diffusion of the hydrophobic phenolic preservatives, and the generally larger diffusion coefficients characteristic of the smaller sized molecules.

35

Insulin compositions are usually stored for extended periods of time e.g. in vials or cartridges. Furthermore, insulin pumps are becoming more widely used, which places an additional demand on the chemical and physical stability of the insulin composition due to the elevated temperatures and physical stress these preparations are exposed to. There is thus
5 a need for insulin compositions that are more physically and chemically stable. It has been found that stabilising Zn^{2+} -site ligands may be added to insulin compositions to improve these properties.

SUMMARY OF THE INVENTION

10

The present invention provides pharmaceutical compositions comprising insulin and novel ligands for the $\text{His}^{\text{B10}} \text{Zn}^{2+}$ sites of the R-state insulin hexamer. The ligands belong to different subclasses of compounds, e.g. benzotriazoles, 3-hydroxy 2-napthoic acids, salicylic acids, tetrazoles, thiazolidinediones, 5-mercaptotetrazoles, or 4-cyano-1,2,3-triazoles. The insulin
15 may be rapid-acting. The insulin may be selected from human insulin, or an analogue or derivative thereof. The formulation may also comprise a phenolic compound, an isotonicity agent, and buffer. Also claimed are methods of treating type 1 or 2 diabetes comprising administration of a pharmaceutical composition of the invention.

DESCRIPTION OF THE DRAWINGS

20

Figures 1-8 show ThT assays of various combinations of insulin formulations and ligands of the invention.

Figure 9 shows disappearance rate of various combinations of insulin formulations and ligands of the invention from the subcutaneous depot following injection in pigs.

25 Figures 10-14 show reverse phase chromatography of various combinations of insulin formulations and ligands of the invention.

DEFINITIONS

The following is a detailed definition of the terms used to describe the invention:

30 "Halogen" designates an atom selected from the group consisting of F, Cl, Br and I.

The term "alkyl" as used herein represents a saturated, branched or straight hydrocarbon group having the indicated number of carbon atoms. Representative examples include, but

are not limited to, methyl, ethyl, n-propyl, isopropyl, butyl, isobutyl, *sec*-butyl, *tert*-butyl, n-pentyl, isopentyl, neopentyl, *tert*-pentyl, n-hexyl, isohexyl and the like.

The term "alkylene" as used herein represents a saturated, branched or straight bivalent hydrocarbon group having the indicated number of carbon atoms. Representative examples
5 include, but are not limited to, methylene, 1,2-ethylene, 1,3-propylene, 1,2-propylene, 1,4-butylene, 1,5-pentylene, 1,6-hexylene, and the like.

The term "alkenyl" as used herein represents a branched or straight hydrocarbon group having the indicated number of carbon atoms and at least one double bond. Examples of such groups include, but are not limited to, vinyl, 1-propenyl, 2-propenyl, iso-propenyl, 1,3-
10 butadienyl, 1-butenyl, 2-butenyl, 3-butenyl, 2-methyl-1-propenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 3-methyl-2-butenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 2,4-hexadienyl, 5-hexenyl and the like.

The term "alkynyl" as used herein represents a branched or straight hydrocarbon group having the indicated number of carbon atoms and at least one triple bond. Examples of such
15 groups include, but are not limited to, ethynyl, 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl, 3-butylnyl, 1-pentylnyl, 2-pentylnyl, 3-pentylnyl, 4-pentylnyl, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 5-hexynyl, 2,4-hexadiynyl and the like.

The term "alkoxy" as used herein refers to the radical -O- alkyl, wherein alkyl is as defined above. Representative examples are methoxy, ethoxy, n-propoxy, isopropoxy, butoxy, *sec*-
20 butoxy, *tert*-butoxy, pentoxy, isopentoxy, hexoxy, isohexoxy and the like.

The term "cycloalkyl" as used herein represents a saturated, carbocyclic group having the indicated number of carbon atoms. Representative examples are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl and the like.

The term "cycloalkenyl" as used herein represents a non-aromatic, carbocyclic group having
25 the indicated number of carbon atoms containing one or two double bonds. Representative examples are 1-cyclopentenyl, 2-cyclopentenyl, 3-cyclopentenyl, 1-cyclohexenyl, 2-cyclohexenyl, 3-cyclohexenyl, 2-cycloheptenyl, 3-cycloheptenyl, 2-cyclooctenyl, 1,4-cyclooctadienyl and the like.

The term "heterocyclyl" as used herein represents a non-aromatic 3 to 10 membered ring
30 containing one or more heteroatoms selected from nitrogen, oxygen and sulphur and optionally containing one or two double bonds. Representative examples are pyrrolidinyl, piperidyl, piperazinyl, morpholinyl, thiomorpholinyl, aziridinyl, tetrahydrofuranyl and the like.

The term "aryl" as used herein is intended to include carbocyclic, aromatic ring systems such as 6 membered monocyclic and 9 to 14 membered bi- and tricyclic, carbocyclic, aromatic
35 ring systems. Representative examples are phenyl, biphenyl, naphthyl, anthracenyl,

phenanthrenyl, fluorenyl, indenyl, azulenyl and the like. Aryl is also intended to include the partially hydrogenated derivatives of the ring systems enumerated above. Non-limiting examples of such partially hydrogenated derivatives are 1,2,3,4-tetrahydronaphthyl, 1,4-dihydronaphthyl and the like.

- 5 The term "arylene" as used herein is intended to include divalent, carbocyclic, aromatic ring systems such as 6 membered monocyclic and 9 to 14 membered bi- and tricyclic, divalent, carbocyclic, aromatic ring systems. Representative examples are phenylene, biphenylene, naphthylene, anthracenylene, phenanthrenylene, fluorenylene, indenylene, azulenylene and the like. Arylene is also intended to include the partially hydrogenated derivatives of the ring systems enumerated above. Non-limiting examples of such partially hydrogenated derivatives are 1,2,3,4-tetrahydronaphthylene, 1,4-dihydronaphthylene and the like.

10 The term "aryloxy" as used herein denotes a group -O-aryl, wherein aryl is as defined above.

The term "aroyl" as used herein denotes a group -C(O)-aryl, wherein aryl is as defined above.

- 15 The term "heteroaryl" as used herein is intended to include aromatic, heterocyclic ring systems containing one or more heteroatoms selected from nitrogen, oxygen and sulphur such as 5 to 7 membered monocyclic and 8 to 14 membered bi- and tricyclic aromatic, heterocyclic ring systems containing one or more heteroatoms selected from nitrogen, oxygen and sulphur. Representative examples are furyl, thienyl, pyrrolyl, pyrazolyl, 3-oxopyrazolyl, oxazolyl, thiazolyl, imidazolyl, isoxazolyl, isothiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, pyranlyl, pyridyl, pyridaziny, pyrimidinyl, pyrazinyl, 1,2,3-triazinyl, 1,2,4-triazinyl, 1,3,5-triazinyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, tetrazolyl, thiadiazinyl, indolyl, isoindolyl, benzofuryl, benzothieryl, indazolyl, benzimidazolyl, benzthiazolyl, 25 benzisothiazolyl, benzoxazolyl, benzisoxazolyl, purinyl, quinazolinyl, quinoliziny, quinoliny, isoquinoliny, quinoxaliny, naphthyridiny, pteridiny, carbazolyl, azepiny, diazepiny, acridiny, thiazolidiny, 2-thiooxothiazolidiny and the like. Heteroaryl is also intended to include the partially hydrogenated derivatives of the ring systems enumerated above. Non-limiting examples of such partially hydrogenated derivatives are 2,3-dihydrobenzofuranyl, pyrroliny, pyrazoliny, indoliny, oxazolidiny, oxazoliny, oxazepiny and the like.

- 30 The term "heteroarylene" as used herein is intended to include divalent, aromatic, heterocyclic ring systems containing one or more heteroatoms selected from nitrogen, oxygen and sulphur such as 5 to 7 membered monocyclic and 8 to 14 membered bi- and tricyclic aromatic, heterocyclic ring systems containing one or more heteroatoms selected from nitrogen, oxygen and sulphur. Representative examples are furylene, thienylene,
- 35

pyrrolylene, oxazolylenes, thiazolylenes, imidazolylenes, isoxazolylenes, isothiazolylenes, 1,2,3-triazolylenes, 1,2,4-triazolylenes, pyranylene, pyridylene, pyridazinylenes, pyrimidinylene, pyrazinylenes, 1,2,3-triazinylenes, 1,2,4-triazinylenes, 1,3,5-triazinylenes, 1,2,3-oxadiazolylenes, 1,2,4-oxadiazolylenes, 1,2,5-oxadiazolylenes, 1,3,4-oxadiazolylenes, 1,2,3-thiadiazolylenes, 1,2,4-thiadiazolylenes, 1,2,5-thiadiazolylenes, 1,3,4-thiadiazolylenes, tetrazolylenes, thiadiazinylenes, indolylenes, isoindolylenes, benzofurylenes, benzothienylenes, indazolylenes, benzimidazolylenes, benzthiazolylenes, benzisothiazolylenes, benzoxazolylenes, benzisoxazolylenes, purinylenes, quinazolinylene, quinolizinylenes, quinolinylene, isoquinolinylene, quinoxalinylenes, naphthyridinylenes, pteridinylenes, carbazolylenes, azepinylenes, diazepinylenes, acridinylenes and the like. Heteroaryl is also intended to include the partially hydrogenated derivatives of the ring systems enumerated above. Non-limiting examples of such partially hydrogenated derivatives are 2,3-dihydrobenzofuranylene, pyrrolinylenes, pyrazolinylene, indolinylene, oxazolidinylenes, oxazolinylene, oxazepinylenes and the like.

The term "ArG1" as used herein is intended to include an aryl or arylene radical as applicable, where aryl or arylene are as defined above but limited to phenyl, biphenyl, naphthyl, anthracenyl, phenanthrenyl, fluorenyl, indenyl, and azulenyl as well as the corresponding divalent radicals.

The term "ArG2" as used herein is intended to include an aryl or arylene radical as applicable, where aryl or arylene are as defined above but limited to phenyl, biphenyl, naphthyl, fluorenyl, and indenyl, as well as the corresponding divalent radicals.


The term "Het1" as used herein is intended to include a heteroaryl or heteroarylene radical as applicable, where heteroaryl or heteroarylene are as defined above but limited to furyl, thienyl, pyrrolyl, pyrazolyl, 3-oxopyrazolyl, oxazolyl, thiazolyl, imidazolyl, isoxazolyl, isothiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, pyranyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, 1,2,3-triazinyl, 1,2,4-triazinyl, 1,3,5-triazinyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, tetrazolyl, thiadiazinyl, indolyl, isoindolyl, benzofuryl, benzothienyl, indazolyl, benzimidazolyl, benzthiazolyl, benzisothiazolyl, benzoxazolyl, benzisoxazolyl, purinyl, quinazolinyl, quinolizinylenes, quinolinyl, isoquinolinyl, quinoxalinylenes, naphthyridinyl, pteridinyl, carbazolyl, azepinyl, diazepinyl, acridinyl, thiazolidinyl, 2-thiooxothiazolidinyl, as well as the corresponding divalent radicals.

The term "Het2" as used herein is intended to include a heteroaryl or heteroarylene radical as applicable, where heteroaryl or heteroarylene are as defined above but limited to furyl, thienyl, pyrrolyl, pyrazolyl, 3-oxopyrazolyl, oxazolyl, thiazolyl, imidazolyl, isoxazolyl, isothiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, pyranyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, 1,2,3-

triazinyl, 1,2,4-triazinyl, 1,3,5- triazinyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, tetrazolyl, thiadiazinyl, indolyl, isoindolyl, benzofuryl, benzothienyl, benzimidazolyl, benzthiazolyl, benzisothiazolyl, benzoxazolyl, benzisoxazolyl, quinoliny, isoquinoliny, quinoxaliny, carbazolyl, thiazolidiny, 2-thiooxothiazolidiny, as well as the corresponding divalent radicals.

The term "Het3" as used herein is intended to include a heteroaryl or heteroarylene radical as applicable, where heteroaryl or heteroarylene are as defined above but limited to furyl, thienyl, pyrrolyl, pyrazolyl, 3-oxopyrazolyl, oxazolyl, thiazolyl, imidazolyl, isoxazolyl, isothiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, pyridyl, tetrazolyl, indolyl, isoindolyl, benzofuryl, benzothienyl, benzimidazolyl, benzthiazolyl, benzisothiazolyl, benzoxazolyl, benzisoxazolyl, quinolyl, isoquinolyl, quinoxaliny, carbazolyl, thiazolidiny, 2-thiooxothiazolidiny, as well as the corresponding divalent radicals.

"Aryl-C₁-C₆-alkyl", "heteroaryl-C₁-C₆-alkyl", "aryl-C₂-C₆-alkenyl" etc. is intended to mean C₁-C₆-alkyl or C₂-C₆-alkenyl as defined above, substituted with an aryl or heteroaryl as defined above, for example:



The term "optionally substituted" as used herein means that the groups in question are either unsubstituted or substituted with one or more of the substituents specified. When the groups in question are substituted with more than one substituent the substituents may be the same or different.

Certain of the above defined terms may occur more than once in the structural formulae, and upon such occurrence each term shall be defined independently of the other.

Furthermore, when using the terms "independently are" and "independently selected from" it should be understood that the groups in question may be the same or different.

The term "substituted with one or more substituents" as used herein is intended to include one to four substituents, such as one to three substituents, one to two substituents, or even one substituent.

The term "treatment" as used herein means the management and care of a patient for the purpose of combating a disease, disorder or condition. The term is intended to include the delaying of the progression of the disease, disorder or condition, the alleviation or relief of

symptoms and complications, and/or the cure or elimination of the disease, disorder or condition. The patient to be treated is preferably a mammal, in particular a human being.

When in the specification or claims mention is made of groups of compounds such as benzotriazoles, 3-hydroxy 2-napthoic acids, salicylic acids, tetrazoles, thiazolidinediones, 5-mercaptotetrazoles, or 4-cyano-1,2,3-triazoles, these groups of compounds are intended to include also derivatives of the compounds from which the groups take their name.

The term "insulin" as used herein refers to human insulin as well as derivatives and analogues hereof as defined below.

10 The term "human insulin" as used herein refers to insulin naturally produced in the human body or recombinantly produced insulin identical thereto. Recombinant human insulin may be produced in any suitable host cell, for example the host cells may be bacterial, fungal (including yeast), insect, animal or plant cells.

The term "insulin derivative" as used herein (and related terms) refers to human insulin or an analogue thereof in which at least one organic substituent is bound to one or more of the amino acids.

By the term "analogue of human insulin" as used herein (and related terms) is meant human insulin in which one or more amino acids have been deleted and/or replaced by other amino acids, including non-codeable amino acids, or human insulin comprising additional amino acids, i.e. more than 51 amino acids, such that the resulting analogue possesses insulin activity.

Rapid acting insulin is intended to mean human insulin, insulin analogues or insulin derivatives having an onset of action after injection or any other form of administration faster or equal to that of soluble and neutral formulations of human insulin.

25 The term "phenolic compound" or similar expressions as used herein refers to a compound in which a hydroxyl group is bound directly to a benzene or substituted benzene ring. Examples of such compounds include, but are not limited to, phenol, o-cresol, m-cresol, p-cresol, chloro-cresol, thymol, and 7-hydroxyindole.

DESCRIPTION OF THE INVENTION

30

The present invention is based on the discovery that the two known ligand binding sites of the R-state insulin hexamer can be used to obtain an insulin composition having improved physical and chemical stability.

The basic concept underlying the present invention involves reversible attachment of a ligand to the His^{B10} Zn²⁺ site of the R-state hexamer. The anions currently used in insulin compositions as allosteric ligands for the R-state hexamers (notably chloride ion) bind only weakly to the His^{B10} anion site.


- 5 The His^{B10} Zn²⁺ site consists of a tunnel or cavity with a triangular-shaped cross-section that extends ~12 Å from the surface of the hexamer down to the His^{B10} Zn²⁺ ion. The diameter of the tunnel varies along its length and, depending on the nature of the ligand occupying the site, the opening can be capped over by the Asn^{B3} and Phe^{B1} side chains. The walls of the tunnel are made up of the side chains of the amino acid residues along one face each of the
- 10 three alpha-helices. The side chains from each helix that make up the lining of the tunnel are Phe^{B1}, Asn^{B3}, and Leu^{B6}. Therefore, except for the zinc ion, which is coordinated to three His^{B10} residues and is positioned at the bottom of the tunnel, the site is principally hydrophobic. Depending on the ligand structure, it may be possible for substituents on the ligand to make H-bonding interactions with Asn^{B3} and with the peptide linkage to Cys^{B7}.

15

In one aspect the invention provides a pharmaceutical composition comprising insulin and a zinc-binding ligand which reversibly binds to a HisB10 Zn²⁺ site of an insulin hexamer, wherein the ligand is selected from the group consisting of benzotriazoles, 3-hydroxy 2-naphthoic acids, salicylic acids, tetrazoles, thiazolidinediones, 5-mercaptotetrazoles, or 4-

20 cyano-1,2,3-triazoles, or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.

In one embodiment the invention provides a pharmaceutical composition wherein the zinc-binding ligand is



wherein

X is =O, =S or =NH

Y is -S-, -O- or -NH-

- 30 R¹ and R⁴ are independently selected from hydrogen or C₁-C₆-alkyl, R² is hydrogen or C₁-C₆-alkyl or aryl, R¹ and R² may optionally be combined to form a double bond,

R³ and R⁵ are independently selected from hydrogen, halogen, aryl, C₁-C₆-alkyl, or -C(O)NR¹¹R¹²,

A and B are independently selected from C₁-C₆-alkyl, aryl, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl or heteroaryl, wherein the alkyl or alkenyl is optionally substituted with one or more substituents independently selected from R⁸ and the aryl or heteroaryl is optionally substituted with up to four substituents R⁷, R⁸, R⁹, and R¹⁰,

A and R³ may be connected through one or two valence bonds, B and R⁵ may be connected through one or two valence bonds,

R⁶ is independently selected from halogen, -CN, -CF₃, -OCF₃, aryl, -COOH and -NH₂,
R⁷, R⁸, R⁹ and R¹⁰ are independently selected from

• hydrogen, halogen, -CN, -CH₂CN, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃,
-OCF₂CHF₂, -S(O)₂CF₃, -OS(O)₂CF₃, -SCF₃, -NO₂, -OR¹¹, -NR¹¹R¹², -SR¹¹,
-NR¹¹S(O)₂R¹², -S(O)₂NR¹¹R¹², -S(O)NR¹¹R¹², -S(O)R¹¹, -S(O)₂R¹¹, -OS(O)₂R¹¹,
-C(O)NR¹¹R¹², -OC(O)NR¹¹R¹², -NR¹¹C(O)R¹², -CH₂C(O)NR¹¹R¹²,
-OC₁-C₆-alkyl-C(O)NR¹¹R¹², -CH₂OR¹¹, -CH₂OC(O)R¹¹, -CH₂NR¹¹R¹², -OC(O)R¹¹,
-OC₁-C₁₅-alkyl-C(O)OR¹¹, -OC₁-C₆-alkyl-OR¹¹, -SC₁-C₆-alkyl-C(O)OR¹¹,
-C₂-C₆-alkenyl-C(=O)OR¹¹, -NR¹¹-C(=O)-C₁-C₆-alkyl-C(=O)OR¹¹,
-NR¹¹-C(=O)-C₁-C₆-alkenyl-C(=O)OR¹¹, -C(O)OR¹¹, C(O)R¹¹, or -C₂-C₆-alkenyl-C(=O)R¹¹, =O, or -C₂-C₆-alkenyl-C(=O)-NR¹¹R¹²,

• C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl, each of which may optionally be substituted with one or more substituents independently selected from R¹³,

• aryl, aryloxy, aryloxycarbonyl, aroyl, arylsulfanyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aroyl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl, heteroaryl-C₂-C₆-alkynyl, or C₃-C₆ cycloalkyl,

of which each cyclic moiety may optionally be substituted with one or more substituents independently selected from R¹⁴,

R¹¹ and R¹² are independently selected from hydrogen, OH, C₁-C₂₀-alkyl, aryl-C₁-C₆-alkyl or aryl, wherein the alkyl groups may optionally be substituted with one or more substituents independently selected from R¹⁵, and the aryl groups may optionally be substituted one or

more substituents independently selected from R^{16} ; R^{11} and R^{12} when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom, the heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds,

5

R^{13} is independently selected from halogen, -CN, -CF₃, -OCF₃, -OR¹¹, -C(O)OR¹¹, -NR¹¹R¹², and -C(O)NR¹¹R¹²,

10 R^{14} is independently selected from halogen, -C(O)OR¹¹, -CH₂C(O)OR¹¹, -CH₂OR¹¹, -CN, -CF₃, -OCF₃, -NO₂, -OR¹¹, -NR¹¹R¹², S(O)₂R¹¹, aryl and C₁-C₆-alkyl,

R^{15} is independently selected from halogen, -CN, -CF₃, -OCF₃, -OC₁-C₆-alkyl, -C(O)OC₁-C₆-alkyl, -COOH and -NH₂,

15 R^{16} is independently selected from halogen, -C(O)OC₁-C₆-alkyl, -COOH, -CN, -CF₃, -OCF₃, -NO₂, -OH, -OC₁-C₆-alkyl, -NH₂, C(=O) or C₁-C₆-alkyl, or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.

20 In another embodiment the invention provides a pharmaceutical composition wherein X is =O or =S.

In another embodiment the invention provides a pharmaceutical composition wherein X is =O.

25 In another embodiment the invention provides a pharmaceutical composition wherein X is =S.

In another embodiment the invention provides a pharmaceutical composition wherein Y is -O- or -S-.

In another embodiment the invention provides a pharmaceutical composition wherein Y is -O-.


30 In another embodiment the invention provides a pharmaceutical composition wherein Y is -S-.

In another embodiment the invention provides a pharmaceutical composition wherein A is aryl optionally substituted with up to four substituents, R^7 , R^8 , R^9 , and R^{10} which may be the same or different.

In another embodiment the invention provides a pharmaceutical composition wherein A is selected from ArG1 optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.

5 In another embodiment the invention provides a pharmaceutical composition wherein A is phenyl or naphthyl optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.

In another embodiment the invention provides a pharmaceutical composition wherein A is



10 In another embodiment the invention provides a pharmaceutical composition wherein A is phenyl.

In another embodiment the invention provides a pharmaceutical composition wherein A is heteroaryl optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.

15 In another embodiment the invention provides a pharmaceutical composition wherein A is selected from Het1 optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.


In another embodiment the invention provides a pharmaceutical composition wherein A is selected from Het2 optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.

20 In another embodiment the invention provides a pharmaceutical composition wherein A is selected from Het3 optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.

25 In another embodiment the invention provides a pharmaceutical composition wherein A is selected from the group consisting of indolyl, benzofuranyl, quinolyl, furyl, thienyl, or pyrrolyl, wherein each heteroaryl may optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.


In another embodiment the invention provides a pharmaceutical composition wherein A is benzofuranyl optionally substituted with up to four substituents R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.

30 In another embodiment the invention provides a pharmaceutical composition wherein A is




In another embodiment the invention provides a pharmaceutical composition wherein A is carbazolyl optionally substituted with up to four substituents R^7 , R^8 , R^9 , and R^{10} which may be the same or different.

- 5 In another embodiment the invention provides a pharmaceutical composition wherein A is




In another embodiment the invention provides a pharmaceutical composition wherein A is quinolyl optionally substituted with up to four substituents R^7 , R^8 , R^9 , and R^{10} which may be the same or different.

- 10 In another embodiment the invention provides a pharmaceutical composition wherein A is



In another embodiment the invention provides a pharmaceutical composition wherein A is indolyl optionally substituted with up to four substituents R^7 , R^8 , R^9 , and R^{10} which may be the same or different.

- 15 In another embodiment the invention provides a pharmaceutical composition wherein A is



In another embodiment the invention provides a pharmaceutical composition wherein R^1 is hydrogen.

- 20 In another embodiment the invention provides a pharmaceutical composition wherein R^2 is hydrogen.

In another embodiment the invention provides a pharmaceutical composition wherein R¹ and R² are combined to form a double bond.

In another embodiment the invention provides a pharmaceutical composition wherein R³ is C₁-C₆-alkyl, halogen, or C(O)NR¹⁶R¹⁷.

5 In another embodiment the invention provides a pharmaceutical composition wherein R³ is C₁-C₆-alkyl or C(O)NR¹⁶R¹⁷.

In another embodiment the invention provides a pharmaceutical composition wherein R³ is methyl.

10 In another embodiment the invention provides a pharmaceutical composition wherein B is phenyl optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.

In another embodiment the invention provides a pharmaceutical composition wherein R⁴ is hydrogen.

15 In another embodiment the invention provides a pharmaceutical composition wherein R⁵ is hydrogen.

In another embodiment the invention provides a pharmaceutical composition wherein R⁶ is aryl.

In another embodiment the invention provides a pharmaceutical composition wherein R⁶ is phenyl.

20 In another embodiment the invention provides a pharmaceutical composition wherein R⁷, R⁸, R⁹ and R¹⁰ are independently selected from

25 • hydrogen, halogen, -NO₂, -OR¹¹, -NR¹¹R¹², -SR¹¹, -NR¹¹S(O)₂R¹², -S(O)₂NR¹¹R¹², -S(O)NR¹¹R¹², -S(O)R¹¹, -S(O)₂R¹¹, -OS(O)₂R¹¹, -NR¹¹C(O)R¹², -CH₂OR¹¹, -CH₂OC(O)R¹¹, -CH₂NR¹¹R¹², -OC(O)R¹¹, -OC₁-C₆-alkyl-C(O)OR¹¹, -OC₁-C₆-alkyl-C(O)NR¹¹R¹², -OC₁-C₆-alkyl-OR¹¹, -SC₁-C₆-alkyl-C(O)OR¹¹, -C₂-C₆-alkenyl-C(=O)OR¹¹, -C(O)OR¹¹, or -C₂-C₆-alkenyl-C(=O)R¹¹,

30 • C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl, which may each optionally be substituted with one or more substituents independently selected from R¹³

35 • aryl, aryloxy, aroyl, arylsulfanyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aroyl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, wherein each of the cyclic moieties optionally may be substituted with one or more substituents independently selected from R¹⁴

In another embodiment the invention provides a pharmaceutical composition wherein R^7 , R^8 , R^9 and R^{10} are independently selected from

- 5 • hydrogen, halogen, $-\text{NO}_2$, $-\text{OR}^{11}$, $-\text{NR}^{11}\text{R}^{12}$, $-\text{SR}^{11}$, $-\text{S}(\text{O})_2\text{R}^{11}$, $-\text{OS}(\text{O})_2\text{R}^{11}$, $-\text{CH}_2\text{OC}(\text{O})\text{R}^{11}$, $-\text{OC}(\text{O})\text{R}^{11}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{11}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-OR}^{11}$, $-\text{SC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{11}$, $-\text{C}(\text{O})\text{OR}^{11}$, or $-\text{C}_2\text{-C}_6\text{-alkenyl-C}(\text{=O})\text{R}^{11}$,
- 10 • $\text{C}_1\text{-C}_6\text{-alkyl}$ or $\text{C}_1\text{-C}_6\text{-alkenyl}$ which may each optionally be substituted with one or more substituents independently selected from R^{13}
- aryl, aryloxy, aroyl, aryl- $\text{C}_1\text{-C}_6\text{-alkoxy}$, aryl- $\text{C}_1\text{-C}_6\text{-alkyl}$, heteroaryl,
- 15 of which each of the cyclic moieties optionally may be substituted with one or more substituents independently selected from R^{14}

In another embodiment the invention provides a pharmaceutical composition wherein R^7 , R^8 , R^9 and R^{10} are independently selected from

- 20 • hydrogen, halogen, $-\text{NO}_2$, $-\text{OR}^{11}$, $-\text{NR}^{11}\text{R}^{12}$, $-\text{SR}^{11}$, $-\text{S}(\text{O})_2\text{R}^{11}$, $-\text{OS}(\text{O})_2\text{R}^{11}$, $-\text{CH}_2\text{OC}(\text{O})\text{R}^{11}$, $-\text{OC}(\text{O})\text{R}^{11}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{11}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-OR}^{11}$, $-\text{SC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{11}$, $-\text{C}(\text{O})\text{OR}^{11}$, or $-\text{C}_2\text{-C}_6\text{-alkenyl-C}(\text{=O})\text{R}^{11}$,
- 25 • $\text{C}_1\text{-C}_6\text{-alkyl}$ or $\text{C}_1\text{-C}_6\text{-}$ which may each optionally be substituted with one or more substituents independently selected from R^{13}
- aryl, aryloxy, aroyl, aryl- $\text{C}_1\text{-C}_6\text{-alkoxy}$, aryl- $\text{C}_1\text{-C}_6\text{-alkyl}$, heteroaryl,
- 30 of which each of the cyclic moieties optionally may be substituted with one or more substituents independently selected from R^{14} .

In another embodiment the invention provides a pharmaceutical composition wherein R^7 , R^8 , R^9 and R^{10} are independently selected from

- 35 • hydrogen, halogen, $-\text{OR}^{11}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{11}$, or $-\text{C}(\text{O})\text{OR}^{11}$,

• C₁-C₆-alkyl which may each optionally be substituted with one or more substituents independently selected from R¹³

• aryl, aryloxy, aryl-C₁-C₆-alkoxy,

5

of which each of the cyclic moieties optionally may be substituted with one or more substituents independently selected from R¹⁴.

10 In another embodiment the invention provides a pharmaceutical composition wherein R⁷, R⁸, R⁹ and R¹⁰ are independently selected from

• hydrogen, halogen, -OR¹¹, -OC₁-C₆-alkyl-C(O)OR¹¹, or -C(O)OR¹¹,

• C₁-C₆-alkyl which may optionally be substituted with one or more substituents independently selected from R¹³

15

• phenyl, phenyloxy, phenyl-C₁-C₆-alkoxy, wherein each of the cyclic moieties optionally may be substituted with one or more substituents independently selected from R¹⁴.

20 In another embodiment the invention provides a pharmaceutical composition wherein R¹¹ and R¹² are independently selected from hydrogen, C₁-C₂₀-alkyl, aryl or aryl-C₁-C₆-alkyl, wherein the alkyl groups may optionally be substituted with one or more substituents independently selected from R¹⁵, and the aryl groups may optionally be substituted one or more substituents independently selected from R¹⁶; R¹¹ and R¹² when attached to the same
25 nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom, the heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds.

In another embodiment the invention provides a pharmaceutical composition wherein R¹¹ and R¹² are independently selected from hydrogen, C₁-C₂₀-alkyl, aryl or aryl-C₁-C₆-alkyl, wherein
30 the alkyl groups may optionally be substituted with one or more substituents independently selected from R¹⁵, and the aryl groups may optionally be substituted one or more substituents independently selected from R¹⁶.

In another embodiment the invention provides a pharmaceutical composition wherein R¹¹ and R¹² are independently selected from phenyl or phenyl-C₁-C₆-alkyl.

In another embodiment the invention provides a pharmaceutical composition wherein one or both of R^{11} and R^{12} are methyl.

In another embodiment the invention provides a pharmaceutical composition wherein R^{13} is independently selected from halogen, CF_3 , OR^{11} or $NR^{11}R^{12}$.

5 In another embodiment the invention provides a pharmaceutical composition wherein R^{13} is independently selected from halogen or OR^{11} .

In another embodiment the invention provides a pharmaceutical composition wherein R^{13} is OR^{11} .

10 In another embodiment the invention provides a pharmaceutical composition wherein R^{14} is independently selected from halogen, $-C(O)OR^{11}$, $-CN$, $-CF_3$, $-OR^{11}$, $S(O)_2R^{11}$, and C_1 - C_6 -alkyl.

In another embodiment the invention provides a pharmaceutical composition wherein R^{14} is independently selected from halogen, $-C(O)OR^{11}$, or $-OR^{11}$.


15 In another embodiment the invention provides a pharmaceutical composition wherein R^{15} is independently selected from halogen, $-CN$, $-CF_3$, $-C(O)OC_1$ - C_6 -alkyl, and $-COOH$.

In another embodiment the invention provides a pharmaceutical composition wherein R^{15} is independently selected from halogen or $-C(O)OC_1$ - C_6 -alkyl.

20 In another embodiment the invention provides a pharmaceutical composition wherein R^{16} is independently selected from halogen, $-C(O)OC_1$ - C_6 -alkyl, $-COOH$, $-NO_2$, $-OC_1$ - C_6 -alkyl, $-NH_2$, $C(=O)$ or C_1 - C_6 -alkyl.

In another embodiment the invention provides a pharmaceutical composition wherein R^{16} is independently selected from halogen, $-C(O)OC_1$ - C_6 -alkyl, $-COOH$, $-NO_2$, or C_1 - C_6 -alkyl.

In another embodiment the invention provides a pharmaceutical composition wherein the zinc-binding ligand is



wherein

R^{19} is hydrogen or C_1 - C_6 -alkyl,

R^{20} is hydrogen or C_1 - C_6 -alkyl,

30

D and F are a valence bond or C_1 - C_6 -alkylene optionally substituted with one or more substituents independently selected from R^{72} ,

R⁷² is independently selected from hydroxy, C₁-C₆-alkyl, or aryl,

E is C₁-C₆-alkyl, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with up to three substituents R²¹, R²² and R²³,

G is C₁-C₆-alkyl, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with up to three substituents R²⁴, R²⁵ and R²⁶,

R¹⁷, R¹⁸, R²¹, R²², R²³, R²⁴, R²⁵ and R²⁶ are independently selected from

• hydrogen, halogen, -CN, -CH₂CN, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -S(O)₂CF₃, -SCF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷S(O)₂R²⁸, -S(O)₂NR²⁷R²⁸, -S(O)NR²⁷R²⁸, -S(O)R²⁷, -S(O)₂R²⁷, -C(O)NR²⁷R²⁸, -OC(O)NR²⁷R²⁸, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -CH₂C(O)NR²⁷R²⁸, -OCH₂C(O)NR²⁷R²⁸, -CH₂OR²⁷, -CH₂NR²⁷R²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -NR²⁷-C(=O)-C₁-C₆-alkyl-C(=O)OR²⁷, -NR²⁷-C(=O)-C₁-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

• C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl,

which may optionally be substituted with one or more substituents independently selected from R²⁹,

• aryl, aryloxy, aryloxycarbonyl, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl or heteroaryl-C₂-C₆-alkynyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰,

R²⁷ and R²⁸ are independently selected from hydrogen, C₁-C₆-alkyl, aryl-C₁-C₆-alkyl or aryl, or R²⁷ and R²⁸ when attached to the same nitrogen atom together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further

heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds,

R^{29} is independently selected from halogen, -CN, -CF₃, -OCF₃, -OR²⁷, and -NR²⁷R²⁸,

5

R^{30} is independently selected from halogen, -C(O)OR²⁷, -CN, -CF₃, -OCF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸ and C₁-C₆-alkyl, or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.

10 In another embodiment the invention provides a pharmaceutical composition wherein D is a valence bond.

In another embodiment the invention provides a pharmaceutical composition wherein D is C₁-C₆-alkylene optionally substituted with one or more hydroxy, C₁-C₆-alkyl, or aryl.


15 In another embodiment the invention provides a pharmaceutical composition wherein E is aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with up to three substituents independently selected from R²¹, R²² and R²³.

In another embodiment the invention provides a pharmaceutical composition wherein E is aryl optionally substituted with up to three substituents independently selected from R²¹, R²² and R²³.

20 In another embodiment the invention provides a pharmaceutical composition wherein E is selected from ArG1 and optionally substituted with up to three substituents independently selected from R²¹, R²² and R²³.

In another embodiment the invention provides a pharmaceutical composition wherein E is phenyl optionally substituted with up to three substituents independently selected from R²¹, R²² and R²³.

25 In another embodiment the invention provides a pharmaceutical composition wherein the zinc-binding ligand is



30 In another embodiment the invention provides a pharmaceutical composition wherein R²¹, R²² and R²³ are independently selected from

• hydrogen, halogen, $-\text{CHF}_2$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OCHF}_2$, $-\text{OCH}_2\text{CF}_3$, $-\text{OCF}_2\text{CHF}_2$, $-\text{SCF}_3$, $-\text{NO}_2$, $-\text{OR}^{27}$, $-\text{NR}^{27}\text{R}^{28}$, $-\text{SR}^{27}$, $-\text{C}(\text{O})\text{NR}^{27}\text{R}^{28}$, $-\text{OC}(\text{O})\text{NR}^{27}\text{R}^{28}$, $-\text{NR}^{27}\text{C}(\text{O})\text{R}^{28}$, $-\text{NR}^{27}\text{C}(\text{O})\text{OR}^{28}$, $-\text{CH}_2\text{C}(\text{O})\text{NR}^{27}\text{R}^{28}$, $-\text{OCH}_2\text{C}(\text{O})\text{NR}^{27}\text{R}^{28}$, $-\text{CH}_2\text{OR}^{27}$, $-\text{CH}_2\text{NR}^{27}\text{R}^{28}$, $-\text{OC}(\text{O})\text{R}^{27}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{SC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}_2\text{-C}_6\text{-alkenyl-C}(\text{O})\text{OR}^{27}$, $-\text{NR}^{27}\text{-C}(\text{O})\text{-C}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{NR}^{27}\text{-C}(\text{O})\text{-C}_1\text{-C}_6\text{-alkenyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}(\text{O})\text{NR}^{27}\text{-C}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, or $-\text{C}(\text{O})\text{OR}^{27}$,

• $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_2\text{-C}_6\text{-alkenyl}$ or $\text{C}_2\text{-C}_6\text{-alkynyl}$,

which may optionally be substituted with one or more substituents independently selected from R^{29}

• aryl, aryloxy, aryloxycarbonyl, aroyl, aryl- $\text{C}_1\text{-C}_6\text{-alkoxy}$, aryl- $\text{C}_1\text{-C}_6\text{-alkyl}$, aryl- $\text{C}_2\text{-C}_6\text{-alkenyl}$, aryl- $\text{C}_2\text{-C}_6\text{-alkynyl}$, heteroaryl, heteroaryl- $\text{C}_1\text{-C}_6\text{-alkyl}$, heteroaryl- $\text{C}_2\text{-C}_6\text{-alkenyl}$ or heteroaryl- $\text{C}_2\text{-C}_6\text{-alkynyl}$,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R^{30} .

In another embodiment the invention provides a pharmaceutical composition wherein R^{21} , R^{22} and R^{23} are independently selected from

• hydrogen, halogen, $-\text{OCF}_3$, $-\text{OR}^{27}$, $-\text{NR}^{27}\text{R}^{28}$, $-\text{SR}^{27}$, $-\text{NR}^{27}\text{C}(\text{O})\text{R}^{28}$, $-\text{NR}^{27}\text{C}(\text{O})\text{OR}^{28}$, $-\text{OC}(\text{O})\text{R}^{27}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{SC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}_2\text{-C}_6\text{-alkenyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}(\text{O})\text{NR}^{27}\text{-C}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, or $-\text{C}(\text{O})\text{OR}^{27}$,

• $\text{C}_1\text{-C}_6\text{-alkyl}$ optionally substituted with one or more substituents independently selected from R^{29}

• aryl, aryloxy, aroyl, aryl- $\text{C}_1\text{-C}_6\text{-alkoxy}$, aryl- $\text{C}_1\text{-C}_6\text{-alkyl}$, heteroaryl, heteroaryl- $\text{C}_1\text{-C}_6\text{-alkyl}$,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

In another embodiment the invention provides a pharmaceutical composition wherein R²¹,

5 R²² and R²³ are independently selected from

• hydrogen, halogen, -OCF₃, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸,
-OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-
C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or
10 -C(O)OR²⁷,

• methyl, ethyl propyl optionally substituted with one or more substituents
independently selected from R²⁹

15 • aryl, aryloxy, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, heteroaryl, heteroaryl-C₁-C₆-
alkyl

of which the cyclic moieties optionally may be substituted with one or more
substituents selected from R³⁰.

In another embodiment the invention provides a pharmaceutical composition wherein R²¹,

20 R²² and R²³ are independently selected from

• hydrogen, halogen, -OCF₃, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸,
-OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-
C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or
25 -C(O)OR²⁷,

• methyl, ethyl propyl optionally substituted with one or more substituents
independently selected from R²⁹

30 • ArG1, ArG1-O-, ArG1-C(O)-, ArG1-C₁-C₆-alkoxy, ArG1-C₁-C₆-alkyl, Het3, Het3-C₁-
C₆-alkyl

of which the cyclic moieties optionally may be substituted with one or more substituents
selected from R³⁰.

In another embodiment the invention provides a pharmaceutical composition wherein R²¹,

35 R²² and R²³ are independently selected from

• hydrogen, halogen, $-\text{OCF}_3$, $-\text{OR}^{27}$, $-\text{NR}^{27}\text{R}^{28}$, $-\text{SR}^{27}$, $-\text{NR}^{27}\text{C}(\text{O})\text{R}^{28}$, $-\text{NR}^{27}\text{C}(\text{O})\text{OR}^{28}$, $-\text{OC}(\text{O})\text{R}^{27}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{SC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}_2\text{-C}_6\text{-alkenyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}(\text{=O})\text{NR}^{27}\text{-C}_1\text{-C}_6\text{-alkyl-C}(\text{=O})\text{OR}^{27}$, $-\text{C}_1\text{-C}_6\text{-alkyl-C}(\text{=O})\text{OR}^{27}$, or $-\text{C}(\text{O})\text{OR}^{27}$,

• $\text{C}_1\text{-C}_6\text{-alkyl}$ optionally substituted with one or more substituents independently selected from R^{29}

• phenyl, phenyloxy, phenyl- $\text{C}_1\text{-C}_6\text{-alkoxy}$, phenyl- $\text{C}_1\text{-C}_6\text{-alkyl}$, of which the cyclic moieties optionally may be substituted with one or more substituents selected from R^{30} .

In another embodiment the invention provides a pharmaceutical composition wherein R^{19} is hydrogen or methyl.

In another embodiment the invention provides a pharmaceutical composition wherein R^{19} is hydrogen.

In another embodiment the invention provides a pharmaceutical composition wherein R^{27} is Hydrogen, $\text{C}_1\text{-C}_6\text{-alkyl}$ or aryl.

In another embodiment the invention provides a pharmaceutical composition wherein R^{27} is hydrogen or $\text{C}_1\text{-C}_6\text{-alkyl}$.

In another embodiment the invention provides a pharmaceutical composition wherein R^{28} is hydrogen or $\text{C}_1\text{-C}_6\text{-alkyl}$.

In another embodiment the invention provides a pharmaceutical composition wherein F is a valence bond.

In another embodiment the invention provides a pharmaceutical composition wherein F is $\text{C}_1\text{-C}_6\text{-alkylene}$ optionally substituted with one or more hydroxy, $\text{C}_1\text{-C}_6\text{-alkyl}$, or aryl.

In another embodiment the invention provides a pharmaceutical composition wherein G is $\text{C}_1\text{-C}_6\text{-alkyl}$ or aryl, wherein the aryl is optionally substituted with up to three substituents R^{24} , R^{25} and R^{26} .

In another embodiment the invention provides a pharmaceutical composition wherein G is $\text{C}_1\text{-C}_6\text{-alkyl}$ or ArG1 , wherein the aryl is optionally substituted with up to three substituents R^{24} , R^{25} and R^{26} .

In another embodiment the invention provides a pharmaceutical composition wherein G is $\text{C}_1\text{-C}_6\text{-alkyl}$.

In another embodiment the invention provides a pharmaceutical composition wherein G is phenyl optionally substituted with up to three substituents R²⁴, R²⁵ and R²⁶.

In another embodiment the invention provides a pharmaceutical composition wherein R²⁴, R²⁵ and R²⁶ are independently selected from

• hydrogen, halogen, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -SCF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -C(O)NR²⁷R²⁸, -OC(O)NR²⁷R²⁸, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -CH₂C(O)NR²⁷R²⁸, -OCH₂C(O)NR²⁷R²⁸, -CH₂OR²⁷, -CH₂NR²⁷R²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -NR²⁷-C(=O)-C₁-C₆-alkyl-C(=O)OR²⁷, -NR²⁷-C(=O)-C₁-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

• C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl,

which may optionally be substituted with one or more substituents independently selected from R²⁹

• aryl, aryloxy, aryloxycarbonyl, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl or heteroaryl-C₂-C₆-alkynyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

In another embodiment the invention provides a pharmaceutical composition wherein R²⁴, R²⁵ and R²⁶ are independently selected from

• hydrogen, halogen, -OCF₃, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

• C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl,

which may optionally be substituted with one or more substituents independently selected from R²⁹

•aryl, aryloxy, aryloxycarbonyl, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl or heteroaryl-C₂-C₆-alkynyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

In another embodiment the invention provides a pharmaceutical composition wherein R²⁴, R²⁵ and R²⁶ are independently selected from

•hydrogen, halogen, -OCF₃, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

•C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R²⁹

•aryl, aryloxy, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, heteroaryl, heteroaryl-C₁-C₆-alkyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

In another embodiment the invention provides a pharmaceutical composition wherein R²¹, R²² and R²³ are independently selected from

•hydrogen, halogen, -OCF₃, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

•methyl, ethyl propyl optionally substituted with one or more substituents independently selected from R²⁹

•ArG1, ArG1-O-, ArG1-C(O)-, ArG1-C₁-C₆-alkoxy, ArG1-C₁-C₆-alkyl, Het3, Het3-C₁-C₆-alkyl

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

In another embodiment the invention provides a pharmaceutical composition wherein R²¹, R²² and R²³ are independently selected from

•hydrogen, halogen, -OCF₃, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

•methyl, ethyl propyl optionally substituted with one or more substituents independently selected from R²⁹

•ArG1, ArG1-O-, ArG1-C(O)-, ArG1-C₁-C₆-alkoxy, ArG1-C₁-C₆-alkyl, Het3, Het3-C₁-C₆-alkyl

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

In another embodiment the invention provides a pharmaceutical composition wherein R²¹, R²² and R²³ are independently selected from

•hydrogen, halogen, -OCF₃, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

•methyl, ethyl propyl optionally substituted with one or more substituents independently selected from R²⁹

•ArG1, ArG1-O-, ArG1-C₁-C₆-alkoxy, ArG1-C₁-C₆-alkyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

In another embodiment the invention provides a pharmaceutical composition wherein R²⁰ is hydrogen or methyl.

5 In another embodiment the invention provides a pharmaceutical composition wherein R²⁰ is hydrogen.

In another embodiment the invention provides a pharmaceutical composition wherein R²⁷ is hydrogen, C₁-C₆-alkyl or aryl.

In another embodiment the invention provides a pharmaceutical composition wherein R²⁷ is hydrogen or C₁-C₆-alkyl or ArG1.

10 In another embodiment the invention provides a pharmaceutical composition wherein R²⁷ is hydrogen or C₁-C₆-alkyl.

In another embodiment the invention provides a pharmaceutical composition wherein R²⁸ is hydrogen or C₁-C₆-alkyl.

15 In another embodiment the invention provides a pharmaceutical composition wherein R¹⁷ and R¹⁸ are independently selected from

• hydrogen, halogen, -CN, -CF₃, -OCF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -S(O)R²⁷, -S(O)₂R²⁷, -C(O)NR²⁷R²⁸, -CH₂OR²⁷, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, or -C(O)OR²⁷,

20

• C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl, optionally substituted with one or more substituents independently selected from R²⁹

• aryl, aryloxy, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, heteroaryl, heteroaryl-C₁-C₆-alkyl,

25

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

30 In another embodiment the invention provides a pharmaceutical composition wherein R¹⁷ and R¹⁸ are independently selected from

• hydrogen, halogen, -CN, -CF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, or -C(O)OR²⁷,

• C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R²⁹

35

•aryl, aryloxy, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, heteroaryl, heteroaryl-C₁-C₆-alkyl,

5 of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

In another embodiment the invention provides a pharmaceutical composition wherein R¹⁷ and R¹⁸ are independently selected from

- 10
- hydrogen, halogen, -CN, -CF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, or -C(O)OR²⁷
 - methyl, ethyl propyl optionally substituted with one or more substituents independently selected from R²⁹
 - aryl, aryloxy, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, heteroaryl, heteroaryl-C₁-C₆-alkyl

15 of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

In another embodiment the invention provides a pharmaceutical composition wherein R¹⁷ and R¹⁸ are independently selected from

- 20
- hydrogen, halogen, -CN, -CF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, or -C(O)OR²⁷
 - methyl, ethyl propyl optionally substituted with one or more substituents independently selected from R²⁹
 - ArG1, ArG1-O-, ArG1-C(O)-, ArG1-C₁-C₆-alkoxy, ArG1-C₁-C₆-alkyl, Het3, Het3-C₁-C₆-alkyl

25 of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

In another embodiment the invention provides a pharmaceutical composition wherein R¹⁷ and R¹⁸ are independently selected from

- 30
- hydrogen, halogen, -CN, -CF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, or -C(O)OR²⁷
 - C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R²⁹
 - phenyl, phenyloxy, phenyl-C₁-C₆-alkoxy, phenyl-C₁-C₆-alkyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

35 In another embodiment the invention provides a pharmaceutical composition wherein R²⁷ is hydrogen or C₁-C₆-alkyl.


In another embodiment the invention provides a pharmaceutical composition wherein R^{27} is hydrogen, methyl or ethyl.

In another embodiment the invention provides a pharmaceutical composition wherein R^{28} is hydrogen or C_1 - C_6 -alkyl.

- 5 In another embodiment the invention provides a pharmaceutical composition wherein R^{28} is hydrogen, methyl or ethyl.


In another embodiment the invention provides a pharmaceutical composition wherein R^{72} is -OH or phenyl.

- 10 In another embodiment the invention provides a pharmaceutical composition wherein the zinc-binding ligand is



- 15 In another embodiment the invention provides a pharmaceutical composition wherein the zinc-binding ligand is of the form H-I-J


wherein H is



- 25 wherein the phenyl, naphthalene or benzocarbazole rings are optionally substituted with one or more substituents independently selected from R^{31}

I is selected from

- a valence bond,
- $-CH_2N(R^{32})-$ or $-SO_2N(R^{33})-$,



- 30 • wherein Z^1 is $S(O)_2$ or CH_2 , Z^2 is -NH-, -O- or -S-, and n is 1 or 2,

J is

- C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl, which may each optionally be substituted with one or more substituents selected from R³⁴,
- Aryl, aryloxy, aryl-oxycarbonyl-, aroyl, aryl-C₁-C₆-alkoxy-, aryl-C₁-C₆-alkyl-, aryl-C₂-C₆-alkenyl-, aryl-C₂-C₆-alkynyl-, heteroaryl, heteroaryl-C₁-C₆-alkyl-, heteroaryl-C₂-C₆-alkenyl- or heteroaryl-C₂-C₆-alkynyl-, wherein the cyclic moieties are optionally substituted with one or more substituents selected from R³⁷,
- hydrogen,

- 10 R³¹ is independently selected from hydrogen, halogen, -CN, -CH₂CN, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -S(O)₂CF₃, -SCF₃, -NO₂, -OR³⁵, -C(O)R³⁵, -NR³⁵R³⁶, -SR³⁵, -NR³⁵S(O)₂R³⁶, -S(O)₂NR³⁵R³⁶, -S(O)NR³⁵R³⁶, -S(O)R³⁵, -S(O)₂R³⁵, -C(O)NR³⁵R³⁶, -OC(O)NR³⁵R³⁶, -NR³⁵C(O)R³⁶, -CH₂C(O)NR³⁵R³⁶, -OCH₂C(O)NR³⁵R³⁶, -CH₂OR³⁵, -CH₂NR³⁵R³⁶, -OC(O)R³⁵, -OC₁-C₆-alkyl-C(O)OR³⁵, -SC₁-C₆-alkyl-C(O)OR³⁵, -C₂-C₆-alkenyl-C(=O)OR³⁵, -NR³⁵-C(=O)-C₁-C₆-alkyl-C(=O)OR³⁵, -NR³⁵-C(=O)-C₁-C₆-alkenyl-C(=O)OR³⁵, C₁-C₆-alkyl, C₁-C₆-alkanoyl or -C(O)OR³⁵,
- 15

R³² and R³³ are independently selected from hydrogen, C₁-C₆-alkyl or C₁-C₆-alkanoyl,

- 20 R³⁴ is independently selected from halogen, -CN, -CF₃, -OCF₃, -OR³⁵, and -NR³⁵R³⁶,

R³⁵ and R³⁶ are independently selected from hydrogen, C₁-C₆-alkyl, aryl-C₁-C₆-alkyl or aryl, or R³⁵ and R³⁶ when attached to the same nitrogen atom together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds,


25

R³⁷ is independently selected from halogen, -C(O)OR³⁵, -C(O)H, -CN, -CF₃, -OCF₃, -NO₂, -OR³⁵, -NR³⁵R³⁶, C₁-C₆-alkyl or C₁-C₆-alkanoyl,

30

or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.


- 35 In another embodiment the invention provides a pharmaceutical composition wherein the zinc-binding ligand is of the form H-I-J, wherein H is



wherein the phenyl, naphthalene or benzocarbazole rings are optionally substituted with one or more substituents independently selected from R^{31} ,

I is selected from

- 5 • a valence bond,
 • $-\text{CH}_2\text{N}(\text{R}^{32})-$ or $-\text{SO}_2\text{N}(\text{R}^{33})-$,



- wherein Z^1 is $\text{S}(\text{O})_2$ or CH_2 , Z^2 is N,-O- or -S-, and n is 1 or 2,

J is

- 10 • $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_2\text{-C}_6\text{-alkenyl}$ or $\text{C}_2\text{-C}_6\text{-alkynyl}$, which may each optionally be substituted with one or more substituents selected from R^{34} ,
 • Aryl, aryloxy, aryl-oxycarbonyl-, aroyl, aryl- $\text{C}_1\text{-C}_6\text{-alkoxy-}$, aryl- $\text{C}_1\text{-C}_6\text{-alkyl-}$, aryl- $\text{C}_2\text{-C}_6\text{-alkenyl-}$, aryl- $\text{C}_2\text{-C}_6\text{-alkynyl-}$, heteroaryl, heteroaryl- $\text{C}_1\text{-C}_6\text{-alkyl-}$, heteroaryl- $\text{C}_2\text{-C}_6\text{-alkenyl-}$ or heteroaryl- $\text{C}_2\text{-C}_6\text{-alkynyl-}$, wherein the cyclic moieties are optionally substituted with one or more substituents selected from R^{37} ,
 15 • hydrogen,

R^{31} is independently selected from hydrogen, halogen, $-\text{CN}$, $-\text{CH}_2\text{CN}$, $-\text{CHF}_2$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OCHF}_2$, $-\text{OCH}_2\text{CF}_3$, $-\text{OCF}_2\text{CHF}_2$, $-\text{S}(\text{O})_2\text{CF}_3$, $-\text{SCF}_3$, $-\text{NO}_2$, $-\text{OR}^{35}$, $-\text{C}(\text{O})\text{R}^{35}$, $-\text{NR}^{35}\text{R}^{36}$, $-\text{SR}^{35}$,
 20 $-\text{NR}^{35}\text{S}(\text{O})_2\text{R}^{36}$, $-\text{S}(\text{O})_2\text{NR}^{35}\text{R}^{36}$, $-\text{S}(\text{O})\text{NR}^{35}\text{R}^{36}$, $-\text{S}(\text{O})\text{R}^{35}$, $-\text{S}(\text{O})_2\text{R}^{35}$, $-\text{C}(\text{O})\text{NR}^{35}\text{R}^{36}$, $-\text{OC}(\text{O})\text{NR}^{35}\text{R}^{36}$, $-\text{NR}^{35}\text{C}(\text{O})\text{R}^{36}$, $-\text{CH}_2\text{C}(\text{O})\text{NR}^{35}\text{R}^{36}$, $-\text{OCH}_2\text{C}(\text{O})\text{NR}^{35}\text{R}^{36}$, $-\text{CH}_2\text{OR}^{35}$, $-\text{CH}_2\text{NR}^{35}\text{R}^{36}$, $-\text{OC}(\text{O})\text{R}^{35}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{35}$, $-\text{SC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{35}$, $-\text{C}_2\text{-C}_6\text{-alkenyl-C}(\text{O})\text{OR}^{35}$, $-\text{NR}^{35}\text{-C}(\text{O})\text{-C}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{35}$, $-\text{NR}^{35}\text{-C}(\text{O})\text{-C}_1\text{-C}_6\text{-alkenyl-C}(\text{O})\text{OR}^{35}$, $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_1\text{-C}_6\text{-alkanoyl}$ or $-\text{C}(\text{O})\text{OR}^{35}$,

25

R^{32} and R^{33} are independently selected from hydrogen, $\text{C}_1\text{-C}_6\text{-alkyl}$ or $\text{C}_1\text{-C}_6\text{-alkanoyl}$,

R^{34} is independently selected from halogen, $-\text{CN}$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OR}^{35}$, and $-\text{NR}^{35}\text{R}^{36}$,

- 30 R^{35} and R^{36} are independently selected from hydrogen, $\text{C}_1\text{-C}_6\text{-alkyl}$, aryl- $\text{C}_1\text{-C}_6\text{-alkyl}$ or aryl, or R^{35} and R^{36} when attached to the same nitrogen atom together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further

heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds,


- 5 R^{37} is independently selected from halogen, $-C(O)OR^{35}$, $-C(O)H$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OR^{35}$, $-NR^{35}R^{36}$, C_1 - C_6 -alkyl or C_1 - C_6 -alkanoyl,

or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base,


With the proviso that R^{31} and J cannot both be hydrogen.

10


In another embodiment the invention provides a pharmaceutical composition wherein H is



In another embodiment the invention provides a pharmaceutical composition wherein H is



In another embodiment the invention provides a pharmaceutical composition wherein H is



15

In another embodiment the invention provides a pharmaceutical composition wherein I is a valence bond, $-CH_2N(R^{32})-$, or $-SO_2N(R^{33})-$.

In another embodiment the invention provides a pharmaceutical composition wherein I is a valence bond.

20

In another embodiment the invention provides a pharmaceutical composition wherein J is

- hydrogen,
- C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl or C_2 - C_6 -alkynyl,

which may optionally be substituted with one or more substituents selected from halogen, $-CN$, $-CF_3$, $-OCF_3$, $-OR^{35}$, and $-NR^{35}R^{36}$,

- aryl, or heteroaryl, wherein the cyclic moieties are optionally substituted with one or more substituents independently selected from R³⁷.

In another embodiment the invention provides a pharmaceutical composition wherein J is

- hydrogen,

- 5 • aryl or heteroaryl, wherein the cyclic moieties are optionally substituted with one or more substituents independently selected from R³⁷.

In another embodiment the invention provides a pharmaceutical composition wherein J is

- hydrogen,

- 10 • ArG1 or Het3, wherein the cyclic moieties are optionally substituted with one or more substituents independently selected from R³⁷.

In another embodiment the invention provides a pharmaceutical composition wherein J is

- hydrogen,
- phenyl or naphthyl optionally substituted with one or more substituents independently selected from R³⁷.

- 15 In another embodiment the invention provides a pharmaceutical composition wherein J is hydrogen.

In another embodiment the invention provides a pharmaceutical composition wherein R³² and R³³ are independently selected from hydrogen or C₁-C₆-alkyl.

- 20 In another embodiment the invention provides a pharmaceutical composition wherein R³⁴ is hydrogen, halogen, -CN, -CF₃, -OCF₃, -SCF₃, -NO₂, -OR³⁵, -C(O)R³⁵, -NR³⁵R³⁶, -SR³⁵, -C(O)NR³⁵R³⁶, -OC(O)NR³⁵R³⁶, -NR³⁵C(O)R³⁶, -OC(O)R³⁵, -OC₁-C₆-alkyl-C(O)OR³⁵, -SC₁-C₆-alkyl-C(O)OR³⁵ or -C(O)OR³⁵.

In another embodiment the invention provides a pharmaceutical composition wherein R³⁴ is hydrogen, halogen, -CF₃, -NO₂, -OR³⁵, -NR³⁵R³⁶, -SR³⁵, -NR³⁵C(O)R³⁶, or -C(O)OR³⁵.

- 25 In another embodiment the invention provides a pharmaceutical composition wherein R³⁴ is hydrogen, halogen, -CF₃, -NO₂, -OR³⁵, -NR³⁵R³⁶, or -NR³⁵C(O)R³⁶.

In another embodiment the invention provides a pharmaceutical composition wherein R³⁴ is hydrogen, halogen, or -OR³⁵.

- 30 In another embodiment the invention provides a pharmaceutical composition wherein R³⁵ and R³⁶ are independently selected from hydrogen, C₁-C₆-alkyl, or aryl.


In another embodiment the invention provides a pharmaceutical composition wherein R³⁵ and R³⁶ are independently selected from hydrogen or C₁-C₆-alkyl.

In another embodiment the invention provides a pharmaceutical composition wherein R³⁷ is halogen, -C(O)OR³⁵, -CN, -CF₃, -OR³⁵, -NR³⁵R³⁶, C₁-C₆-alkyl or C₁-C₆-alkanoyl.

In another embodiment the invention provides a pharmaceutical composition wherein R^{37} is halogen, $-C(O)OR^{35}$, $-OR^{35}$, $-NR^{35}R^{36}$, C_1 - C_6 -alkyl or C_1 - C_6 -alkanoyl.

In another embodiment the invention provides a pharmaceutical composition wherein R^{37} is halogen, $-C(O)OR^{35}$ or $-OR^{35}$.

- 5 In another embodiment the invention provides a pharmaceutical composition wherein the zinc-binding ligand is



- 10 wherein K is a valence bond, C_1 - C_6 -alkylene, $-NH-C(=O)-U-$, $-C_1$ - C_6 -alkyl-S-, $-C_1$ - C_6 -alkyl-O-, $-C(=O)-$, or $-C(=O)-NH-$, wherein any C_1 - C_6 -alkyl moiety is optionally substituted with R^{38} ,

U is a valence bond, C_1 - C_6 -alkenylene, $-C_1$ - C_6 -alkyl-O- or C_1 - C_6 -alkylene wherein any C_1 - C_6 -alkyl moiety is optionally substituted with C_1 - C_6 -alkyl,

- 15 R^{38} is C_1 - C_6 -alkyl, aryl, wherein the alkyl or aryl moieties are optionally substituted with one or more substituents independently selected from R^{39} ,

R^{39} is independently selected from halogen, cyano, nitro, amino,

- 20 M is a valence bond, arylene or heteroarylene, wherein the aryl or heteroaryl moieties are optionally substituted with one or more substituents independently selected from R^{40} ,

R^{40} is selected from

- 25 • hydrogen, halogen, $-CN$, $-CH_2CN$, $-CHF_2$, $-CF_3$, $-OCF_3$, $-OCHF_2$, $-OCH_2CF_3$, $-OCF_2CHF_2$, $-S(O)_2CF_3$, $-OS(O)_2CF_3$, $-SCF_3$, $-NO_2$, $-OR^{41}$, $-NR^{41}R^{42}$, $-SR^{41}$, $-NR^{41}S(O)_2R^{42}$, $-S(O)_2NR^{41}R^{42}$, $-S(O)NR^{41}R^{42}$, $-S(O)R^{41}$, $-S(O)_2R^{41}$, $-OS(O)_2R^{41}$, $-C(O)NR^{41}R^{42}$, $-OC(O)NR^{41}R^{42}$, $-NR^{41}C(O)R^{42}$, $-CH_2C(O)NR^{41}R^{42}$, $-OC_1$ - C_6 -alkyl- $C(O)NR^{41}R^{42}$, $-CH_2OR^{41}$, $-CH_2OC(O)R^{41}$, $-CH_2NR^{41}R^{42}$, $-OC(O)R^{41}$, $-OC_1$ - C_6 -alkyl- $C(O)OR^{41}$, $-OC_1$ - C_6 -alkyl- OR^{41} , $-S-C_1$ - C_6 -alkyl- $C(O)OR^{41}$, $-C_2$ - C_6 -alkenyl- $C(=O)OR^{41}$, $-NR^{41}-C(=O)-C_1$ - C_6 -alkyl- $C(=O)OR^{41}$, $-NR^{41}-C(=O)-C_1$ - C_6 -alkenyl- $C(=O)OR^{41}$, $-C(O)OR^{41}$, $-C_2$ - C_6 -alkenyl- $C(=O)R^{41}$, $=O$, $-NH-C(=O)-O-C_1$ - C_6 -alkyl, or $-NH-C(=O)-C(=O)-O-C_1$ - C_6 -alkyl,
- 30

• C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl, which may each optionally be substituted with one or more substituents selected from R⁴³,

• aryl, aryloxy, aryloxy carbonyl, aroyl, arylsulfanyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aroyl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl or heteroaryl-C₂-C₆-alkynyl, wherein the cyclic moieties optionally may be substituted with one or more substituents selected from R⁴⁴,

R⁴¹ and R⁴² are independently selected from hydrogen, -OH, C₁-C₆-alkyl, C₁-C₆-alkenyl, aryl-C₁-C₆-alkyl or aryl, wherein the alkyl moieties may optionally be substituted with one or more substituents independently selected from R⁴⁵, and the aryl moieties may optionally be substituted with one or more substituents independently selected from R⁴⁶; R⁴¹ and R⁴² when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom, the heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds,

R⁴³ is independently selected from halogen, -CN, -CF₃, -OCF₃, -OR⁴¹, and -NR⁴¹R⁴²

R⁴⁴ is independently selected from halogen, -C(O)OR⁴¹, -CH₂C(O)OR⁴¹, -CH₂OR⁴¹, -CN, -CF₃, -OCF₃, -NO₂, -OR⁴¹, -NR⁴¹R⁴² and C₁-C₆-alkyl,

R⁴⁵ is independently selected from halogen, -CN, -CF₃, -OCF₃, -O-C₁-C₆-alkyl, -C(O)-O-C₁-C₆-alkyl, -COOH and -NH₂,

R⁴⁶ is independently selected from halogen, -C(O)OC₁-C₆-alkyl, -COOH, -CN, -CF₃, -OCF₃, -NO₂, -OH, -OC₁-C₆-alkyl, -NH₂, C(=O) or C₁-C₆-alkyl,

Q is a valence bond, C₁-C₆-alkylene, -C₁-C₆-alkyl-O-, -C₁-C₆-alkyl-NH-, -NH-C₁-C₆-alkyl, -NH-C(=O)-, -C(=O)-NH-, -O-C₁-C₆-alkyl, -C(=O)-, or -C₁-C₆-alkyl-C(=O)-N(R⁴⁷)- wherein the alkyl moieties are optionally substituted with one or more substituents independently selected from R⁴⁸,

R⁴⁷ and R⁴⁸ are independently selected from hydrogen, C₁-C₆-alkyl, aryl optionally substituted with one or more R⁴⁹,

R⁴⁹ is independently selected from halogen and -COOH,

T is

- 5 • hydrogen,
 • C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₆-alkyloxy-carbonyl, wherein the alkyl, alkenyl and alkynyl moieties are optionally substituted with one or more substituents independently selected from R⁵⁰,
 10 • aryl, aryloxy, aryloxy-carbonyl, aryl-C₁-C₆-alkyl, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl-, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl, heteroaryl-C₂-C₆-alkynyl,

wherein any alkyl, alkenyl, alkynyl, aryl and heteroaryl moiety is optionally substituted with one or more substituents independently selected from R⁵⁰,

15

R⁵⁰ is C₁-C₆-alkyl, C₁-C₆-alkoxy, aryl, aryloxy, aryl-C₁-C₆-alkoxy, -C(=O)-NH-C₁-C₆-alkyl-aryl, heteroaryl, heteroaryl-C₁-C₆-alkoxy, -C₁-C₆-alkyl-COOH, -O-C₁-C₆-alkyl-COOH, -S(O)₂R⁵¹, -C₂-C₆-alkenyl-COOH, -OR⁵¹, -NO₂, halogen, -COOH, -CF₃, -CN, =O, -N(R⁵¹R⁵²), wherein the aryl or heteroaryl moieties are optionally substituted with one or more R⁵³,

20

R⁵¹ and R⁵² are independently selected from hydrogen and C₁-C₆-alkyl,
 R⁵³ is independently selected from C₁-C₆-alkyl, C₁-C₆-alkoxy, -C₁-C₆-alkyl-COOH, -C₂-C₆-alkenyl-COOH, -OR⁵¹, -NO₂, halogen, -COOH, -CF₃, -CN, or -N(R⁵¹R⁵²),

- 25 or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.

In another embodiment the invention provides a pharmaceutical composition wherein K is a valence bond, C₁-C₆-alkylene, -NH-C(=O)-U-, -C₁-C₆-alkyl-S-, -C₁-C₆-alkyl-O-, or -C(=O)-, wherein any C₁-C₆-alkyl moiety is optionally substituted with R³⁸.

- 30 In another embodiment the invention provides a pharmaceutical composition wherein K is a valence bond, C₁-C₆-alkylene, -NH-C(=O)-U-, -C₁-C₆-alkyl-S-, or -C₁-C₆-alkyl-O-, wherein any C₁-C₆-alkyl moiety is optionally substituted with R³⁸.

In another embodiment the invention provides a pharmaceutical composition wherein K is a valence bond, C₁-C₆-alkylene, or -NH-C(=O)-U-, wherein any C₁-C₆-alkyl moiety is optionally

- 35 substituted with R³⁸.

In another embodiment the invention provides a pharmaceutical composition wherein K is a valence bond or C₁-C₆-alkylene, wherein any C₁-C₆-alkyl moiety is optionally substituted with R³⁸.

5 In another embodiment the invention provides a pharmaceutical composition wherein K is a valence bond or -NH-C(=O)-U.

In another embodiment the invention provides a pharmaceutical composition wherein K is a valence bond.

In another embodiment the invention provides a pharmaceutical composition wherein U is a valence bond or -C₁-C₆-alkyl-O-.

10 In another embodiment the invention provides a pharmaceutical composition wherein U is a valence bond.

In another embodiment the invention provides a pharmaceutical composition wherein M is arylene or heteroarylene, wherein the arylene or heteroarylene moieties are optionally substituted with one or more substituents independently selected from R⁴⁰.

15 In another embodiment the invention provides a pharmaceutical composition wherein M is ArG1 or Het1, wherein the arylene or heteroarylene moieties are optionally substituted with one or more substituents independently selected from R⁴⁰.


In another embodiment the invention provides a pharmaceutical composition wherein M is ArG1 or Het2, wherein the arylene or heteroarylene moieties are optionally substituted with
20 one or more substituents independently selected from R⁴⁰.

In another embodiment the invention provides a pharmaceutical composition wherein M is ArG1 or Het3, wherein the arylene or heteroarylene moieties are optionally substituted with one or more substituents independently selected from R⁴⁰.

In another embodiment the invention provides a pharmaceutical composition wherein M is
25 phenylene optionally substituted with one or more substituents independently selected from R⁴⁰.


In another embodiment the invention provides a pharmaceutical composition wherein M is indolylene optionally substituted with one or more substituents independently selected from R⁴⁰.

30 In another embodiment the invention provides a pharmaceutical composition wherein M is



In another embodiment the invention provides a pharmaceutical composition wherein M is carbazolyne optionally substituted with one or more substituents independently selected from R⁴⁰.

In another embodiment the invention provides a pharmaceutical composition wherein M is



5

In another embodiment the invention provides a pharmaceutical composition wherein R⁴⁰ is selected from

- hydrogen, halogen, -CN, -CF₃, -OCF₃, -NO₂, -OR⁴¹, -NR⁴¹R⁴², -SR⁴¹, -S(O)₂R⁴¹, -NR⁴¹C(O)R⁴², -OC₁-C₆-alkyl-C(O)NR⁴¹R⁴², -C₂-C₆-alkenyl-C(=O)OR⁴¹, -C(O)OR⁴¹, =O, -NH-C(=O)-O-C₁-C₆-alkyl, or -NH-C(=O)-C(=O)-O-C₁-C₆-alkyl,

10

C₁-C₆-alkyl or C₂-C₆- alkenyl which may each optionally be substituted with one or more substituents independently selected from R⁴³,

15

- aryl, aryloxy, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, or heteroaryl-C₂-C₆-alkenyl, wherein the cyclic moieties optionally may be substituted with one or more substituents selected from R⁴⁴.

In another embodiment the invention provides a pharmaceutical composition wherein R⁴⁰ is selected from

20

- hydrogen, halogen, -CN, -CF₃, -OCF₃, -NO₂, -OR⁴¹, -NR⁴¹R⁴², -SR⁴¹, -S(O)₂R⁴¹, -NR⁴¹C(O)R⁴², -OC₁-C₆-alkyl-C(O)NR⁴¹R⁴², -C₂-C₆-alkenyl-C(=O)OR⁴¹, -C(O)OR⁴¹, =O, -NH-C(=O)-O-C₁-C₆-alkyl, or -NH-C(=O)-C(=O)-O-C₁-C₆-alkyl,

25

C₁-C₆-alkyl or C₂-C₆- alkenyl which may each optionally be substituted with one or more substituents independently selected from R⁴³,

- ArG1, ArG1-O-, ArG1-C₁-C₆-alkoxy, ArG1-C₁-C₆-alkyl, ArG1-C₂-C₆-alkenyl, Het3, Het3-C₁-C₆-alkyl, or Het3-C₂-C₆-alkenyl, wherein the cyclic moieties optionally may be substituted with one or more substituents selected from R⁴⁴.

30

In another embodiment the invention provides a pharmaceutical composition wherein R⁴⁰ is selected from

- hydrogen, halogen, -CF₃, -NO₂, -OR⁴¹, -NR⁴¹R⁴², -C(O)OR⁴¹, =O, or -NR⁴¹C(O)R⁴²,

- C₁-C₆-alkyl,
- ArG1.

In another embodiment the invention provides a pharmaceutical composition wherein R⁴⁰ is selected from

- 5 • Halogen, -NO₂, -OR⁴¹, -NR⁴¹R⁴², -C(O)OR⁴¹, or -NR⁴¹C(O)R⁴²,
 • Methyl,
 • Phenyl.

10 In another embodiment the invention provides a pharmaceutical composition wherein R⁴¹ and R⁴² are independently selected from hydrogen, C₁-C₆-alkyl, or aryl, wherein the aryl moieties may optionally be substituted with halogen or -COOH.

In another embodiment the invention provides a pharmaceutical composition wherein R⁴¹ and R⁴² are independently selected from hydrogen, methyl, ethyl, or phenyl, wherein the phenyl moieties may optionally be substituted with halogen or -COOH.

15 In another embodiment the invention provides a pharmaceutical composition wherein Q is a valence bond, C₁-C₆-alkylene, -C₁-C₆-alkyl-O-, -C₁-C₆-alkyl-NH-, -NH-C₁-C₆-alkyl, -NH-C(=O)-, -C(=O)-NH-, -O-C₁-C₆-alkyl, -C(=O)-, or -C₁-C₆-alkyl-C(=O)-N(R⁴⁷)- wherein the alkyl moieties are optionally substituted with one or more substituents independently selected from R⁴⁸.

20 In another embodiment the invention provides a pharmaceutical composition wherein Q is a valence bond, -CH₂-, -CH₂-CH₂-, -CH₂-O-, -CH₂-CH₂-O-, -CH₂-NH-, -CH₂-CH₂-NH-, -NH-CH₂-, -NH-CH₂-CH₂-, -NH-C(=O)-, -C(=O)-NH-, -O-CH₂-, -O-CH₂-CH₂-, or -C(=O)-.

In another embodiment the invention provides a pharmaceutical composition wherein R⁴⁷ and R⁴⁸ are independently selected from hydrogen, methyl and phenyl.

In another embodiment the invention provides a pharmaceutical composition wherein T is

- 25 • hydrogen,
 • C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R⁵⁰,
 • aryl, aryl-C₁-C₆-alkyl, heteroaryl, wherein the alkyl, aryl and heteroaryl moieties are optionally substituted with one or more substituents independently selected from R⁵⁰.

30 In another embodiment the invention provides a pharmaceutical composition wherein T is

- hydrogen,
 • C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R⁵⁰,
 • ArG1, ArG1-C₁-C₆-alkyl, Het3, wherein the alkyl, aryl and heteroaryl moieties are
35 optionally substituted with one or more substituents independently selected from R⁵⁰.

In another embodiment the invention provides a pharmaceutical composition wherein T is

- hydrogen,
- C₁-C₆-alkyl, optionally substituted with one or more substituents independently selected from R⁵⁰,
- phenyl, phenyl-C₁-C₆-alkyl, wherein the alkyl and phenyl moieties are optionally substituted with one or more substituents independently selected from R⁵⁰.

10 In another embodiment the invention provides a pharmaceutical composition wherein R⁵⁰ is C₁-C₆-alkyl, C₁-C₆-alkoxy, aryl, aryloxy, aryl-C₁-C₆-alkoxy, -C(=O)-NH-C₁-C₆-alkyl-aryl, heteroaryl, -C₁-C₆-alkyl-COOH, -O-C₁-C₆-alkyl-COOH, -S(O)₂R⁵¹, -C₂-C₆-alkenyl-COOH, -OR⁵¹, -NO₂, halogen, -COOH, -CF₃, -CN, =O, -N(R⁵¹R⁵²), wherein the aryl or heteroaryl moieties are optionally substituted with one or more R⁵³.

15 In another embodiment the invention provides a pharmaceutical composition wherein R⁵⁰ is C₁-C₆-alkyl, C₁-C₆-alkoxy, aryl, aryloxy, aryl-C₁-C₆-alkoxy, -OR⁵¹, -NO₂, halogen, -COOH, -CF₃, wherein any aryl moiety is optionally substituted with one or more R⁵³.

In another embodiment the invention provides a pharmaceutical composition wherein R⁵⁰ is C₁-C₆-alkyl, aryloxy, aryl-C₁-C₆-alkoxy, -OR⁵¹, halogen, -COOH, -CF₃, wherein any aryl moiety is optionally substituted with one or more R⁵³.

20 In another embodiment the invention provides a pharmaceutical composition wherein R⁵⁰ is C₁-C₆-alkyl, ArG1-O-, ArG1-C₁-C₆-alkoxy, -OR⁵¹, halogen, -COOH, -CF₃, wherein any aryl moiety is optionally substituted with one or more R⁵³.


In another embodiment the invention provides a pharmaceutical composition wherein R⁵⁰ is phenyl, methyl or ethyl.

25 In another embodiment the invention provides a pharmaceutical composition wherein R⁵⁰ is methyl or ethyl.

In another embodiment the invention provides a pharmaceutical composition wherein R⁵¹ is methyl.

30 In another embodiment the invention provides a pharmaceutical composition wherein R⁵³ is C₁-C₆-alkyl, C₁-C₆-alkoxy, -OR⁵¹, halogen, or -CF₃.

In another embodiment the invention provides a pharmaceutical composition wherein the zinc-binding ligand is



wherein V is C₁-C₆-alkyl, aryl, heteroaryl, aryl-C₁₋₆-alkyl- or aryl-C₂₋₆-alkenyl-, wherein the alkyl or alkenyl is optionally substituted with one or more substituents independently selected from R⁵⁴, and the aryl or heteroaryl is optionally substituted with one or more substituents independently selected from R⁵⁵,

R⁵⁴ is independently selected from halogen, -CN, -CF₃, -OCF₃, aryl, -COOH and -NH₂,

R⁵⁵ is independently selected from

• hydrogen, halogen, -CN, -CH₂CN, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -S(O)₂CF₃, -OS(O)₂CF₃, -SCF₃, -NO₂, -OR⁵⁶, -NR⁵⁶R⁵⁷, -SR⁵⁶, -NR⁵⁶S(O)₂R⁵⁷, -S(O)₂NR⁵⁶R⁵⁷, -S(O)NR⁵⁶R⁵⁷, -S(O)R⁵⁶, -S(O)₂R⁵⁶, -OS(O)₂R⁵⁶, -C(O)NR⁵⁶R⁵⁷, -OC(O)NR⁵⁶R⁵⁷, -NR⁵⁶C(O)R⁵⁷, -CH₂C(O)NR⁵⁶R⁵⁷, -OC₁-C₆-alkyl-C(O)NR⁵⁶R⁵⁷, -CH₂OR⁵⁶, -CH₂OC(O)R⁵⁶, -CH₂NR⁵⁶R⁵⁷, -OC(O)R⁵⁶, -OC₁-C₆-alkyl-C(O)OR⁵⁶, -OC₁-C₆-alkyl-OR⁵⁶, -SC₁-C₆-alkyl-C(O)OR⁵⁶, -C₂-C₆-alkenyl-C(=O)OR⁵⁶, -NR⁵⁶-C(=O)-C₁-C₆-alkyl-C(=O)OR⁵⁶, -NR⁵⁶-C(=O)-C₁-C₆-alkenyl-C(=O)OR⁵⁶, -C(O)OR⁵⁶, or -C₂-C₆-alkenyl-C(=O)R⁵⁶,

• C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl,

which may optionally be substituted with one or more substituents selected from R⁵⁸,

• aryl, aryloxy, aryloxycarbonyl, aroyl, arylsulfanyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aroyl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl or heteroaryl-C₂-C₆-alkynyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R⁵⁹,

R⁵⁶ and R⁵⁷ are independently selected from hydrogen, OH, CF₃, C₁-C₁₂-alkyl, aryl-C₁-C₆-alkyl, -C(=O)-C₁-C₆-alkyl or aryl, wherein the alkyl groups may optionally be substituted with one or more substituents independently selected from R⁶⁰, and the aryl groups may optionally be substituted with one or more substituents independently selected from R⁶¹; R⁵⁸

and R⁵⁷ when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom, the heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds,

5

R⁵⁸ is independently selected from halogen, -CN, -CF₃, -OCF₃, -OR⁵⁶, and -NR⁵⁶R⁵⁷,

R⁵⁹ is independently selected from halogen, -C(O)OR⁵⁶, -CH₂C(O)OR⁵⁶, -CH₂OR⁵⁶, -CN, -CF₃, -OCF₃, -NO₂, -OR⁵⁶, -NR⁵⁶R⁵⁷ and C₁-C₆-alkyl,

10

R⁶⁰ is independently selected from halogen, -CN, -CF₃, -OCF₃, -OC₁-C₆-alkyl, -C(O)OC₁-C₆-alkyl, -C(=O)-R⁶², -COOH and -NH₂,

R⁶¹ is independently selected from halogen, -C(O)OC₁-C₆-alkyl, -COOH, -CN, -CF₃, -OCF₃, -NO₂, -OH, -OC₁-C₆-alkyl, -NH₂, C(=O) or C₁-C₆-alkyl,

15

R⁶² is C₁-C₆-alkyl, aryl optionally substituted with one or more substituents independently selected from halogen, or heteroaryl optionally substituted with one or more C₁-C₆-alkyl independently,

20

or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.

In another embodiment the invention provides a pharmaceutical composition wherein V is aryl, heteroaryl, or aryl-C₁₋₆-alkyl-, wherein the alkyl is optionally substituted with one or more substituents independently selected R⁵⁴, and the aryl or heteroaryl is optionally substituted with one or more substituents independently selected from R⁵⁵.

25

In another embodiment the invention provides a pharmaceutical composition wherein V is aryl, Het1, or aryl-C₁₋₆-alkyl-, wherein the alkyl is optionally substituted with one or more substituents independently selected from R⁵⁴, and the aryl or heteroaryl moiety is optionally substituted with one or more substituents independently selected from R⁵⁵.

30

In another embodiment the invention provides a pharmaceutical composition wherein V is aryl, Het2, or aryl-C₁₋₆-alkyl-, wherein the alkyl is optionally substituted with one or more substituents independently selected from R⁵⁴, and the aryl or heteroaryl moiety is optionally substituted with one or more substituents independently selected from R⁵⁵.

In another embodiment the invention provides a pharmaceutical composition wherein V is aryl, Het3, or aryl-C₁₋₆-alkyl-, wherein the alkyl is optionally substituted with one or more substituents independently selected from R⁵⁴, and the aryl or heteroaryl moiety is optionally substituted with one or more substituents independently selected from R⁵⁵.

5 In another embodiment the invention provides a pharmaceutical composition wherein V is aryl optionally substituted with one or more substituents independently selected from R⁵⁵.

In another embodiment the invention provides a pharmaceutical composition wherein V is ArG1 optionally substituted with one or more substituents independently selected from R⁵⁵.

10 In another embodiment the invention provides a pharmaceutical composition wherein V is phenyl, naphthyl or anthranyl optionally substituted with one or more substituents independently selected from R⁵⁵.

In another embodiment the invention provides a pharmaceutical composition wherein V is phenyl optionally substituted with one or more substituents independently selected from R⁵⁵.

15 In another embodiment the invention provides a pharmaceutical composition wherein R⁵⁵ is independently selected from

- halogen, C₁-C₆-alkyl, -CN, -OCF₃, -CF₃, -NO₂, -OR⁵⁶, -NR⁵⁶R⁵⁷, -NR⁵⁶C(O)R⁵⁷-SR⁵⁶, -OC₁-C₈-alkyl-C(O)OR⁵⁶, or -C(O)OR⁵⁶,

- C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R⁵⁸

20 • aryl, aryl-C₁-C₆-alkyl, heteroaryl, or heteroaryl-C₁-C₆-alkyl
of which the cyclic moieties optionally may be substituted with one or more substituents independently selected from R⁵⁹.

In another embodiment the invention provides a pharmaceutical composition wherein R⁵⁵ is independently selected from

25 • halogen, C₁-C₆-alkyl, -CN, -OCF₃, -CF₃, -NO₂, -OR⁵⁶, -NR⁵⁶R⁵⁷, -NR⁵⁶C(O)R⁵⁷-SR⁵⁶, -OC₁-C₈-alkyl-C(O)OR⁵⁶, or -C(O)OR⁵⁶

- C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R⁵⁸

- ArG1, ArG1-C₁-C₆-alkyl, Het3, or Het3-C₁-C₆-alkyl

30 of which the cyclic moieties optionally may be substituted with one or more substituents independently selected from R⁵⁹.

In another embodiment the invention provides a pharmaceutical composition wherein R⁵⁵ is independently selected from halogen, -OR⁵⁶, -NR⁵⁶R⁵⁷, -C(O)OR⁵⁶, -OC₁-C₈-alkyl-C(O)OR⁵⁶,

35 -NR⁵⁶C(O)R⁵⁷ or C₁-C₆-alkyl.


In another embodiment the invention provides a pharmaceutical composition wherein R^{55} is independently selected from halogen, $-OR^{56}$, $-NR^{56}R^{57}$, $-C(O)OR^{56}$, $-OC_1-C_8\text{-alkyl}-C(O)OR^{56}$, $-NR^{56}C(O)R^{57}$, methyl or ethyl.

In another embodiment the invention provides a pharmaceutical composition wherein R^{56} and R^{57} are independently selected from hydrogen, CF_3 , $C_1-C_{12}\text{-alkyl}$, or $-C(=O)-C_1-C_8\text{-alkyl}$; R^{56} and R^{57} when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom.

In another embodiment the invention provides a pharmaceutical composition wherein R^{56} and R^{57} are independently selected from hydrogen or $C_1-C_{12}\text{-alkyl}$, R^{56} and R^{57} when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom.

In another embodiment the invention provides a pharmaceutical composition wherein R^{56} and R^{57} are independently selected from hydrogen or methyl, ethyl, propyl butyl, R^{56} and R^{57} when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom.

In another embodiment the invention provides a pharmaceutical composition 1 wherein the zinc-binding ligand is



wherein AA is $C_1-C_6\text{-alkyl}$, aryl, heteroaryl, aryl- $C_{1-6}\text{-alkyl}$ - or aryl- $C_{2-6}\text{-alkenyl}$ -, wherein the alkyl or alkenyl is optionally substituted with one or more substituents independently selected from R^{63} , and the aryl or heteroaryl is optionally substituted with one or more substituents independently selected from R^{64} ,

R^{63} is independently selected from halogen, $-CN$, $-CF_3$, $-OCF_3$, aryl, $-COOH$ and $-NH_2$,

R^{64} is independently selected from

- hydrogen, halogen, $-CN$, $-CH_2CN$, $-CHF_2$, $-CF_3$, $-OCF_3$, $-OCHF_2$, $-OCH_2CF_3$, $-OCF_2CHF_2$, $-S(O)_2CF_3$, $-OS(O)_2CF_3$, $-SCF_3$, $-NO_2$, $-OR^{65}$, $-NR^{65}R^{66}$, $-SR^{65}$, $-NR^{65}S(O)_2R^{66}$, $-S(O)_2NR^{65}R^{66}$, $-S(O)NR^{65}R^{66}$, $-S(O)R^{65}$, $-S(O)_2R^{65}$, $-OS(O)_2R^{65}$,

-C(O)NR⁶⁵R⁶⁶, -OC(O)NR⁶⁵R⁶⁶, -NR⁶⁵C(O)R⁶⁶, -CH₂C(O)NR⁶⁵R⁶⁶, -OC₁-C₆-alkyl-C(O)NR⁶⁵R⁶⁶, -CH₂OR⁶⁵, -CH₂OC(O)R⁶⁵, -CH₂NR⁶⁵R⁶⁶, -OC(O)R⁶⁵, -OC₁-C₆-alkyl-C(O)OR⁶⁵, -OC₁-C₆-alkyl-OR⁶⁵, -SC₁-C₆-alkyl-C(O)OR⁶⁵, -C₂-C₆-alkenyl-C(=O)OR⁶⁵, -NR⁶⁵-C(=O)-C₁-C₆-alkyl-C(=O)OR⁶⁵, -NR⁶⁵-C(=O)-C₁-C₆-alkenyl-C(=O)OR⁶⁵, -C(O)OR⁶⁵, or -C₂-C₆-alkenyl-C(=O)R⁶⁵,

• C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl, each of which may optionally be substituted with one or more substituents selected from R⁶⁷,

• aryl, aryloxy, aryloxycarbonyl, aroyl, arylsulfanyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aroyl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl or heteroaryl-C₂-C₆-alkynyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R⁶⁸,

R⁶⁵ and R⁶⁶ are independently selected from hydrogen, OH, CF₃, C₁-C₁₂-alkyl, aryl-C₁-C₆-alkyl, -C(=O)-R⁶⁹, aryl or heteroaryl, wherein the alkyl groups may optionally be substituted with one or more substituents selected from R⁷⁰, and the aryl and heteroaryl groups may optionally be substituted with one or more substituents independently selected from R⁷¹; R⁶⁵ and R⁶⁶ when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom, the heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds,

R⁶⁷ is independently selected from halogen, -CN, -CF₃, -OCF₃, -OR⁶⁵, and -NR⁶⁵R⁶⁶,

R⁶⁸ is independently selected from halogen, -C(O)OR⁶⁵, -CH₂C(O)OR⁶⁵, -CH₂OR⁶⁵, -CN, -CF₃, -OCF₃, -NO₂, -OR⁶⁵, -NR⁶⁵R⁶⁶ and C₁-C₆-alkyl,

R⁶⁹ is independently selected from C₁-C₆-alkyl, aryl optionally substituted with one or more halogen, or heteroaryl optionally substituted with one or more C₁-C₆-alkyl,

R⁷⁰ is independently selected from halogen, -CN, -CF₃, -OCF₃, -OC₁-C₆-alkyl, -C(O)OC₁-C₆-alkyl, -COOH and -NH₂,

R⁷¹ is independently selected from halogen, -C(O)OC₁₋₆-alkyl, -COOH, -CN, -CF₃, -OCF₃, -NO₂, -OH, -OC₁₋₆-alkyl, -NH₂, C(=O) or C₁₋₆-alkyl,

- 5 or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.

In another embodiment the invention provides a pharmaceutical composition wherein AA is aryl, heteroaryl or aryl-C₁₋₆-alkyl-, wherein the alkyl is optionally substituted with one or more R⁶³, and the aryl or heteroaryl is optionally substituted with one or more substituents

- 10 independently selected from R⁶⁴.

In another embodiment the invention provides a pharmaceutical composition wherein AA is aryl or heteroaryl optionally substituted with one or more substituents independently selected from R⁶⁴.

- 15 In another embodiment the invention provides a pharmaceutical composition wherein AA is ArG1 or Het1 optionally substituted with one or more substituents independently selected from R⁶⁴.

In another embodiment the invention provides a pharmaceutical composition wherein AA is ArG1 or Het2 optionally substituted with one or more substituents independently selected from R⁶⁴.

- 20 In another embodiment the invention provides a pharmaceutical composition wherein AA is ArG1 or Het3 optionally substituted with one or more substituents independently selected from R⁶⁴.

- 25 In another embodiment the invention provides a pharmaceutical composition wherein AA is phenyl, naphthyl, anthryl, carbazolyl, thienyl, pyridyl, or benzodioxyl optionally substituted with one or more substituents independently selected from R⁶⁴.

In another embodiment the invention provides a pharmaceutical composition wherein AA is phenyl or naphthyl optionally substituted with one or more substituents independently selected from R⁶⁴.

- 30 In another embodiment the invention provides a pharmaceutical composition wherein R⁶⁴ is independently selected from hydrogen, halogen, -CF₃, -OCF₃, -OR⁶⁵, -NR⁶⁵R⁶⁶, C₁₋₆-alkyl, -OC(O)R⁶⁵, -OC₁₋₆-alkyl-C(O)OR⁶⁵, aryl-C₂₋₆-alkenyl, aryloxy or aryl, wherein C₁₋₆-alkyl is optionally substituted with one or more substituents independently selected from R⁶⁷, and the cyclic moieties optionally are substituted with one or more substituents independently selected from R⁶⁸.

In another embodiment the invention provides a pharmaceutical composition wherein R^{64} is independently selected from halogen, $-CF_3$, $-OCF_3$, $-OR^{65}$, $-NR^{65}R^{66}$, methyl, ethyl, propyl, $-OC(O)R^{65}$, $-OCH_2-C(O)OR^{65}$, $-OCH_2-CH_2-C(O)OR^{65}$, phenoxy optionally substituted with one or more substituents independently selected from R^{68} .

- 5 In another embodiment the invention provides a pharmaceutical composition wherein R^{65} and R^{66} are independently selected from hydrogen, CF_3 , C_1 - C_{12} -alkyl, aryl, or heteroaryl optionally substituted with one or more substituents independently selected from R^{71} .

In another embodiment the invention provides a pharmaceutical composition wherein R^{65} and R^{66} are independently hydrogen, C_1 - C_{12} -alkyl, aryl, or heteroaryl optionally substituted with one or more substituents independently selected from R^{71} .

In another embodiment the invention provides a pharmaceutical composition wherein R^{65} and R^{66} are independently hydrogen, methyl, ethyl, propyl, butyl, 2,2-dimethyl-propyl, ArG1 or Het1 optionally substituted with one or more substituents independently selected from R^{71} .

- 15 In another embodiment the invention provides a pharmaceutical composition wherein R^{65} and R^{66} are independently hydrogen, methyl, ethyl, propyl, butyl, 2,2-dimethyl-propyl, ArG1 or Het2 optionally substituted with one or more substituents independently selected from R^{71} .

In another embodiment the invention provides a pharmaceutical composition wherein R^{65} and R^{66} are independently hydrogen, methyl, ethyl, propyl, butyl, 2,2-dimethyl-propyl, ArG1 or Het3 optionally substituted with one or more substituents independently selected from R^{71} .


- 20 In another embodiment the invention provides a pharmaceutical composition wherein R^{65} and R^{66} are independently hydrogen, methyl, ethyl, propyl, butyl, 2,2-dimethyl-propyl, phenyl, naphthyl, thiadiazolyl optionally substituted with one or more R^{71} independently; or isoxazolyl optionally substituted with one or more substituents independently selected from R^{71} .

In another embodiment the invention provides a pharmaceutical composition wherein R^{71} is halogen or C_1 - C_6 -alkyl.

In another embodiment the invention provides a pharmaceutical composition wherein R^{71} is halogen or methyl.

- In another aspect the invention provides a pharmaceutical composition comprising
30 insulin and a zinc-binding ligand which reversibly binds to a $His^{B10} Zn^{2+}$ site of an insulin hexamer, wherein the ligand is selected from the group consisting of
benzotriazoles, 3-hydroxy 2-napthoic acids, salicylic acids, tetrazoles, thiazolidinediones, 5-
mercaptotetrazoles, or 4-cyano-1,2,3-triazoles, or any enantiomer, diastereomer, including a
racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid
35 or base.

In one embodiment hereof the zinc-binding ligand is



wherein

5 X is =O, =S or =NH

Y is -S-, -O- or -NH-

R^1 , R^{1A} and R^4 are independently selected from hydrogen or C_1 - C_6 -alkyl,

10 R^2 and R^{2A} are hydrogen or C_1 - C_6 -alkyl or aryl, R^1 and R^2 may optionally be combined to form a double bond, R^{1A} and R^{2A} may optionally be combined to form a double bond, R^3 , R^{3A} and R^5 are independently selected from hydrogen, halogen, aryl optionally substituted with one or more substituents independently selected from R^{16} , C_1 - C_6 -alkyl, or $-C(O)NR^{11}R^{12}$,

15 A, A^1 and B are independently selected from C_1 - C_6 -alkyl, aryl, aryl- C_1 - C_6 -alkyl, $-NR^{11}$ -aryl, aryl- C_2 - C_6 -alkenyl or heteroaryl, wherein the alkyl or alkenyl is optionally substituted with one or more substituents independently selected from R^6 and the aryl or heteroaryl is optionally substituted with up to four substituents R^7 , R^8 , R^9 , and R^{10} ,

20 A and R^3 may be connected through one or two valence bonds, B and R^5 may be connected through one or two valence bonds,

R^6 is independently selected from halogen, -CN, $-CF_3$, $-OCF_3$, aryl, $-COOH$ and $-NH_2$,

R^7 , R^8 , R^9 and R^{10} are independently selected from

25 •hydrogen, halogen, -CN, $-CH_2CN$, $-CHF_2$, $-CF_3$, $-OCF_3$, $-OCHF_2$, $-OCH_2CF_3$, $-OCF_2CHF_2$, $-S(O)_2CF_3$, $-OS(O)_2CF_3$, $-SCF_3$, $-NO_2$, $-OR^{11}$, $-NR^{11}R^{12}$, $-SR^{11}$, $-NR^{11}S(O)_2R^{12}$, $-S(O)_2NR^{11}R^{12}$, $-S(O)NR^{11}R^{12}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-OS(O)_2R^{11}$, $-C(O)NR^{11}R^{12}$, $-OC(O)NR^{11}R^{12}$, $-NR^{11}C(O)R^{12}$, $-CH_2C(O)NR^{11}R^{12}$, $-OC_1$ - C_6 -alkyl- $C(O)NR^{11}R^{12}$, $-CH_2OR^{11}$, $-CH_2OC(O)R^{11}$, $-CH_2NR^{11}R^{12}$, $-OC(O)R^{11}$, $-OC_1$ - C_{15} -alkyl- $C(O)OR^{11}$, $-OC_1$ - C_6 -alkyl- OR^{11} , $-SC_1$ - C_6 -alkyl- $C(O)OR^{11}$, $-NR^{11}$ - $C(=O)$ - C_1 - C_6 -alkyl- $C(=O)OR^{11}$, $-C_2$ - C_6 -alkenyl- $C(=O)OR^{11}$,

30

$-\text{NR}^{11}-\text{C}(=\text{O})-\text{C}_1-\text{C}_6\text{-alkenyl}-\text{C}(=\text{O})\text{OR}^{11}$, $-\text{C}(\text{O})\text{OR}^{11}$, $\text{C}(\text{O})\text{R}^{11}$, or $-\text{C}_2-\text{C}_6\text{-alkenyl}-\text{C}(=\text{O})\text{R}^{11}$, $=\text{O}$, or $-\text{C}_2-\text{C}_6\text{-alkenyl}-\text{C}(=\text{O})-\text{NR}^{11}\text{R}^{12}$,

• $\text{C}_1-\text{C}_6\text{-alkyl}$, $\text{C}_2-\text{C}_6\text{-alkenyl}$ or $\text{C}_2-\text{C}_6\text{-alkynyl}$, each of which may optionally be substituted with one or more substituents independently selected from R^{13} ,

• aryl , aryloxy , aryloxycarbonyl , aroyl , arylsulfanyl , aryl- $\text{C}_1-\text{C}_6\text{-alkoxy}$, aryl- $\text{C}_1-\text{C}_6\text{-alkyl}$, aryl- $\text{C}_2-\text{C}_6\text{-alkenyl}$, aroyl- $\text{C}_2-\text{C}_6\text{-alkenyl}$, aryl- $\text{C}_2-\text{C}_6\text{-alkynyl}$, heteroaryl , heteroaryl- $\text{C}_1-\text{C}_6\text{-alkyl}$, heteroaryl- $\text{C}_2-\text{C}_6\text{-alkenyl}$, heteroaryl- $\text{C}_2-\text{C}_6\text{-alkynyl}$, or C_3-C_6 cycloalkyl ,

of which each cyclic moiety may optionally be substituted with one or more substituents independently selected from R^{14} ,

R^{11} and R^{12} are independently selected from hydrogen , OH , $\text{C}_1-\text{C}_{20}\text{-alkyl}$, aryl- $\text{C}_1-\text{C}_6\text{-alkyl}$ or aryl , wherein the alkyl groups may optionally be substituted with one or more substituents independently selected from R^{15} , and the aryl groups may optionally be substituted one or more substituents independently selected from R^{16} ; R^{11} and R^{12} when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom , the heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen , oxygen and sulphur , and optionally containing one or two double bonds ,

R^{13} is independently selected from halogen , -CN , $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OR}^{11}$, $-\text{C}(\text{O})\text{OR}^{11}$, $-\text{NR}^{11}\text{R}^{12}$, and $-\text{C}(\text{O})\text{NR}^{11}\text{R}^{12}$,

R^{14} is independently selected from halogen , $-\text{C}(\text{O})\text{OR}^{11}$, $-\text{CH}_2\text{C}(\text{O})\text{OR}^{11}$, $-\text{CH}_2\text{OR}^{11}$, -CN , $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{NO}_2$, $-\text{OR}^{11}$, $-\text{NR}^{11}\text{R}^{12}$, $-\text{NR}^{11}\text{C}(\text{O})\text{R}^{11}$, $-\text{S}(\text{O})_2\text{R}^{11}$, aryl and $\text{C}_1-\text{C}_6\text{-alkyl}$,

R^{15} is independently selected from halogen , -CN , $-\text{CF}_3$, $=\text{O}$, $-\text{OCF}_3$, $-\text{OC}_1-\text{C}_6\text{-alkyl}$, $-\text{C}(\text{O})\text{OC}_1-\text{C}_6\text{-alkyl}$, $-\text{COOH}$ and $-\text{NH}_2$,

R^{16} is independently selected from halogen , $-\text{C}(\text{O})\text{OC}_1-\text{C}_6\text{-alkyl}$, $-\text{COOH}$, -CN , $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{NO}_2$, -OH , $-\text{OC}_1-\text{C}_6\text{-alkyl}$, $-\text{NH}_2$, $\text{C}(=\text{O})$ or $\text{C}_1-\text{C}_6\text{-alkyl}$, or any enantiomer , diastereomer , including a racemic mixture , tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base .

In another embodiment hereof X is =O or =S.

In another embodiment hereof X is =O.

In another embodiment hereof X is =S.

In another embodiment hereof Y is -O- or -S-.

5 In another embodiment hereof Y is -O-.

In another embodiment hereof Y is -NH-.


In another embodiment hereof Y is -S-.

In another embodiment hereof A is aryl optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.

10 In another embodiment hereof A is selected from ArG1 optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.

In another embodiment hereof A is phenyl or naphthyl optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.

In another embodiment hereof A is



In another embodiment hereof A is phenyl.

In another embodiment hereof A is heteroaryl optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.

20 In another embodiment hereof A is selected from Het1 optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.


In another embodiment hereof A is selected from Het2 optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.

In another embodiment hereof A is selected from Het3 optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.

25 In another embodiment hereof A is selected from the group consisting of indolyl, benzofuranyl, quinolyl, furyl, thienyl, or pyrrolyl, wherein each heteroaryl may optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.


30 In another embodiment hereof A is benzofuranyl optionally substituted with up to four substituents R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.

In another embodiment hereof A is



In another embodiment hereof A is carbazolyl optionally substituted with up to four substituents R^7 , R^8 , R^9 , and R^{10} which may be the same or different.


In another embodiment hereof A is



5


In another embodiment hereof A is quinolyl optionally substituted with up to four substituents R^7 , R^8 , R^9 , and R^{10} which may be the same or different.

In another embodiment hereof A is



10 In another embodiment hereof A is indolyl optionally substituted with up to four substituents R^7 , R^8 , R^9 , and R^{10} which may be the same or different.

In another embodiment hereof A is



In another embodiment hereof R^1 is hydrogen.

15 In another embodiment hereof R^2 is hydrogen.

In another embodiment hereof R^1 and R^2 are combined to form a double bond.

In another embodiment hereof R^3 is C_1 - C_6 -alkyl, halogen, or $C(O)NR^{16}R^{17}$.

In another embodiment hereof R^3 is C_1 - C_6 -alkyl or $C(O)NR^{16}R^{17}$.

In another embodiment hereof R^3 is methyl.

In another embodiment hereof B is phenyl optionally substituted with up to four substituents, R^7 , R^8 , R^9 , and R^{10} which may be the same or different.

In another embodiment hereof R^4 is hydrogen.

In another embodiment hereof R^5 is hydrogen.

5 In another embodiment hereof R^6 is aryl.

In another embodiment hereof R^6 is phenyl.

In another embodiment hereof R^7 , R^8 , R^9 and R^{10} are independently selected from

10 •hydrogen, halogen, $-NO_2$, $-OR^{11}$, $-NR^{11}R^{12}$, $-SR^{11}$, $-NR^{11}S(O)_2R^{12}$, $-S(O)_2NR^{11}R^{12}$, $-S(O)NR^{11}R^{12}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-OS(O)_2R^{11}$, $-NR^{11}C(O)R^{12}$, $-CH_2OR^{11}$, $-CH_2OC(O)R^{11}$, $-CH_2NR^{11}R^{12}$, $-OC(O)R^{11}$, $-OC_1-C_6-alkyl-C(O)OR^{11}$, $-OC_1-C_6-alkyl-C(O)NR^{11}R^{12}$, $-OC_1-C_6-alkyl-OR^{11}$, $-SC_1-C_6-alkyl-C(O)OR^{11}$, $-C_2-C_6-alkenyl-C(=O)OR^{11}$, $-C(O)OR^{11}$, or $-C_2-C_6-alkenyl-C(=O)R^{11}$,

15 • $C_1-C_6-alkyl$, $C_2-C_6-alkenyl$ or $C_2-C_6-alkynyl$, which may each optionally be substituted with one or more substituents independently selected from R^{13}

20 •aryl, aryloxy, aroyl, arylsulfanyl, aryl- $C_1-C_6-alkoxy$, aryl- $C_1-C_6-alkyl$, aryl- $C_2-C_6-alkenyl$, aroyl- $C_2-C_6-alkenyl$, aryl- $C_2-C_6-alkynyl$, heteroaryl, heteroaryl- $C_1-C_6-alkyl$, wherein each of the cyclic moieties optionally may be substituted with one or more substituents independently selected from R^{14}

In another embodiment hereof R^7 , R^8 , R^9 and R^{10} are independently selected from

25 •hydrogen, halogen, $-NO_2$, $-OR^{11}$, $-NR^{11}R^{12}$, $-SR^{11}$, $-S(O)_2R^{11}$, $-OS(O)_2R^{11}$, $-CH_2OC(O)R^{11}$, $-OC(O)R^{11}$, $-OC_1-C_6-alkyl-C(O)OR^{11}$, $-OC_1-C_6-alkyl-OR^{11}$, $-SC_1-C_6-alkyl-C(O)OR^{11}$, $-C(O)OR^{11}$, or $-C_2-C_6-alkenyl-C(=O)R^{11}$,

30 • $C_1-C_6-alkyl$ or $C_1-C_6-alkenyl$ which may each optionally be substituted with one or more substituents independently selected from R^{13}

•aryl, aryloxy, aroyl, aryl- $C_1-C_6-alkoxy$, aryl- $C_1-C_6-alkyl$, heteroaryl,

35 of which each of the cyclic moieties optionally may be substituted with one or more substituents independently selected from R^{14}

In another embodiment hereof R^7 , R^8 , R^9 and R^{10} are independently selected from

5 • hydrogen, halogen, $-NO_2$, $-OR^{11}$, $-NR^{11}R^{12}$, $-SR^{11}$, $-S(O)_2R^{11}$, $-OS(O)_2R^{11}$, $-CH_2OC(O)R^{11}$, $-OC(O)R^{11}$, $-OC_1-C_6\text{-alkyl-C(O)OR}^{11}$, $-OC_1-C_6\text{-alkyl-OR}^{11}$, $-SC_1-C_6\text{-alkyl-C(O)OR}^{11}$, $-C(O)OR^{11}$, or $-C_2-C_6\text{-alkenyl-C(=O)R}^{11}$,

10 • $C_1-C_6\text{-alkyl}$ or $C_1-C_6\text{-}$ which may each optionally be substituted with one or more substituents independently selected from R^{13}

15 • aryl, aryloxy, aroyl, aryl- $C_1-C_6\text{-alkoxy}$, aryl- $C_1-C_6\text{-alkyl}$, heteroaryl,

of which each of the cyclic moieties optionally may be substituted with one or more substituents independently selected from R^{14} .

20 In another embodiment hereof R^7 , R^8 , R^9 and R^{10} are independently selected from

 • hydrogen, halogen, $-OR^{11}$, $-OC_1-C_6\text{-alkyl-C(O)OR}^{11}$, or $-C(O)OR^{11}$,

25 • $C_1-C_6\text{-alkyl}$ which may each optionally be substituted with one or more substituents independently selected from R^{13}

 • aryl, aryloxy, aryl- $C_1-C_6\text{-alkoxy}$,

of which each of the cyclic moieties optionally may be substituted with one or more substituents independently selected from R^{14} .

30 In another embodiment hereof R^7 , R^8 , R^9 and R^{10} are independently selected from

 • hydrogen, halogen, $-OR^{11}$, $-OC_1-C_6\text{-alkyl-C(O)OR}^{11}$, or $-C(O)OR^{11}$,

 • $C_1-C_6\text{-alkyl}$ which may each optionally be substituted with one or more substituents independently selected from R^{13}

 • ArG1, ArG1oxy, ArG1- $C_1-C_6\text{-alkoxy}$,

of which each of the cyclic moieties optionally may be substituted with one or more substituents independently selected from R¹⁴.

In another embodiment hereof R⁷, R⁸, R⁹ and R¹⁰ are independently selected from

- hydrogen, halogen, -OR¹¹, -OC₁-C₆-alkyl-C(O)OR¹¹, or -C(O)OR¹¹,

5

- C₁-C₆-alkyl which may optionally be substituted with one or more substituents independently selected from R¹³

10

- phenyl, phenyloxy, phenyl-C₁-C₆-alkoxy, wherein each of the cyclic moieties optionally may be substituted with one or more substituents independently selected from R¹⁴.

In another embodiment hereof R¹¹ and R¹² are independently selected from hydrogen, C₁-C₂₀-alkyl, aryl or aryl-C₁-C₆-alkyl, wherein the alkyl groups may optionally be substituted with one or more substituents independently selected from R¹⁵, and the aryl groups may optionally be substituted one or more substituents independently selected from R¹⁶, R¹¹ and R¹² when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom, the heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds.

15

20

In another embodiment hereof R¹¹ and R¹² are independently selected from hydrogen, C₁-C₂₀-alkyl, aryl or aryl-C₁-C₆-alkyl, wherein the alkyl groups may optionally be substituted with one or more substituents independently selected from R¹⁵, and the aryl groups may optionally be substituted one or more substituents independently selected from R¹⁶.

25

In another embodiment hereof R¹¹ and R¹² are independently selected from phenyl or phenyl-C₁-C₆-alkyl.

In another embodiment hereof one or both of R¹¹ and R¹² are methyl.

In another embodiment hereof R¹³ is independently selected from halogen, CF₃, OR¹¹ or NR¹¹R¹².

30

In another embodiment hereof R¹³ is independently selected from halogen or OR¹¹.

In another embodiment hereof R¹³ is OR¹¹.

In another embodiment hereof R¹⁴ is independently selected from halogen, -C(O)OR¹¹, -CN, -CF₃, -OR¹¹, S(O)₂R¹¹, and C₁-C₆-alkyl.

In another embodiment hereof R¹⁴ is independently selected from halogen, -C(O)OR¹¹, or -

35

OR¹¹.


In another embodiment hereof R^{15} is independently selected from halogen, -CN, -CF₃, -C(O)OC₁-C₆-alkyl, and -COOH.

In another embodiment hereof R^{15} is independently selected from halogen or -C(O)OC₁-C₆-alkyl.

- 5 In another embodiment hereof R^{16} is independently selected from halogen, -C(O)OC₁-C₆-alkyl, -COOH, -NO₂, -OC₁-C₆-alkyl, -NH₂, C(=O) or C₁-C₆-alkyl.

In another embodiment hereof R^{16} is independently selected from halogen, -C(O)OC₁-C₆-alkyl, -COOH, -NO₂, or C₁-C₆-alkyl.

In another embodiment hereof the zinc-binding ligand is



wherein

R^{19} is hydrogen or C₁-C₆-alkyl,

R^{20} is hydrogen or C₁-C₆-alkyl,

15

D, D¹ and F are a valence bond, C₁-C₆-alkylene or C₁-C₆-alkenylene optionally substituted with one or more substituents independently selected from R^{72} ,

R^{72} is independently selected from hydroxy, C₁-C₆-alkyl, or aryl,

20

E is C₁-C₆-alkyl, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with up to three substituents R^{21} , R^{22} and R^{23} ,

G and G¹ are C₁-C₆-alkyl, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with up to three substituents R^{24} , R^{25} and R^{26} ,

25

R^{17} , R^{18} , R^{21} , R^{22} , R^{23} , R^{24} , R^{25} and R^{26} are independently selected from

- hydrogen, halogen, -CN, -CH₂CN, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -S(O)₂CF₃, -SCF₃, -NO₂, =O, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷S(O)₂R²⁸,

-S(O)₂NR²⁷R²⁸, -S(O)NR²⁷R²⁸, -S(O)R²⁷, -S(O)₂R²⁷, -C(O)NR²⁷R²⁸, -OC(O)NR²⁷R²⁸,
 -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -CH₂C(O)NR²⁷R²⁸, -OCH₂C(O)NR²⁷R²⁸, -CH₂OR²⁷,
 -CH₂NR²⁷R²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-
 alkenyl-C(=O)OR²⁷, -NR²⁷-C(=O)-C₁-C₆-alkyl-C(=O)OR²⁷, -NR²⁷-C(=O)-C₁-C₆-
 alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or
 -C(O)OR²⁷,

• C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl,

which may optionally be substituted with one or more substituents independently selected from R²⁹,

• aryl, aryloxy, aryloxycarbonyl, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl or heteroaryl-C₂-C₆-alkynyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰,

R²⁷ and R²⁸ are independently selected from hydrogen, C₁-C₆-alkyl, aryl-C₁-C₆-alkyl or aryl, or R²⁷ and R²⁸ when attached to the same nitrogen atom together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds,

R²⁹ is independently selected from halogen, -CN, -CF₃, -OCF₃, -OR²⁷, and -NR²⁷R²⁸,

R³⁰ is independently selected from halogen, -C(O)OR²⁷, -CN, -CF₃, -OCF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸ and C₁-C₆-alkyl, or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.

In another embodiment hereof D is a valence bond.

In another embodiment hereof D is C₁-C₆-alkylene optionally substituted with one or more hydroxy, C₁-C₆-alkyl, or aryl.


In another embodiment hereof E is aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with up to three substituents independently selected from R^{21} , R^{22} and R^{23} .

5 In another embodiment hereof E is aryl optionally substituted with up to three substituents independently selected from R^{21} , R^{22} and R^{23} .

In another embodiment hereof E is selected from ArG1 and optionally substituted with up to three substituents independently selected from R^{21} , R^{22} and R^{23} .

In another embodiment hereof E is phenyl optionally substituted with up to three substituents independently selected from R^{21} , R^{22} and R^{23} .

10 In another embodiment hereof the zinc-binding ligand is



In another embodiment hereof R^{21} , R^{22} and R^{23} are independently selected from

15 • hydrogen, halogen, $-\text{CHF}_2$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OCHF}_2$, $-\text{OCH}_2\text{CF}_3$, $-\text{OCF}_2\text{CHF}_2$, $-\text{SCF}_3$, $-\text{NO}_2$, $-\text{OR}^{27}$, $-\text{NR}^{27}\text{R}^{28}$, $-\text{SR}^{27}$, $-\text{C}(\text{O})\text{NR}^{27}\text{R}^{28}$, $-\text{OC}(\text{O})\text{NR}^{27}\text{R}^{28}$, $-\text{NR}^{27}\text{C}(\text{O})\text{R}^{28}$, $-\text{NR}^{27}\text{C}(\text{O})\text{OR}^{28}$, $-\text{CH}_2\text{C}(\text{O})\text{NR}^{27}\text{R}^{28}$, $-\text{OCH}_2\text{C}(\text{O})\text{NR}^{27}\text{R}^{28}$, $-\text{CH}_2\text{OR}^{27}$, $-\text{CH}_2\text{NR}^{27}\text{R}^{28}$, $-\text{OC}(\text{O})\text{R}^{27}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{SC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}_2\text{-C}_6\text{-alkenyl-C}(\text{O})\text{OR}^{27}$, $-\text{NR}^{27}\text{-C}(\text{O})\text{-C}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{NR}^{27}\text{-C}(\text{O})\text{-C}_1\text{-C}_6\text{-alkenyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}(\text{O})\text{NR}^{27}\text{-C}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, or $-\text{C}(\text{O})\text{OR}^{27}$,

20

• $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_2\text{-C}_6\text{-alkenyl}$ or $\text{C}_2\text{-C}_6\text{-alkynyl}$,

25 which may optionally be substituted with one or more substituents independently selected from R^{29}

30 • aryl, aryloxy, aryloxycarbonyl, aroyl, aryl- $\text{C}_1\text{-C}_6\text{-alkoxy}$, aryl- $\text{C}_1\text{-C}_6\text{-alkyl}$, aryl- $\text{C}_2\text{-C}_6\text{-alkenyl}$, aryl- $\text{C}_2\text{-C}_6\text{-alkynyl}$, heteroaryl, heteroaryl- $\text{C}_1\text{-C}_6\text{-alkyl}$, heteroaryl- $\text{C}_2\text{-C}_6\text{-alkenyl}$ or heteroaryl- $\text{C}_2\text{-C}_6\text{-alkynyl}$,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R^{30} .

In another embodiment hereof R^{21} , R^{22} and R^{23} are independently selected from

5

• hydrogen, halogen, $-OCF_3$, $-OR^{27}$, $-NR^{27}R^{28}$, $-SR^{27}$, $-NR^{27}C(O)R^{28}$, $-NR^{27}C(O)OR^{28}$, $-OC(O)R^{27}$, $-OC_1-C_6\text{-alkyl-C}(O)OR^{27}$, $-SC_1-C_6\text{-alkyl-C}(O)OR^{27}$, $-C_2-C_6\text{-alkenyl-C}(=O)OR^{27}$, $-C(=O)NR^{27}-C_1-C_6\text{-alkyl-C}(=O)OR^{27}$, $-C_1-C_6\text{-alkyl-C}(=O)OR^{27}$, or $-C(O)OR^{27}$,

10

• $C_1-C_6\text{-alkyl}$ optionally substituted with one or more substituents independently selected from R^{29}

15

• aryl, aryloxy, aroyl, aryl- $C_1-C_6\text{-alkoxy}$, aryl- $C_1-C_6\text{-alkyl}$, heteroaryl, heteroaryl- $C_1-C_6\text{-alkyl}$,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R^{30} .

20

In another embodiment hereof R^{21} , R^{22} and R^{23} are independently selected from

25

• hydrogen, halogen, $-OCF_3$, $-OR^{27}$, $-NR^{27}R^{28}$, $-SR^{27}$, $-NR^{27}C(O)R^{28}$, $-NR^{27}C(O)OR^{28}$, $-OC(O)R^{27}$, $-OC_1-C_6\text{-alkyl-C}(O)OR^{27}$, $-SC_1-C_6\text{-alkyl-C}(O)OR^{27}$, $-C_2-C_6\text{-alkenyl-C}(=O)OR^{27}$, $-C(=O)NR^{27}-C_1-C_6\text{-alkyl-C}(=O)OR^{27}$, $-C_1-C_6\text{-alkyl-C}(=O)OR^{27}$, or $-C(O)OR^{27}$,

• methyl, ethyl propyl optionally substituted with one or more substituents independently selected from R^{29}

30

• aryl, aryloxy, aroyl, aryl- $C_1-C_6\text{-alkoxy}$, aryl- $C_1-C_6\text{-alkyl}$, heteroaryl, heteroaryl- $C_1-C_6\text{-alkyl}$

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R^{30} .

In another embodiment hereof R^{21} , R^{22} and R^{23} are independently selected from

35

• hydrogen, halogen, $-\text{OCF}_3$, $-\text{OR}^{27}$, $-\text{NR}^{27}\text{R}^{28}$, $-\text{SR}^{27}$, $-\text{NR}^{27}\text{C}(\text{O})\text{R}^{28}$, $-\text{NR}^{27}\text{C}(\text{O})\text{OR}^{28}$, $-\text{OC}(\text{O})\text{R}^{27}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{SC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}_2\text{-C}_6\text{-alkenyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}(\text{O})\text{NR}^{27}\text{-C}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, or $-\text{C}(\text{O})\text{OR}^{27}$,

5

• methyl, ethyl propyl optionally substituted with one or more substituents independently selected from R^{29}

• ArG1 , ArG1-O- , $\text{ArG1-C}(\text{O})\text{-}$, $\text{ArG1-C}_1\text{-C}_6\text{-alkoxy}$, $\text{ArG1-C}_1\text{-C}_6\text{-alkyl}$, Het3 , $\text{Het3-C}_1\text{-C}_6\text{-alkyl}$

10

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R^{30} .

In another embodiment hereof R^{21} , R^{22} and R^{23} are independently selected from

15

• hydrogen, halogen, $-\text{OCF}_3$, $-\text{OR}^{27}$, $-\text{NR}^{27}\text{R}^{28}$, $-\text{SR}^{27}$, $-\text{NR}^{27}\text{C}(\text{O})\text{R}^{28}$, $-\text{NR}^{27}\text{C}(\text{O})\text{OR}^{28}$, $-\text{OC}(\text{O})\text{R}^{27}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{SC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}_2\text{-C}_6\text{-alkenyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}(\text{O})\text{NR}^{27}\text{-C}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, or $-\text{C}(\text{O})\text{OR}^{27}$,

20

• $\text{C}_1\text{-C}_6\text{-alkyl}$ optionally substituted with one or more substituents independently selected from R^{29}

• phenyl, phenyloxy, phenyl- $\text{C}_1\text{-C}_6\text{-alkoxy}$, phenyl- $\text{C}_1\text{-C}_6\text{-alkyl}$,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R^{30} .

25

In another embodiment hereof R^{19} is hydrogen or methyl.

In another embodiment hereof R^{19} is hydrogen.

In another embodiment hereof R^{27} is hydrogen, $\text{C}_1\text{-C}_6\text{-alkyl}$ or aryl.

30

In another embodiment hereof R^{27} is hydrogen or $\text{C}_1\text{-C}_6\text{-alkyl}$.

In another embodiment hereof R^{28} is hydrogen or $\text{C}_1\text{-C}_6\text{-alkyl}$.

In another embodiment hereof F is a valence bond.

In another embodiment hereof F is $\text{C}_1\text{-C}_6\text{-alkylene}$ optionally substituted with one or more hydroxy, $\text{C}_1\text{-C}_6\text{-alkyl}$, or aryl.

In another embodiment hereof G is C₁-C₆-alkyl or aryl, wherein the aryl is optionally substituted with up to three substituents R²⁴, R²⁵ and R²⁶.

In another embodiment hereof G is C₁-C₆-alkyl or ArG1, wherein the aryl is optionally substituted with up to three substituents R²⁴, R²⁵ and R²⁶.

5 In another embodiment hereof G is C₁-C₆-alkyl.

In another embodiment hereof G is phenyl optionally substituted with up to three substituents R²⁴, R²⁵ and R²⁶.

In another embodiment hereof R²⁴, R²⁵ and R²⁶ are independently selected from

10 • hydrogen, halogen, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -SCF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -C(O)NR²⁷R²⁸, -OC(O)NR²⁷R²⁸, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -CH₂C(O)NR²⁷R²⁸, -OCH₂C(O)NR²⁷R²⁸, -CH₂OR²⁷, -CH₂NR²⁷R²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -NR²⁷-C(=O)-C₁-C₆-alkyl-C(=O)OR²⁷, -NR²⁷-C(=O)-C₁-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

• C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl,

20 which may optionally be substituted with one or more substituents independently selected from R²⁹

• aryl, aryloxy, aryloxycarbonyl, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl or heteroaryl-C₂-C₆-alkynyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

30 In another embodiment hereof R²⁴, R²⁵ and R²⁶ are independently selected from

• hydrogen, halogen, -OCF₃, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

- C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl,

which may optionally be substituted with one or more substituents independently selected from R²⁹

- aryl, aryloxy, aryloxycarbonyl, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl or heteroaryl-C₂-C₆-alkynyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

In another embodiment hereof R²⁴, R²⁵ and R²⁶ are independently selected from

- hydrogen, halogen, -OCF₃, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

- C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R²⁹

- aryl, aryloxy, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, heteroaryl, heteroaryl-C₁-C₆-alkyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

In another embodiment hereof R²¹, R²² and R²³ are independently selected from

- hydrogen, halogen, -OCF₃, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

•methyl, ethyl propyl optionally substituted with one or more substituents independently selected from R²⁹

5 •ArG1, ArG1-O-, ArG1-C(O)-, ArG1-C₁-C₆-alkoxy, ArG1-C₁-C₆-alkyl, Het3, Het3-C₁-C₆-alkyl

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

In another embodiment hereof R²¹, R²² and R²³ are independently selected from

10

•hydrogen, halogen, -OCF₃, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

15

•methyl, ethyl propyl optionally substituted with one or more substituents independently selected from R²⁹

20

•ArG1, ArG1-O-, ArG1-C(O)-, ArG1-C₁-C₆-alkoxy, ArG1-C₁-C₆-alkyl, Het3, Het3-C₁-C₆-alkyl

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

In another embodiment hereof R²¹, R²² and R²³ are independently selected from

25

•hydrogen, halogen, -OCF₃, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

30

•methyl, ethyl propyl optionally substituted with one or more substituents independently selected from R²⁹

•ArG1, ArG1-O-, ArG1-C₁-C₆-alkoxy, ArG1-C₁-C₆-alkyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

35

In another embodiment hereof R^{20} is hydrogen or methyl.

In another embodiment hereof R^{20} is hydrogen.

In another embodiment hereof R^{27} is hydrogen, C_1 - C_6 -alkyl or aryl.

In another embodiment hereof R^{27} is hydrogen or C_1 - C_6 -alkyl or ArG1.

5 In another embodiment hereof R^{27} is hydrogen or C_1 - C_6 -alkyl.

In another embodiment hereof R^{28} is hydrogen or C_1 - C_6 -alkyl.

In another embodiment hereof R^{17} and R^{18} are independently selected from

10 • hydrogen, halogen, -CN, -CF₃, -OCF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -S(O)R²⁷,
-S(O)₂R²⁷, -C(O)NR²⁷R²⁸, -CH₂OR²⁷, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-
alkyl-C(O)OR²⁷, or -C(O)OR²⁷,

15 • C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl or C_2 - C_6 -alkynyl, optionally substituted with one or more
substituents independently selected from R^{29}

• aryl, aryloxy, aroyl, aryl- C_1 - C_6 -alkoxy, aryl- C_1 - C_6 -alkyl, heteroaryl, heteroaryl- C_1 - C_6 -
alkyl,

20 of which the cyclic moieties optionally may be substituted with one or more
substituents selected from R^{30} .

In another embodiment hereof R^{17} and R^{18} are independently selected from

• hydrogen, halogen, -CN, -CF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, or -C(O)OR²⁷,

25 • C_1 - C_6 -alkyl optionally substituted with one or more substituents independently
selected from R^{29}

• aryl, aryloxy, aroyl, aryl- C_1 - C_6 -alkoxy, aryl- C_1 - C_6 -alkyl, heteroaryl, heteroaryl- C_1 - C_6 -
alkyl,

30 of which the cyclic moieties optionally may be substituted with one or more
substituents selected from R^{30} .

In another embodiment hereof R^{17} and R^{18} are independently selected from

35 • hydrogen, halogen, -CN, -CF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, or -C(O)OR²⁷

- methyl, ethyl propyl optionally substituted with one or more substituents independently selected from R^{29}

- aryl, aryloxy, aroyl, aryl- C_1-C_6 -alkoxy, aryl- C_1-C_6 -alkyl, heteroaryl, heteroaryl- C_1-C_6 -alkyl

5 of which the cyclic moieties optionally may be substituted with one or more substituents selected from R^{30} .

In another embodiment hereof R^{17} and R^{18} are independently selected from

- hydrogen, halogen, -CN, -CF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, or -C(O)OR²⁷

10 • methyl, ethyl propyl optionally substituted with one or more substituents independently selected from R^{29}

- ArG1, ArG1-O-, ArG1-C(O)-, ArG1- C_1-C_6 -alkoxy, ArG1- C_1-C_6 -alkyl, Het3, Het3- C_1-C_6 -alkyl

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R^{30} .

15 In another embodiment hereof R^{17} and R^{18} are independently selected from

- hydrogen, halogen, -CN, -CF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, or -C(O)OR²⁷

- C_1-C_6 -alkyl optionally substituted with one or more substituents independently selected from R^{29}

- phenyl, phenyloxy, phenyl- C_1-C_6 -alkoxy, phenyl- C_1-C_6 -alkyl,

20 of which the cyclic moieties optionally may be substituted with one or more substituents selected from R^{30} .

In another embodiment hereof R^{27} is hydrogen or C_1-C_6 -alkyl.


In another embodiment hereof R^{27} is hydrogen, methyl or ethyl.

In another embodiment hereof R^{28} is hydrogen or C_1-C_6 -alkyl.

25 In another embodiment hereof R^{28} is hydrogen, methyl or ethyl.

In another embodiment hereof R^{72} is -OH or phenyl.


In another embodiment hereof the zinc-binding ligand is



30

In another embodiment hereof the zinc-binding ligand is of the form H-I-J


wherein H is



wherein the phenyl, naphthalene or benzocarbazole rings are optionally substituted with one or more substituents independently selected from R^{31}

10 I is selected from

- a valence bond,
- $-\text{CH}_2\text{N}(\text{R}^{32})-$ or $-\text{SO}_2\text{N}(\text{R}^{33})-$,



- $-\text{Z}^1-\text{N}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{Z}^2-$ wherein Z^1 is $\text{S}(\text{O})_2$ or CH_2 , Z^2 is $-\text{NH}-$, $-\text{O}-$ or $-\text{S}-$, and n is 1 or 2,

15 J is

- $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_2\text{-C}_6\text{-alkenyl}$ or $\text{C}_2\text{-C}_6\text{-alkynyl}$, which may each optionally be substituted with one or more substituents selected from R^{34} ,
- Aryl, aryloxy, aryl-oxycarbonyl-, aroyl, aryl- $\text{C}_1\text{-C}_6\text{-alkoxy-}$, aryl- $\text{C}_1\text{-C}_6\text{-alkyl-}$, aryl- $\text{C}_2\text{-C}_6\text{-alkenyl-}$, aryl- $\text{C}_2\text{-C}_6\text{-alkynyl-}$, heteroaryl, heteroaryl- $\text{C}_1\text{-C}_6\text{-alkyl-}$, heteroaryl- $\text{C}_2\text{-C}_6\text{-alkenyl-}$ or heteroaryl- $\text{C}_2\text{-C}_6\text{-alkynyl-}$, wherein the cyclic moieties are optionally substituted with one or more substituents selected from R^{37} ,
- hydrogen,

20

R^{31} is independently selected from hydrogen, halogen, $-\text{CN}$, $-\text{CH}_2\text{CN}$, $-\text{CHF}_2$, $-\text{CF}_3$, $-\text{OCF}_3$,
 25 $-\text{OCHF}_2$, $-\text{OCH}_2\text{CF}_3$, $-\text{OCF}_2\text{CHF}_2$, $-\text{S}(\text{O})_2\text{CF}_3$, $-\text{SCF}_3$, $-\text{NO}_2$, $-\text{OR}^{35}$, $-\text{C}(\text{O})\text{R}^{35}$, $-\text{NR}^{35}\text{R}^{36}$, $-\text{SR}^{35}$,
 $-\text{NR}^{35}\text{S}(\text{O})_2\text{R}^{36}$, $-\text{S}(\text{O})_2\text{NR}^{35}\text{R}^{36}$, $-\text{S}(\text{O})\text{NR}^{35}\text{R}^{36}$, $-\text{S}(\text{O})\text{R}^{35}$, $-\text{S}(\text{O})_2\text{R}^{35}$, $-\text{C}(\text{O})\text{NR}^{35}\text{R}^{36}$,
 $-\text{OC}(\text{O})\text{NR}^{35}\text{R}^{36}$, $-\text{NR}^{35}\text{C}(\text{O})\text{R}^{36}$, $-\text{CH}_2\text{C}(\text{O})\text{NR}^{35}\text{R}^{36}$, $-\text{OCH}_2\text{C}(\text{O})\text{NR}^{35}\text{R}^{36}$, $-\text{CH}_2\text{OR}^{35}$,
 $-\text{CH}_2\text{NR}^{35}\text{R}^{36}$, $-\text{OC}(\text{O})\text{R}^{35}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{35}$, $-\text{SC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{35}$, $-\text{C}_2\text{-C}_6\text{-alkenyl-}$
 $\text{C}(\text{O})\text{OR}^{35}$, $-\text{NR}^{35}\text{-C}(\text{O})\text{-C}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{35}$, $-\text{NR}^{35}\text{-C}(\text{O})\text{-C}_1\text{-C}_6\text{-alkenyl-C}(\text{O})\text{OR}^{35}$,
 30 $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_1\text{-C}_6\text{-alkanoyl}$ or $-\text{C}(\text{O})\text{OR}^{35}$,

R^{32} and R^{33} are independently selected from hydrogen, $\text{C}_1\text{-C}_6\text{-alkyl}$ or $\text{C}_1\text{-C}_6\text{-alkanoyl}$,

R^{34} is independently selected from halogen, -CN, -CF₃, -OCF₃, -OR³⁵, and -NR³⁵R³⁶,


5 R^{35} and R^{36} are independently selected from hydrogen, C₁-C₆-alkyl, aryl-C₁-C₆-alkyl or aryl, or R^{35} and R^{36} when attached to the same nitrogen atom together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds,

10 R^{37} is independently selected from halogen, -C(O)OR³⁵, -C(O)H, -CN, -CF₃, -OCF₃, -NO₂, -OR³⁵, -NR³⁵R³⁶, C₁-C₆-alkyl or C₁-C₆-alkanoyl,

or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.

15


In another embodiment hereof the zinc-binding ligand is of the form H-I-J, wherein H is wherein the phenyl, naphthalene or benzocarbazole rings are optionally substituted with one



or more substituents independently selected from R^{31} ,

20 I is selected from

- a valence bond,
- -CH₂N(R³²)- or -SO₂N(R³³)-,



- wherein Z¹ is S(O)₂ or CH₂, Z² is N, -O- or -S-, and n is 1 or 2,

25 J is

- C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl, which may each optionally be substituted with one or more substituents selected from R^{34} ,

• Aryl, aryloxy, aryl-oxycarbonyl-, aroyl, aryl-C₁-C₆-alkoxy-, aryl-C₁-C₆-alkyl-, aryl-C₂-C₆-alkenyl-, aryl-C₂-C₆-alkynyl-, heteroaryl, heteroaryl-C₁-C₆-alkyl-, heteroaryl-C₂-C₆-alkenyl- or heteroaryl-C₂-C₆-alkynyl-, wherein the cyclic moieties are optionally substituted with one or more substituents selected from R³⁷,

5 • hydrogen,

R³¹ is independently selected from hydrogen, halogen, -CN, -CH₂CN, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -S(O)₂CF₃, -SCF₃, -NO₂, -OR³⁵, -C(O)R³⁵, -NR³⁵R³⁶, -SR³⁵, -NR³⁵S(O)₂R³⁶, -S(O)₂NR³⁵R³⁶, -S(O)NR³⁵R³⁶, -S(O)R³⁵, -S(O)₂R³⁵, -C(O)NR³⁵R³⁶,
 10 -OC(O)NR³⁵R³⁶, -NR³⁵C(O)R³⁶, -CH₂C(O)NR³⁵R³⁶, -OCH₂C(O)NR³⁵R³⁶, -CH₂OR³⁵, -CH₂NR³⁵R³⁶, -OC(O)R³⁵, -OC₁-C₆-alkyl-C(O)OR³⁵, -SC₁-C₆-alkyl-C(O)OR³⁵ -C₂-C₆-alkenyl-C(=O)OR³⁵, -NR³⁵-C(=O)-C₁-C₆-alkyl-C(=O)OR³⁵, -NR³⁵-C(=O)-C₁-C₆-alkenyl-C(=O)OR³⁵, C₁-C₆-alkyl, C₁-C₆-alkanoyl or -C(O)OR³⁵,

15 R³² and R³³ are independently selected from hydrogen, C₁-C₆-alkyl or C₁-C₆-alkanoyl,

R³⁴ is independently selected from halogen, -CN, -CF₃, -OCF₃, -OR³⁵, and -NR³⁵R³⁶,


R³⁵ and R³⁶ are independently selected from hydrogen, C₁-C₆-alkyl, aryl-C₁-C₆-alkyl or aryl, or
 20 R³⁵ and R³⁶ when attached to the same nitrogen atom together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds,

25 R³⁷ is independently selected from halogen, -C(O)OR³⁵, -C(O)H, -CN, -CF₃, -OCF₃, -NO₂, -OR³⁵, -NR³⁵R³⁶, C₁-C₆-alkyl or C₁-C₆-alkanoyl,


or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base,

30 With the proviso that R³¹ and J cannot both be hydrogen.


In another embodiment hereof H is



In another embodiment hereof H is



In another embodiment hereof H is



5 In another embodiment hereof I is a valence bond, $-\text{CH}_2\text{N}(\text{R}^{32})-$, or $-\text{SO}_2\text{N}(\text{R}^{33})-$.

In another embodiment hereof I is a valence bond.

In another embodiment hereof J is

- hydrogen,
 - C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl or C_2 - C_6 -alkynyl,
- 10 which may optionally be substituted with one or more substituents selected from halogen, $-\text{CN}$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OR}^{35}$, and $-\text{NR}^{35}\text{R}^{36}$,
- aryl, or heteroaryl, wherein the cyclic moieties are optionally substituted with one or more substituents independently selected from R^{37} .

In another embodiment hereof J is

- 15
- hydrogen,
 - aryl or heteroaryl, wherein the cyclic moieties are optionally substituted with one or more substituents independently selected from R^{37} .

In another embodiment hereof J is

- 20
- hydrogen,
 - ArG1 or Het3, wherein the cyclic moieties are optionally substituted with one or more substituents independently selected from R^{37} .

In another embodiment hereof J is

- 25
- hydrogen,
 - phenyl or naphthyl optionally substituted with one or more substituents independently selected from R^{37} .

In another embodiment hereof J is hydrogen.

In another embodiment hereof R^{32} and R^{33} are independently selected from hydrogen or C_1 - C_6 -alkyl.

In another embodiment hereof R^{34} is hydrogen, halogen, -CN, -CF₃, -OCF₃, -SCF₃, -NO₂, -OR³⁵, -C(O)R³⁵, -NR³⁵R³⁶, -SR³⁵, -C(O)NR³⁵R³⁶, -OC(O)NR³⁵R³⁶, -NR³⁵C(O)R³⁶, -OC(O)R³⁵, -OC₁-C₆-alkyl-C(O)OR³⁵, -SC₁-C₆-alkyl-C(O)OR³⁵ or -C(O)OR³⁵.

In another embodiment hereof R^{34} is hydrogen, halogen, -CF₃, -NO₂, -OR³⁵, -NR³⁵R³⁶, -SR³⁵,
5 -NR³⁵C(O)R³⁶, or -C(O)OR³⁵.

In another embodiment hereof R^{34} is hydrogen, halogen, -CF₃, -NO₂, -OR³⁵, -NR³⁵R³⁶, or -NR³⁵C(O)R³⁶.

In another embodiment hereof R^{34} is hydrogen, halogen, or -OR³⁵.

In another embodiment hereof R^{35} and R^{36} are independently selected from hydrogen, C₁-C₆-alkyl, or aryl.
10


In another embodiment hereof R^{35} and R^{36} are independently selected from hydrogen or C₁-C₆-alkyl.

In another embodiment hereof R^{37} is halogen, -C(O)OR³⁵, -CN, -CF₃, -OR³⁵, -NR³⁵R³⁶, C₁-C₆-alkyl or C₁-C₆-alkanoyl.

15 In another embodiment hereof R^{37} is halogen, -C(O)OR³⁵, -OR³⁵, -NR³⁵R³⁶, C₁-C₆-alkyl or C₁-C₆-alkanoyl.

In another embodiment hereof R^{37} is halogen, -C(O)OR³⁵ or -OR³⁵.

In another embodiment hereof the zinc-binding ligand is



20

wherein K is a valence bond, C₁-C₆-alkylene, -NH-C(=O)-U-, -C₁-C₆-alkyl-S-, -C₁-C₆-alkyl-O-, -C(=O)-, or -C(=O)-NH-, wherein any C₁-C₆-alkyl moiety is optionally substituted with R^{38} ,

25 U is a valence bond, C₁-C₆-alkenylene, -C₁-C₆-alkyl-O- or C₁-C₆-alkylene wherein any C₁-C₆-alkyl moiety is optionally substituted with C₁-C₆-alkyl,

R^{38} is C₁-C₆-alkyl, aryl, wherein the alkyl or aryl moieties are optionally substituted with one or more substituents independently selected from R^{39} ,

30 R^{39} is independently selected from halogen, cyano, nitro, amino,

M is a valence bond, arylene or heteroarylene, wherein the aryl or heteroaryl moieties are optionally substituted with one or more substituents independently selected from R^{40} ,

R⁴⁰ is selected from

• hydrogen, halogen, -CN, -CH₂CN, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃,
 -OCF₂CHF₂, -S(O)₂CF₃, -OS(O)₂CF₃, -SCF₃, -NO₂, -OR⁴¹, -NR⁴¹R⁴², -SR⁴¹,
 -NR⁴¹S(O)₂R⁴², -S(O)₂NR⁴¹R⁴², -S(O)NR⁴¹R⁴², -S(O)R⁴¹, -S(O)₂R⁴¹, -OS(O)₂R⁴¹,
 -C(O)NR⁴¹R⁴², -OC(O)NR⁴¹R⁴², -NR⁴¹C(O)R⁴², -CH₂C(O)NR⁴¹R⁴², -OC₁-C₆-
 alkyl-C(O)NR⁴¹R⁴², -CH₂OR⁴¹, -CH₂OC(O)R⁴¹, -CH₂NR⁴¹R⁴², -OC(O)R⁴¹, -OC₁-C₆-
 alkyl-C(O)OR⁴¹, -OC₁-C₆-alkyl-OR⁴¹, -S-C₁-C₆-alkyl-C(O)OR⁴¹, -C₂-C₆-alkenyl-
 C(=O)OR⁴¹, -NR⁴¹-C(=O)-C₁-C₆-alkyl-C(=O)OR⁴¹, -NR⁴¹-C(=O)-C₁-C₆-
 alkenyl-C(=O)OR⁴¹, -C(O)OR⁴¹, -C₂-C₆-alkenyl-C(=O)R⁴¹, =O, -NH-C(=O)-O-C₁-
 C₆-alkyl, or -NH-C(=O)-C(=O)-O-C₁-C₆-alkyl,

• C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl, which may each optionally be substituted
 with one or more substituents selected from R⁴³,

• aryl, aryloxy, aryloxycarbonyl, aroyl, arylsulfanyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl,
 aryl-C₂-C₆-alkenyl, aroyl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-
 C₆-alkyl, heteroaryl-C₂-C₆-alkenyl or heteroaryl-C₂-C₆-alkynyl, wherein the cyclic
 moieties optionally may be substituted with one or more substituents selected from
 R⁴⁴,

R⁴¹ and R⁴² are independently selected from hydrogen, -OH, C₁-C₆-alkyl, C₁-C₆-alkenyl, aryl-
 C₁-C₆-alkyl or aryl, wherein the alkyl moieties may optionally be substituted with one or more
 substituents independently selected from R⁴⁵, and the aryl moieties may optionally be
 substituted with one or more substituents independently selected from R⁴⁶; R⁴¹ and R⁴² when
 attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the
 said nitrogen atom, the heterocyclic ring optionally containing one or two further heteroatoms
 selected from nitrogen, oxygen and sulphur, and optionally containing one or two double
 bonds,

R⁴³ is independently selected from halogen, -CN, -CF₃, -OCF₃, -OR⁴¹, and -NR⁴¹R⁴²

R⁴⁴ is independently selected from halogen, -C(O)OR⁴¹, -CH₂C(O)OR⁴¹, -CH₂OR⁴¹, -CN, -
 CF₃, -OCF₃, -NO₂, -OR⁴¹, -NR⁴¹R⁴² and C₁-C₆-alkyl,

R⁴⁵ is independently selected from halogen, -CN, -CF₃, -OCF₃, -O-C₁-C₆-alkyl, -C(O)-O-C₁-C₆-alkyl, -COOH and -NH₂,

R⁴⁶ is independently selected from halogen, -C(O)OC₁-C₆-alkyl, -COOH, -CN, -CF₃, -OCF₃, -NO₂, -OH, -OC₁-C₆-alkyl, -NH₂, C(=O) or C₁-C₆-alkyl,

5

Q is a valence bond, C₁-C₆-alkylene, -C₁-C₆-alkyl-O-, -C₁-C₆-alkyl-NH-, -NH-C₁-C₆-alkyl, -NH-C(=O)-, -C(=O)-NH-, -O-C₁-C₆-alkyl, -C(=O)-, or -C₁-C₆-alkyl-C(=O)-N(R⁴⁷)- wherein the alkyl moieties are optionally substituted with one or more substituents independently selected from R⁴⁸,

10

R⁴⁷ and R⁴⁸ are independently selected from hydrogen, C₁-C₆-alkyl, aryl optionally substituted with one or more R⁴⁹,

R⁴⁹ is independently selected from halogen and -COOH,

15

T is

- hydrogen,
- C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₆-alkyloxy-carbonyl, wherein the alkyl, alkenyl and alkynyl moieties are optionally substituted with one or more substituents independently selected from R⁵⁰,
- aryl, aryloxy, aryloxy-carbonyl, aryl-C₁-C₆-alkyl, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl-, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl, heteroaryl-C₂-C₆-alkynyl,

20

25

wherein any alkyl, alkenyl, alkynyl, aryl and heteroaryl moiety is optionally substituted with one or more substituents independently selected from R⁵⁰,

R⁵⁰ is C₁-C₆-alkyl, C₁-C₆-alkoxy, aryl, aryloxy, aryl-C₁-C₆-alkoxy, -C(=O)-NH-C₁-C₆-alkyl-aryl, -C(=O)-NR^{50A}-C₁-C₆-alkyl, -C(=O)-NH-(CH₂CH₂O)_m-C₁-C₆-alkyl-COOH, heteroaryl, heteroaryl-C₁-C₆-alkoxy, -C₁-C₆-alkyl-COOH, -O-C₁-C₆-alkyl-COOH, -S(O)₂R⁵¹, -C₂-C₆-alkenyl-COOH, -OR⁵¹, -NO₂, halogen, -COOH, -CF₃, -CN, =O, -N(R⁵¹R⁵²), wherein m is 1, 2, 3 or 4, and wherein the aryl or heteroaryl moieties are optionally substituted with one or more R⁵³, and the alkyl moieties are optionally substituted with one or more R^{50B}.

30

R^{50A} and R^{50B} are independently selected from $-C(O)OC_1-C_6\text{-alkyl}$, $-COOH$, $-C_1-C_6\text{-alkyl}-C(O)OC_1-C_6\text{-alkyl}$, $-C_1-C_6\text{-alkyl}-COOH$, or $C_1-C_6\text{-alkyl}$,

R^{51} and R^{52} are independently selected from hydrogen and $C_1-C_6\text{-alkyl}$,

R^{53} is independently selected from $C_1-C_6\text{-alkyl}$, $C_1-C_6\text{-alkoxy}$, $-C_1-C_6\text{-alkyl}-COOH$, $-C_2\text{-C}_6\text{-alkenyl}-COOH$, $-OR^{51}$, $-NO_2$, halogen, $-COOH$, $-CF_3$, $-CN$, or $-N(R^{51}R^{52})$,

or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.

In another embodiment hereof K is a valence bond, $C_1-C_6\text{-alkylene}$, $-NH-C(=O)-U-$, $-C_1-C_6\text{-alkyl}-S-$, $-C_1-C_6\text{-alkyl}-O-$, or $-C(=O)-$, wherein any $C_1-C_6\text{-alkyl}$ moiety is optionally substituted with R^{38} .

In another embodiment hereof K is a valence bond, $C_1-C_6\text{-alkylene}$, $-NH-C(=O)-U-$, $-C_1-C_6\text{-alkyl}-S-$, or $-C_1-C_6\text{-alkyl}-O$, wherein any $C_1-C_6\text{-alkyl}$ moiety is optionally substituted with R^{38} .

In another embodiment hereof K is a valence bond, $C_1-C_6\text{-alkylene}$, or $-NH-C(=O)-U$, wherein any $C_1-C_6\text{-alkyl}$ moiety is optionally substituted with R^{38} .

In another embodiment hereof K is a valence bond or $C_1-C_6\text{-alkylene}$, wherein any $C_1-C_6\text{-alkyl}$ moiety is optionally substituted with R^{38} .

In another embodiment hereof K is a valence bond or $-NH-C(=O)-U$.

In another embodiment hereof K is a valence bond.

In another embodiment hereof U is a valence bond or $-C_1-C_6\text{-alkyl}-O-$.

In another embodiment hereof U is a valence bond.

In another embodiment hereof M is arylene or heteroarylene, wherein the arylene or heteroarylene moieties are optionally substituted with one or more substituents independently selected from R^{40} .

In another embodiment hereof M is ArG1 or Het1, wherein the arylene or heteroarylene moieties are optionally substituted with one or more substituents independently selected from R^{40} .


In another embodiment hereof M is ArG1 or Het2, wherein the arylene or heteroarylene moieties are optionally substituted with one or more substituents independently selected from R^{40} .

In another embodiment hereof M is ArG1 or Het3, wherein the arylene or heteroarylene moieties are optionally substituted with one or more substituents independently selected from R^{40} .

In another embodiment hereof M is phenylene optionally substituted with one or more substituents independently selected from R^{40} .


In another embodiment hereof M is indolyne optionally substituted with one or more substituents independently selected from R^{40} .

5 In another embodiment hereof M is



In another embodiment hereof M is carbazolyne optionally substituted with one or more substituents independently selected from R^{40} .

In another embodiment hereof M is



10

In another embodiment hereof R^{40} is selected from

- hydrogen, halogen, -CN, -CF₃, -OCF₃, -NO₂, -OR⁴¹, -NR⁴¹R⁴², -SR⁴¹, -S(O)₂R⁴¹, -NR⁴¹C(O)R⁴², -OC₁-C₆-alkyl-C(O)NR⁴¹R⁴², -C₂-C₆-alkenyl-C(=O)OR⁴¹, -C(O)OR⁴¹, =O, -NH-C(=O)-O-C₁-C₆-alkyl, or -NH-C(=O)-C(=O)-O-C₁-C₆-alkyl,

15

C₁-C₆-alkyl or C₂-C₆- alkenyl which may each optionally be substituted with one or more substituents independently selected from R^{43} ,

- aryl, aryloxy, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, or heteroaryl-C₂-C₆-alkenyl, wherein the cyclic moieties optionally may be substituted with one or more substituents selected from R^{44} .

20

In another embodiment hereof R^{40} is selected from

- hydrogen, halogen, -CN, -CF₃, -OCF₃, -NO₂, -OR⁴¹, -NR⁴¹R⁴², -SR⁴¹, -S(O)₂R⁴¹, -NR⁴¹C(O)R⁴², -OC₁-C₆-alkyl-C(O)NR⁴¹R⁴², -C₂-C₆-alkenyl-C(=O)OR⁴¹, -C(O)OR⁴¹, =O, -NH-C(=O)-O-C₁-C₆-alkyl, or -NH-C(=O)-C(=O)-O-C₁-C₆-alkyl,

25

C₁-C₆-alkyl or C₂-C₆- alkenyl which may each optionally be substituted with one or more substituents independently selected from R^{43} ,

• ArG1, ArG1-O-, ArG1-C₁-C₆-alkoxy, ArG1-C₁-C₆-alkyl, ArG1-C₂-C₆-alkenyl, Het3, Het3-C₁-C₆-alkyl, or Het3-C₂-C₆-alkenyl, wherein the cyclic moieties optionally may be substituted with one or more substituents selected from R⁴⁴.

In another embodiment hereof R⁴⁰ is selected from

- 5 • hydrogen, halogen, -CF₃, -NO₂, -OR⁴¹, -NR⁴¹R⁴², -C(O)OR⁴¹, =O, or -NR⁴¹C(O)R⁴²,
 • C₁-C₆-alkyl,
 • ArG1.

In another embodiment hereof R⁴⁰ is hydrogen.

10 In another embodiment hereof R⁴⁰ is selected from

- Halogen, -NO₂, -OR⁴¹, -NR⁴¹R⁴², -C(O)OR⁴¹, or -NR⁴¹C(O)R⁴²,
 • Methyl,
 • Phenyl.

15 In another embodiment hereof R⁴¹ and R⁴² are independently selected from hydrogen, C₁-C₆-alkyl, or aryl, wherein the aryl moieties may optionally be substituted with halogen or -COOH.

In another embodiment hereof R⁴¹ and R⁴² are independently selected from hydrogen, methyl, ethyl, or phenyl, wherein the phenyl moieties may optionally be substituted with halogen or -COOH.

20 In another embodiment hereof Q is a valence bond, C₁-C₆-alkylene, -C₁-C₆-alkyl-O-, -C₁-C₆-alkyl-NH-, -NH-C₁-C₆-alkyl, -NH-C(=O)-, -C(=O)-NH-, -O-C₁-C₆-alkyl, -C(=O)-, or -C₁-C₆-alkyl-C(=O)-N(R⁴⁷)- wherein the alkyl moieties are optionally substituted with one or more substituents independently selected from R⁴⁸.

25 In another embodiment hereof Q is a valence bond, -CH₂-, -CH₂-CH₂-, -CH₂-O-, -CH₂-CH₂-O-, -CH₂-NH-, -CH₂-CH₂-NH-, -NH-CH₂-, -NH-CH₂-CH₂-, -NH-C(=O)-, -C(=O)-NH-, -O-CH₂-, -O-CH₂-CH₂-, or -C(=O)-.

In another embodiment hereof R⁴⁷ and R⁴⁸ are independently selected from hydrogen, methyl and phenyl.

In another embodiment hereof T is

- 30 • Hydrogen,
 • C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R⁵⁰,
 • aryl, aryl-C₁-C₆-alkyl, heteroaryl, wherein the alkyl, aryl and heteroaryl moieties are optionally substituted with one or more substituents independently selected from R⁵⁰.

35 In another embodiment hereof T is

- hydrogen,
 - C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R⁵⁰,
 - ArG1, ArG1-C₁-C₆-alkyl, Het3, wherein the alkyl, aryl and heteroaryl moieties are
- 5 optionally substituted with one or more substituents independently selected from R⁵⁰.

In another embodiment hereof T is

- hydrogen,
 - C₁-C₆-alkyl, optionally substituted with one or more substituents independently
- 10 selected from R⁵⁰,
- phenyl, phenyl-C₁-C₆-alkyl, wherein the alkyl and phenyl moieties are optionally substituted with one or more substituents independently selected from R⁵⁰.

In another embodiment hereof T is phenyl substituted with R⁵⁰.

- 15 In another embodiment hereof R⁵⁰ is C₁-C₆-alkyl, C₁-C₆-alkoxy, aryl, aryloxy, aryl-C₁-C₆-alkoxy, -C(=O)-NH-C₁-C₆-alkyl-aryl, -C(=O)-NR^{50A}-C₁-C₆-alkyl, -C(=O)-NH-(CH₂CH₂O)_mC₁-C₆-alkyl-COOH, heteroaryl, -C₁-C₆-alkyl-COOH, -O-C₁-C₆-alkyl-COOH, -S(O)₂R⁵¹, -C₂-C₆-alkenyl-COOH, -OR⁵¹, -NO₂, halogen, -COOH, -CF₃, -CN, =O, -N(R⁵¹R⁵²), wherein the aryl or heteroaryl moieties are optionally substituted with one or more R⁵³.

- 20 In another embodiment hereof R⁵⁰ is C₁-C₆-alkyl, C₁-C₆-alkoxy, aryl, aryloxy, -C(=O)-NR^{50A}-C₁-C₆-alkyl, -C(=O)-NH-(CH₂CH₂O)_mC₁-C₆-alkyl-COOH, aryl-C₁-C₆-alkoxy, -OR⁵¹, -NO₂, halogen, -COOH, -CF₃, wherein any aryl moiety is optionally substituted with one or more R⁵³.

- In another embodiment hereof R⁵⁰ is C₁-C₆-alkyl, aryloxy, -C(=O)-NR^{50A}-C₁-C₆-alkyl,
- 25 -C(=O)-NH-(CH₂CH₂O)_mC₁-C₆-alkyl-COOH, aryl-C₁-C₆-alkoxy, -OR⁵¹, halogen, -COOH, -CF₃, wherein any aryl moiety is optionally substituted with one or more R⁵³.

In another embodiment hereof R⁵⁰ is C₁-C₆-alkyl, ArG1-O-, -C(=O)-NR^{50A}-C₁-C₆-alkyl, -C(=O)-NH-(CH₂CH₂O)_mC₁-C₆-alkyl-COOH, ArG1-C₁-C₆-alkoxy, -OR⁵¹, halogen, -COOH, -CF₃, wherein any aryl moiety is optionally substituted with one or more R⁵³.

- 30 In another embodiment hereof R⁵⁰ is -C(=O)-NR^{50A}CH₂, -C(=O)-NH-(CH₂CH₂O)₂CH₂-COOH, or -C(=O)-NR^{50A}CH₂CH₂.

In another embodiment hereof R⁵⁰ is phenyl, methyl or ethyl.

In another embodiment hereof R⁵⁰ is methyl or ethyl.

In another embodiment hereof m is 1 or 2.


- 35 In another embodiment hereof R⁵¹ is methyl.

In another embodiment hereof R^{53} is C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, $-OR^{51}$, halogen, or $-CF_3$.

In another embodiment hereof R^{50A} is $-C(O)OCH_3$, $-C(O)OCH_2CH_3$, $-COOH$, $-CH_2C(O)OCH_3$, $-CH_2C(O)OCH_2CH_3$, $-CH_2CH_2C(O)OCH_3$, $-CH_2CH_2C(O)OCH_2CH_3$, $-CH_2COOH$, methyl, or ethyl.

- 5 In another embodiment hereof R^{50B} is $-C(O)OCH_3$, $-C(O)OCH_2CH_3$, $-COOH$, $-CH_2C(O)OCH_3$, $-CH_2C(O)OCH_2CH_3$, $-CH_2CH_2C(O)OCH_3$, $-CH_2CH_2C(O)OCH_2CH_3$, $-CH_2COOH$, methyl, or ethyl.

In another embodiment hereof the zinc-binding ligand is



wherein V is C_1 - C_6 -alkyl, aryl, heteroaryl, aryl- C_{1-6} -alkyl- or aryl- C_{2-6} -alkenyl-, wherein the alkyl or alkenyl is optionally substituted with one or more substituents independently selected from R^{54} , and the aryl or heteroaryl is optionally substituted with one or more substituents independently selected from R^{55} ,

15

R^{54} is independently selected from halogen, $-CN$, $-CF_3$, $-OCF_3$, aryl, $-COOH$ and $-NH_2$,

R^{55} is independently selected from

- 20 • hydrogen, halogen, $-CN$, $-CH_2CN$, $-CHF_2$, $-CF_3$, $-OCF_3$, $-OCHF_2$, $-OCH_2CF_3$, $-OCF_2CHF_2$, $-S(O)_2CF_3$, $-OS(O)_2CF_3$, $-SCF_3$, $-NO_2$, $-OR^{56}$, $-NR^{56}R^{57}$, $-SR^{56}$, $-NR^{56}S(O)_2R^{57}$, $-S(O)_2NR^{56}R^{57}$, $-S(O)NR^{56}R^{57}$, $-S(O)R^{56}$, $-S(O)_2R^{56}$, $-OS(O)_2R^{56}$, $-C(O)NR^{56}R^{57}$, $-OC(O)NR^{56}R^{57}$, $-NR^{56}C(O)R^{57}$, $-CH_2C(O)NR^{56}R^{57}$, $-OC_1-C_6$ -alkyl- $C(O)NR^{56}R^{57}$, $-CH_2OR^{56}$, $-CH_2OC(O)R^{56}$, $-CH_2NR^{56}R^{57}$, $-OC(O)R^{56}$, $-OC_1-C_8$ -alkyl- $C(O)OR^{56}$, $-OC_1-C_6$ -alkyl- OR^{56} , $-SC_1-C_6$ -alkyl- $C(O)OR^{56}$, $-C_2-C_6$ -alkenyl- $C(=O)OR^{56}$, $-NR^{56}-C(=O)-C_1-C_6$ -alkyl- $C(=O)OR^{56}$, $-NR^{56}-C(=O)-C_1-C_6$ -alkenyl- $C(=O)OR^{56}$, $-C(O)OR^{56}$, or $-C_2-C_6$ -alkenyl- $C(=O)R^{56}$,
- 25

- C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl or C_2 - C_6 -alkynyl,

- 30 which may optionally be substituted with one or more substituents selected from R^{58} ,

• aryl, aryloxy, aryloxycarbonyl, aroyl, arylsulfanyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aroyl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl or heteroaryl-C₂-C₆-alkynyl,

- 5 of which the cyclic moieties optionally may be substituted with one or more substituents selected from R⁵⁹,

R⁵⁶ and R⁵⁷ are independently selected from hydrogen, OH, CF₃, C₁-C₁₂-alkyl, aryl-C₁-C₆-alkyl, -C(=O)-C₁-C₆-alkyl or aryl, wherein the alkyl groups may optionally be substituted with one or more substituents independently selected from R⁶⁰, and the aryl groups may optionally be substituted with one or more substituents independently selected from R⁶¹; R⁵⁶ and R⁵⁷ when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom, the heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds,

R⁵⁸ is independently selected from halogen, -CN, -CF₃, -OCF₃, -OR⁵⁶, and -NR⁵⁶R⁵⁷,

R⁵⁹ is independently selected from halogen, -C(O)OR⁵⁶, -CH₂C(O)OR⁵⁶, -CH₂OR⁵⁶, -CN, -CF₃, -OCF₃, -NO₂, -OR⁵⁶, -NR⁵⁶R⁵⁷ and C₁-C₆-alkyl,

R⁶⁰ is independently selected from halogen, -CN, -CF₃, -OCF₃, -OC₁-C₆-alkyl, -C(O)OC₁-C₆-alkyl, -C(=O)-R⁶², -COOH and -NH₂,

R⁶¹ is independently selected from halogen, -C(O)OC₁-C₆-alkyl, -COOH, -CN, -CF₃, -OCF₃, -NO₂, -OH, -OC₁-C₆-alkyl, -NH₂, C(=O) or C₁-C₆-alkyl,

R⁶² is C₁-C₆-alkyl, aryl optionally substituted with one or more substituents independently selected from halogen, or heteroaryl optionally substituted with one or more C₁-C₆-alkyl independently,

or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.

In another embodiment hereof V is aryl, heteroaryl, or aryl-C₁₋₆-alkyl-, wherein the alkyl is optionally substituted with one or more substituents independently selected R⁵⁴, and the aryl

or heteroaryl is optionally substituted with one or more substituents independently selected from R⁵⁵.

In another embodiment hereof V is aryl, Het1, or aryl-C₁₋₆-alkyl-, wherein the alkyl is optionally substituted with one or more substituents independently selected from R⁵⁴, and the
5 aryl or heteroaryl moiety is optionally substituted with one or more substituents independently selected from R⁵⁵.

In another embodiment hereof V is aryl, Het2, or aryl-C₁₋₆-alkyl-, wherein the alkyl is optionally substituted with one or more substituents independently selected from R⁵⁴, and the
10 aryl or heteroaryl moiety is optionally substituted with one or more substituents independently selected from R⁵⁵.

In another embodiment hereof V is aryl, Het3, or aryl-C₁₋₆-alkyl-, wherein the alkyl is optionally substituted with one or more substituents independently selected from R⁵⁴, and the
aryl or heteroaryl moiety is optionally substituted with one or more substituents independently selected from R⁵⁵.

15 In another embodiment hereof V is aryl optionally substituted with one or more substituents independently selected from R⁵⁵.

In another embodiment hereof V is ArG1 optionally substituted with one or more substituents independently selected from R⁵⁵.

In another embodiment hereof V is phenyl, naphthyl or anthranyl optionally substituted with
20 one or more substituents independently selected from R⁵⁵.

In another embodiment hereof V is phenyl optionally substituted with one or more substituents independently selected from R⁵⁵.

In another embodiment hereof R⁵⁵ is independently selected from

- 25
- halogen, C₁-C₆-alkyl, -CN, -OCF₃, -CF₃, -NO₂, -OR⁵⁶, -NR⁵⁶R⁵⁷, -NR⁵⁶C(O)R⁵⁷-SR⁵⁶, -OC₁-C₈-alkyl-C(O)OR⁵⁶, or -C(O)OR⁵⁶,
 - C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R⁵⁸
 - aryl, aryl-C₁-C₆-alkyl, heteroaryl, or heteroaryl-C₁-C₆-alkyl
- 30 of which the cyclic moieties optionally may be substituted with one or more substituents independently selected from R⁵⁹.

In another embodiment hereof R⁵⁵ is independently selected from

- 35
- halogen, C₁-C₆-alkyl, -CN, -OCF₃, -CF₃, -NO₂, -OR⁵⁶, -NR⁵⁶R⁵⁷, -NR⁵⁶C(O)R⁵⁷-SR⁵⁶, -OC₁-C₈-alkyl-C(O)OR⁵⁶, or -C(O)OR⁵⁶
 - C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R⁵⁸

• ArG1, ArG1-C₁-C₆-alkyl, Het3, or Het3-C₁-C₆-alkyl

of which the cyclic moieties optionally may be substituted with one or more substituents independently selected from R⁵⁹.

5 In another embodiment hereof R⁵⁵ is independently selected from halogen, -OR⁵⁶, -NR⁵⁶R⁵⁷, -C(O)OR⁵⁶, -OC₁-C₈-alkyl-C(O)OR⁵⁶, -NR⁵⁶C(O)R⁵⁷ or C₁-C₆-alkyl.


In another embodiment hereof R⁵⁵ is independently selected from halogen, -OR⁵⁶, -NR⁵⁶R⁵⁷, -C(O)OR⁵⁶, -OC₁-C₈-alkyl-C(O)OR⁵⁶, -NR⁵⁶C(O)R⁵⁷, methyl or ethyl.

10 In another embodiment hereof R⁵⁶ and R⁵⁷ are independently selected from hydrogen, CF₃, C₁-C₁₂-alkyl, or -C(=O)-C₁-C₆-alkyl; R⁵⁶ and R⁵⁷ when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom.

In another embodiment hereof R⁵⁶ and R⁵⁷ are independently selected from hydrogen or C₁-C₁₂-alkyl, R⁵⁶ and R⁵⁷ when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom.

15 In another embodiment hereof R⁵⁶ and R⁵⁷ are independently selected from hydrogen or methyl, ethyl, propyl butyl, R⁵⁶ and R⁵⁷ when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom.

In another embodiment hereof the zinc-binding ligand is



wherein AA is C₁-C₆-alkyl, aryl, heteroaryl, aryl-C₁₋₈-alkyl- or aryl-C₂₋₆-alkenyl-, wherein the alkyl or alkenyl is optionally substituted with one or more substituents independently selected from R⁶³, and the aryl or heteroaryl is optionally substituted with one or more substituents

25 independently selected from R⁶⁴,

R⁶³ is independently selected from halogen, -CN, -CF₃, -OCF₃, aryl, -COOH and -NH₂,

R⁶⁴ is independently selected from

30

• hydrogen, halogen, -CN, -CH₂CN, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -S(O)₂CF₃, -OS(O)₂CF₃, -SCF₃, -NO₂, -OR⁶⁵, -NR⁶⁵R⁶⁶, -SR⁶⁵,

$-\text{NR}^{65}\text{S}(\text{O})_2\text{R}^{66}$, $-\text{S}(\text{O})_2\text{NR}^{65}\text{R}^{66}$, $-\text{S}(\text{O})\text{NR}^{65}\text{R}^{66}$, $-\text{S}(\text{O})\text{R}^{65}$, $-\text{S}(\text{O})_2\text{R}^{65}$, $-\text{OS}(\text{O})_2\text{R}^{65}$,
 $-\text{C}(\text{O})\text{NR}^{65}\text{R}^{66}$, $-\text{OC}(\text{O})\text{NR}^{65}\text{R}^{66}$, $-\text{NR}^{65}\text{C}(\text{O})\text{R}^{66}$, $-\text{CH}_2\text{C}(\text{O})\text{NR}^{65}\text{R}^{66}$, $-\text{OC}_1\text{-C}_6\text{-}$
 $\text{alkyl-C}(\text{O})\text{NR}^{65}\text{R}^{66}$, $-\text{CH}_2\text{OR}^{65}$, $-\text{CH}_2\text{OC}(\text{O})\text{R}^{65}$, $-\text{CH}_2\text{NR}^{65}\text{R}^{66}$, $-\text{OC}(\text{O})\text{R}^{65}$, $-\text{OC}_1\text{-C}_6\text{-}$
 $\text{alkyl-C}(\text{O})\text{OR}^{65}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-OR}^{65}$, $-\text{SC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{65}$, $-\text{C}_2\text{-C}_6\text{-alkenyl-}$
 $\text{C}(=\text{O})\text{OR}^{65}$, $-\text{NR}^{65}\text{-C}(=\text{O})\text{-C}_1\text{-C}_6\text{-alkyl-C}(=\text{O})\text{OR}^{65}$, $-\text{NR}^{65}\text{-C}(=\text{O})\text{-C}_1\text{-C}_6\text{-}$
 $\text{alkenyl-C}(=\text{O})\text{OR}^{65}$, $-\text{C}(\text{O})\text{OR}^{65}$, or $-\text{C}_2\text{-C}_6\text{-alkenyl-C}(=\text{O})\text{R}^{65}$,

• $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_2\text{-C}_6\text{-alkenyl}$ or $\text{C}_2\text{-C}_6\text{-alkynyl}$, each of which may optionally be substituted with one or more substituents selected from R^{67} ,

• aryl, aryloxy, aryloxycarbonyl, aroyl, arylsulfanyl, aryl- $\text{C}_1\text{-C}_6\text{-alkoxy}$, aryl- $\text{C}_1\text{-C}_6\text{-alkyl}$, aryl- $\text{C}_2\text{-C}_6\text{-alkenyl}$, aroyl- $\text{C}_2\text{-C}_6\text{-alkenyl}$, aryl- $\text{C}_2\text{-C}_6\text{-alkynyl}$, heteroaryl, heteroaryl- $\text{C}_1\text{-C}_6\text{-alkyl}$, heteroaryl- $\text{C}_2\text{-C}_6\text{-alkenyl}$ or heteroaryl- $\text{C}_2\text{-C}_6\text{-alkynyl}$,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R^{68} ,

R^{65} and R^{66} are independently selected from hydrogen, OH, CF_3 , $\text{C}_1\text{-C}_{12}\text{-alkyl}$, aryl- $\text{C}_1\text{-C}_6\text{-alkyl}$, $-\text{C}(=\text{O})\text{-R}^{69}$, aryl or heteroaryl, wherein the alkyl groups may optionally be substituted with one or more substituents selected from R^{70} , and the aryl and heteroaryl groups may optionally be substituted with one or more substituents independently selected from R^{71} ; R^{65} and R^{66} when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom, the heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds,

R^{67} is independently selected from halogen, $-\text{CN}$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OR}^{65}$, and $-\text{NR}^{65}\text{R}^{66}$,

R^{68} is independently selected from halogen, $-\text{C}(\text{O})\text{OR}^{65}$, $-\text{CH}_2\text{C}(\text{O})\text{OR}^{65}$, $-\text{CH}_2\text{OR}^{65}$, $-\text{CN}$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{NO}_2$, $-\text{OR}^{65}$, $-\text{NR}^{65}\text{R}^{66}$ and $\text{C}_1\text{-C}_6\text{-alkyl}$,

R^{69} is independently selected from $\text{C}_1\text{-C}_6\text{-alkyl}$, aryl optionally substituted with one or more halogen, or heteroaryl optionally substituted with one or more $\text{C}_1\text{-C}_6\text{-alkyl}$,

R⁷⁰ is independently selected from halogen, -CN, -CF₃, -OCF₃, -OC₁-C₆-alkyl, -C(O)OC₁-C₆-alkyl, -COOH and -NH₂,

- 5 R⁷¹ is independently selected from halogen, -C(O)OC₁-C₆-alkyl, -COOH, -CN, -CF₃, -OCF₃, -NO₂, -OH, -OC₁-C₆-alkyl, -NH₂, C(=O) or C₁-C₆-alkyl,

or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.

- 10 In another embodiment hereof AA is aryl, heteroaryl or aryl-C₁₋₆-alkyl-, wherein the alkyl is optionally substituted with one or more R⁶³, and the aryl or heteroaryl is optionally substituted with one or more substituents independently selected from R⁶⁴.

In another embodiment hereof AA is aryl or heteroaryl optionally substituted with one or more substituents independently selected from R⁶⁴.

- 15 In another embodiment hereof AA is ArG1 or Het1 optionally substituted with one or more substituents independently selected from R⁶⁴.

In another embodiment hereof AA is ArG1 or Het2 optionally substituted with one or more substituents independently selected from R⁶⁴.

In another embodiment hereof AA is ArG1 or Het3 optionally substituted with one or more substituents independently selected from R⁶⁴.

- 20 In another embodiment hereof AA is phenyl, naphthyl, anthryl, carbazolyl, thienyl, pyridyl, or benzodioxyl optionally substituted with one or more substituents independently selected from R⁶⁴.

In another embodiment hereof AA is phenyl or naphthyl optionally substituted with one or more substituents independently selected from R⁶⁴.

- 25 In another embodiment hereof R⁶⁴ is independently selected from hydrogen, halogen, -CF₃, -OCF₃, -OR⁶⁵, -NR⁶⁵R⁶⁶, C₁-C₆-alkyl, -OC(O)R⁶⁵, -OC₁-C₆-alkyl-C(O)OR⁶⁵, aryl-C₂-C₆-alkenyl, aryloxy or aryl, wherein C₁-C₆-alkyl is optionally substituted with one or more substituents independently selected from R⁶⁷, and the cyclic moieties optionally are substituted with one or more substituents independently selected from R⁶⁸.

- 30 In another embodiment hereof R⁶⁴ is independently selected from halogen, -CF₃, -OCF₃, -OR⁶⁵, -NR⁶⁵R⁶⁶, methyl, ethyl, propyl, -OC(O)R⁶⁵, -OCH₂-C(O)OR⁶⁵, -OCH₂-CH₂-C(O)OR⁶⁵, phenoxy optionally substituted with one or more substituents independently selected from R⁶⁸.

In another embodiment hereof R⁶⁵ and R⁶⁶ are independently selected from hydrogen, CF₃, C₁-C₁₂-alkyl, aryl, or heteroaryl optionally substituted with one or more substituents independently selected from R⁷¹.

5 In another embodiment hereof R⁶⁵ and R⁶⁶ are independently hydrogen, C₁-C₁₂-alkyl, aryl, or heteroaryl optionally substituted with one or more substituents independently selected from R⁷¹.

In another embodiment hereof R⁶⁵ and R⁶⁶ are independently hydrogen, methyl, ethyl, propyl, butyl, 2,2-dimethyl-propyl, ArG1 or Het1 optionally substituted with one or more substituents independently selected from R⁷¹.

10 In another embodiment hereof R⁶⁵ and R⁶⁶ are independently hydrogen, methyl, ethyl, propyl, butyl, 2,2-dimethyl-propyl, ArG1 or Het2 optionally substituted with one or more substituents independently selected from R⁷¹.

In another embodiment hereof R⁶⁵ and R⁶⁶ are independently hydrogen, methyl, ethyl, propyl, butyl, 2,2-dimethyl-propyl, ArG1 or Het3 optionally substituted with one or more substituents independently selected from R⁷¹.

15 In another embodiment hereof R⁶⁵ and R⁶⁶ are independently hydrogen, methyl, ethyl, propyl, butyl, 2,2-dimethyl-propyl, phenyl, naphthyl, thiadiazolyl optionally substituted with one or more R⁷¹ independently; or isoxazolyl optionally substituted with one or more substituents independently selected from R⁷¹.

20 In another embodiment hereof R⁷¹ is halogen or C₁-C₆-alkyl.

In another embodiment hereof R⁷¹ is halogen or methyl.

The following aspects are also provided by the present invention, wherein the compounds of the invention may be any of the above described embodiments.

25

In one aspect the invention provides a pharmaceutical composition wherein the insulin is rapid acting insulin.

30 In another embodiment the invention provides a pharmaceutical composition wherein the insulin is selected from the group consisting of human insulin, an analogue thereof, a derivative thereof, and combinations of any of these.

In another embodiment the invention provides a pharmaceutical composition wherein the insulin is an analogue of human insulin selected from the group consisting of

i. An analogue wherein position B28 is Asp, Lys, Leu, Val, or Ala and position B29 is Lys or Pro; and

35 ii. des(B28-B30), des(B27) or des(B30) human insulin.

In another embodiment the invention provides a pharmaceutical composition wherein the insulin is an analogue of human insulin wherein position B28 is Asp or Lys, and position B29 is Lys or Pro.

- 5 In another embodiment the invention provides a pharmaceutical composition wherein the insulin is des(B30) human insulin.

In another embodiment the invention provides a pharmaceutical composition wherein the insulin is an analogue of human insulin wherein position B3 is Lys and position B29 is Glu or Asp.

- 10 In another embodiment the invention provides a pharmaceutical composition wherein the insulin is a derivative of human insulin having one or more lipophilic substituents.

In another embodiment the invention provides a pharmaceutical composition wherein the insulin derivative is selected from the group consisting of B29-N^E-myristoyl-des(B30) human insulin, B29-N^E-palmitoyl-des(B30) human insulin, B29-N^E-myristoyl human insulin, B29-N^E-palmitoyl human insulin, B28-N^E-myristoyl Lys^{B28} Pro^{B29} human insulin, B28-N^E-palmitoyl Lys^{B28} Pro^{B29} human insulin, B30-N^E-myristoyl-Thr^{B29}Lys^{B30} human insulin, B30-N^E-palmitoyl-Thr^{B29}Lys^{B30} human insulin, B29-N^E-(N-palmitoyl-γ-glutamyl)-des(B30) human insulin, B29-N^E-(N-lithocholyl-γ-glutamyl)-des(B30) human insulin, B29-N^E-(ω-carboxyheptadecanoyl)-des(B30) human insulin and B29-N^E-(ω-carboxyheptadecanoyl) human insulin.

- 20 In another embodiment the invention provides a pharmaceutical composition wherein the insulin derivative is B29-N^E-myristoyl-des(B30) human insulin.

In another embodiment the invention provides a pharmaceutical composition comprising 2-6 moles zinc²⁺ ions per mole insulin.

- 25 In another embodiment the invention provides a pharmaceutical composition comprising 2-3 moles zinc²⁺ ions per mole insulin.

In another embodiment the invention provides a pharmaceutical composition further comprising at least 3 molecules of a phenolic compound per insulin hexamer.

- 30 In another embodiment the invention provides a pharmaceutical composition further comprising an isotonicity agent.

In another embodiment the invention provides a pharmaceutical composition further comprising a buffer substance.

A method of stabilising an insulin composition comprising adding a zinc-binding ligand to the insulin composition.

A method of treating type 1 or type 2 diabetes comprising administering to a patient in need thereof a pharmaceutically effective dose of an insulin composition.

5 In one embodiment of the invention the concentration of added ligand for the zinc site is between 0.2 and 10 times that of zinc ion in the preparation. In another embodiment the concentration is between 0.5 and 5 times that of zinc ion. In another embodiment the ligand concentration is identical to that of zinc ion in the preparation.

10 The compounds of the present invention may be chiral, and it is intended that any enantiomers, as separated, pure or partially purified enantiomers or racemic mixtures thereof are included within the scope of the invention.

Furthermore, when a double bond or a fully or partially saturated ring system or more than one centre of asymmetry or a bond with restricted rotatability is present in the molecule diastereomers may be formed. It is intended that any diastereomers, as separated, pure or
15 partially purified diastereomers or mixtures thereof are included within the scope of the invention.

Furthermore, some of the compounds of the present invention may exist in different tautomeric forms and it is intended that any tautomeric forms, which the compounds are able to form, are included within the scope of the present invention.

20 The present invention also encompasses pharmaceutically acceptable salts of the present compounds. Such salts include pharmaceutically acceptable acid addition salts, pharmaceutically acceptable metal salts, ammonium and alkylated ammonium salts. Acid addition salts include salts of inorganic acids as well as organic acids. Representative examples of suitable inorganic acids include hydrochloric, hydrobromic, hydroiodic, phosphoric, sulphuric,
25 nitric acids and the like. Representative examples of suitable organic acids include formic, acetic, trichloroacetic, trifluoroacetic, propionic, benzoic, cinnamic, citric, fumaric, glycolic, lactic, maleic, malic, malonic, mandelic, picric, pyruvic, succinic, methanesulfonic, ethanesulfonic, tartaric, ascorbic, pantoic, , ethanedisulfonic, gluconic, citraconic, aspartic, stearic, palmitic, glycolic, p-aminobenzoic, glutamic, benzenesulfonic, p-toluenesulfonic acids
30 and the like. Further examples of pharmaceutically acceptable inorganic or organic acid addition salts include the pharmaceutically acceptable salts listed in J. Pharm. Sci. 1977, 66, 2, which is incorporated herein by reference. Examples of metal salts include lithium, sodium, potassium, magnesium salts and the like. Examples of ammonium and alkylated ammonium salts include ammonium, methyl-, dimethyl-, trimethyl-, ethyl-, hydroxyethyl-, diethyl-,
35 n-butyl-, sec-butyl-, tert-butyl-, tetramethylammonium salts and the like.

Also intended as pharmaceutically acceptable acid addition salts are the hydrates, which the present compounds, are able to form.

Furthermore, the pharmaceutically acceptable salts comprise basic amino acid salts such as lysine, arginine and ornithine.

- 5 The acid addition salts may be obtained as the direct products of compound synthesis. In the alternative, the free base may be dissolved in a suitable solvent containing the appropriate acid, and the salt isolated by evaporating the solvent or otherwise separating the salt and solvent.

- 10 The compounds of the present invention may form solvates with standard low molecular weight solvents using methods well known to the person skilled in the art. Such solvates are also contemplated as being within the scope of the present invention.

- 15 In one embodiment of the invention the stabilized preparations are used in connection with insulin pumps. The insulin pumps may be prefilled and disposable, or the insulin compositions may be supplied from a reservoir which is removable. Insulin pumps may be skin-mounted or carried, and the path of the insulin composition from the storage compartment of the pump to the patient may be more or less tortuous. The elevated temperature and increased physical stress the insulin composition is thus exposed to challenges the stability of the constituent insulin. Non-limiting examples of insulin pumps are
20 disclosed in US 5,957,895, US 5,858,001, US 4,468,221, US 4,468,221, US 5,957,895, US 5,858,001, US 6,074,369, US 5,858,001, US 5,527,288, and US 6,074,369.

- In another embodiment the stabilized preparations are used in connection with pen-like injection devices, which may be prefilled and disposable, or the insulin compositions may be
25 supplied from a reservoir which is removable. Non-limiting examples of pen-like injection devices are FlexPen®, InnoLet®, InDuo™, Innovo®.

- In a further embodiment stabilized preparations are used in connection with devices for pulmonary administration of aqueous insulin compositions, a non-limiting example of which is
30 the AerX® device.

- In one aspect of the invention, the ligands are added to rapid acting insulin. The resulting preparations have improved physical and chemical stability while still retaining a high rate of absorbtion from subcutaneous tissue.

The present invention also relates to pharmaceutical compositions for the treatment of diabetes in a patient in need of such a treatment comprising an R-state hexamer of insulin according to the invention together with a pharmaceutically acceptable carrier.

In one embodiment of the invention the insulin composition comprises 60 to 3000 nmol/ml of insulin.

In another embodiment of the invention the insulin composition comprises 240 to 1200 nmol/ml of insulin.

In another embodiment of the invention the insulin composition comprises about 600 nmol/ml of insulin.

10 Zinc ions may be present in an amount corresponding to 13 to 33 μg Zn/100 U insulin, more preferably 15 to 26 μg Zn/100 U insulin.

Insulin formulations of the invention are usually administered from multi-dose containers where a preservative effect is desired. Since phenolic preservatives also stabilize the R-state hexamer the formulations may contain up to 50 mM of phenolic molecules. Non-limiting
15 examples of phenolic molecules are phenol, m-cresol, chloro-cresol, thymol, 7-hydroxyindole or any mixture thereof.

In one embodiment of the invention 0.5 to 4.0 mg/ml of phenolic compound may be employed.

In another embodiment of the invention 0.6 to 4.0 mg/ml of m-cresol may be employed.

In another embodiment of the invention 0.5 to 4.0 mg/ml of phenol may be employed.

20 In another embodiment of the invention 1.4 to 4.0 mg/ml of phenol may be employed.

In another embodiment of the invention 0.5 to 4.0 mg/ml of a mixture of m-cresol or phenol may be employed.

In another embodiment of the invention 1.4 to 4.0 mg/ml of a mixture of m-cresol or phenol may be employed.

25 The pharmaceutical composition may further comprise a buffer substance, such as a TRIS, phosphate, glycine or glycyglycine (or another zwitterionic substance) buffer, an isotonicity agent, such as NaCl, glycerol, mannitol and/or lactose. Chloride would be used at moderate concentrations, in one embodiment of the invention up to 50 mM to avoid competition with the zinc-site ligands of the present invention. In another embodiment the chloride
30 concentration would be from 3 to 20 mM.

The *in vivo* action of insulin may be modified by the addition of physiologically acceptable agents that increase the viscosity of the pharmaceutical composition. Thus, the pharmaceutical composition according to the invention may furthermore comprise an agent which increases the viscosity, such as polyethylene glycol, polypropylene glycol, copolymers
35 thereof, dextrans and/or polylactides.

In one embodiment the insulin composition of the invention comprises between 0.0005 % by weight and 1 % by weight of a non-ionic or zwitter-ionic surfactant, for example tween 20 or Polox 188. A nonionic detergent can be added to stabilise insulin against fibrillation during storage and handling.

- 5 The insulin composition of the present invention may have a pH value in the range of 3.0 to 8.5, e.g. 7.4 to 7.9.

EXAMPLES

10

The following examples and general procedures refer to intermediate compounds and final products identified in the specification and in the synthesis schemes. The preparation of the compounds of the present invention is described in detail using the following examples, but the chemical reactions described are disclosed in terms of their general applicability to the preparation of compounds of the invention. Occasionally, the reaction may not be applicable as described to each compound included within the disclosed scope of the invention. The compounds for which this occurs will be readily recognised by those skilled in the art. In these cases the reactions can be successfully performed by conventional modifications known to those skilled in the art, that is, by appropriate protection of interfering groups, by changing to other conventional reagents, or by routine modification of reaction conditions. Alternatively, other reactions disclosed herein or otherwise conventional will be applicable to the preparation of the corresponding compounds of the invention. In all preparative methods, all starting materials are known or may easily be prepared from known starting materials. All temperatures are set forth in degrees Celsius and unless otherwise indicated, all parts and percentages are by weight when referring to yields and all parts are by volume when referring to solvents and eluents.

15

20

25

HPLC-MS (Method A)

The following instrumentation was used:

30

- Hewlett Packard series 1100 G1312A Bin Pump
- Hewlett Packard series 1100 Column compartment
- Hewlett Packard series 1100 G13 15A DAD diode array detector
- Hewlett Packard series 1100 MSD

The instrument was controlled by HP Chemstation software.

The HPLC pump was connected to two eluent reservoirs containing:

- A: 0.01% TFA in water
B: 0.01% TFA in acetonitrile

5

The analysis was performed at 40 °C by injecting an appropriate volume of the sample (preferably 1 µL) onto the column, which was eluted with a gradient of acetonitrile.

The HPLC conditions, detector settings and mass spectrometer settings used are given in the following table.

10

| | |
|-----------|--|
| Column | Waters Xterra MS C-18 X 3 mm id |
| Gradient | 10% - 100% acetonitrile lineary during 7.5 min at 1.0 mL/min |
| Detection | UV: 210 nm (analog output from DAD) |
| MS | Ionisation mode: API-ES Scan 100-1000 amu step 0.1 amu |

HPLC-MS (Method B)

The following instrumentation was used:

Sciex API 100 Single quadropole mass spectrometer

15 Perkin Elmer Series 200 Quard pump

Perkin Elmer Series 200 autosampler

Applied Biosystems 785A UV detector

Sedex 55 evaporative light scattering detector

A Valco column switch with a Valco actuator controlled by timed events from the pump.

20

The Sciex Sample control software running on a Macintosh PowerPC 7200 computer was used for the instrument control and data acquisition.

The HPLC pump was connected to four eluent reservoirs containing:

- A: Acetonitrile
B: Water

C: 0.5% TFA in water
D: 0.02 M ammonium acetate

The requirements for samples are that they contain approximately 500 µg/mL of the compound to be analysed in an acceptable solvent such as methanol, ethanol, acetonitrile, THF, water and mixtures thereof. (High concentrations of strongly eluting solvents will interfere with the chromatography at low acetonitrile concentrations.)

The analysis was performed at room temperature by injecting 20 µL of the sample solution on the column, which was eluted with a gradient of acetonitrile in either 0.05% TFA or 0.002 M ammonium acetate. Depending on the analysis method varying elution conditions were used.

The eluate from the column was passed through a flow splitting T-connector, which passed approximately 20 µL/min through approx. 1 m. 75 µ fused silica capillary to the API interface of API 100 spectrometer.

The remaining 1.48 mL/min was passed through the UV detector and to the ELS detector.

During the LC-analysis the detection data were acquired concurrently from the mass spectrometer, the UV detector and the ELS detector.

The LC conditions, detector settings and mass spectrometer settings used for the different methods are given in the following table.

| | | | |
|-----------|--|--|------------|
| Column | YMC ODS-A 120Å s - 5µ 3 mm x 50 mm id | | |
| Gradient | 5% - 90% acetonitrile in 0.05% TFA linearly during 7.5 min at 1.5 mL/min | | |
| Detection | UV: 214 nm | | ELS: 40 °C |
| MS | Experiment: Start: 100 amu Stop: 800 amu Step: 0.2 amu | | |
| | Dwell: 0.571 msec | | |
| | Method: Scan 284 times = 9.5 min | | |

HPLC-MS (Method C) The following instrumentation is used:

- Hewlett Packard series 1100 G1312A Bin Pump

- Hewlett Packard series 1100 Column compartment
- Hewlett Packard series 1100 G1315A DAD diode array detector
- Hewlett Packard series 1100 MSD
- Sedere 75 Evaporative Light Scattering detector

5 The instrument is controlled by HP Chemstation software.

The HPLC pump is connected to two eluent reservoirs containing:

| | |
|---|---------------------------|
| A | 0.01% TFA in water |
| B | 0.01% TFA in acetonitrile |

The analysis is performed at 40 °C by injecting an appropriate volume of the sample (preferably 1 μ l) onto the column which is eluted with a gradient of acetonitrile.

10 The HPLC conditions, detector settings and mass spectrometer settings used are given in the following table.

| | |
|-----------|---|
| Column | Waters Xterra MS C-18 X 3 mm id 5 μ m |
| Gradient | 5% - 100% acetonitrile linear during 7.5 min at 1.5 ml/min |
| Detection | 210 nm (analogue output from DAD) ELS (analogue output from ELS) |
| MS | ionisation mode API-ES Scan 100-1000 amu step 0.1 amu |

After the DAD the flow is divided yielding approximately 1 ml/min to the ELS and 0.5 ml/min to the MS.

15

HPLC-MS (Method D)

The following instrumentation was used:

Sciex API 150 Single Quadropole mass spectrometer

Hewlett Packard Series 1100 G1312A Bin pump

20 Gilson 215 micro injector

Hewlett Packard Series 1100 G1315A DAD diode array detector

Sedex 55 evaporative light scattering detector

A Valco column switch with a Valco actuator controlled by timed events from the pump.

The Sciex Sample control software running on a Macintosh Power G3 computer was used

25 for the instrument control and data acquisition.

The HPLC pump was connected to two eluent reservoirs containing:

- A: Acetonitrile containing 0.05% TFA
B: Water containing 0.05% TFA

The requirements for the samples are that they contain approximately 500 µg/ml of the compound to be analysed in an acceptable solvent such as methanol, ethanol, acetonitrile, THF, water and mixtures thereof. (High concentrations of strongly eluting solvents will interfere with the chromatography at low acetonitrile concentrations.)

The analysis was performed at room temperature by injecting 20 µl of the sample solution on the column, which was eluted with a gradient of acetonitrile in 0.05% TFA

The eluate from the column was passed through a flow splitting T-connector, which passed approximately 20 µl/min through approx. 1 m 75 µ fused silica capillary to the API interface of API 150 spectrometer.


The remaining 1.48 ml/min was passed through the UV detector and to the ELS detector. During the LC-analysis the detection data were acquired concurrently from the mass spectrometer, the UV detector and the ELS detector.

The LC conditions, detector settings and mass spectrometer settings used for the different methods are given in the following table.

| | | | |
|-----------|--|--------------------------|--------------------------------|
| Column | Waters X-terra C18 5µ 3 mm x 50 mm id | | |
| Gradient | 5% - 90% acetonitrile in 0.05% TFA linearly during 7.5 min at 1.5 ml/min | | |
| Detection | UV: 214 nm | ELS: 40 °C | |
| MS | Experiment: | Start: 100 amu | Stop: 800 amu Step: 0.2 amu |
| | Dwell: | 0.571 msec | |
| | Method: | Scan 284 times = 9.5 min | |


EXAMPLES

0102-0000-0273Example 1 HBOL

1H-Benzotriazole


5

0102-0000-0274Example 2 HBOL

5,6-Dimethyl-1*H*-benzotriazole


10

0102-0000-0275Example 3 HBOL


1*H*-Benzotriazole-5-carboxylic acid

0102-0000-0280Example 4 HBOL

15


4-Nitro-1*H*-benzotriazole

0102-0000-0284Example 5 HBOL

5-Amino-1*H*-benzotriazole


20

0102-0000-0287Example 6 HBOL


5-Chloro-1*H*-benzotriazole

5

0102-0000-0286Example 7 HBOL

5-Nitro-1*H*-benzotriazole


0102-0000-3015Example 8 PEM

10 4-[(1*H*-Benzotriazole-5-carbonyl)amino]benzoic acid

4-[(1*H*-Benzotriazole-5-carbonyl)amino]benzoic acid methyl ester (5.2 g, 17.6 mmol) was dissolved in THF (60 mL) and methanol (10 mL) was added followed by 1N sodium hydroxide (35 mL). The mixture was stirred at room temperature for 16 hours and then 1N hydrochloric acid (45 mL) was added. The mixture was added water (200 mL) and extracted with ethyl acetate (2 x 500 mL). The combined organic phases were evaporated *in vacuo* to afford 0.44 g of 4-[(1*H*-benzotriazole-5-carbonyl)amino]benzoic acid. By filtration of the aqueous phase a further crop of 4-[(1*H*-benzotriazole-5-carbonyl)amino]benzoic acid was isolated (0.52 g).

20

¹H-NMR (DMSO-*d*₆): δ 7.97 (4H, s), 8.03 (2H, m), 8.66 (1H, bs), 10.7 (1H, s), 12.6 (1H, bs); HPLC-MS (Method A): *m/z*: 283 (M+1); *R*_t = 1.85 min.

General procedure (A) for preparation of compounds of general formula I₁:

wherein D, E and R^{19} are as defined above, and E is optionally substituted with up to three substituents R^{21} , R^{22} and R^{23} independently as defined above.

5


The carboxylic acid of 1H-benzotriazole-5-carboxylic acid is activated, i.e. the OH functionality is converted into a leaving group L (selected from eg fluorine, chlorine, bromine, iodine, 1-imidazolyl, 1,2,4-triazolyl, 1-benzotriazolyl, 1-(4-aza benzotriazolyl)oxy, pentafluorophenoxy, N-succinyloxy, 3,4-dihydro-4-oxo-3-(1,2,3-benzotriazinyl)oxy, benzotriazole 5-COO, or any other leaving group known to act as a leaving group in acylation reactions. The activated benzotriazole-5-carboxylic acid is then reacted with $\text{R}^2-(\text{CH}_2)_n-\text{B}'$ in the presence of a base. The base can be either absent (i.e. $\text{R}^2-(\text{CH}_2)_n-\text{B}'$ acts as a base) or triethylamine, N-ethyl-N,N-diisopropylamine, N-methylmorpholine, 2,6-lutidine, 2,2,6,6-tetramethylpiperidine, potassium carbonate, sodium carbonate, caesium carbonate or any other base known to be useful in acylation reactions. The reaction is performed in a solvent solvent such as THF, dioxane, toluene, dichloromethane, DMF, NMP or a mixture of two or more of these. The reaction is performed between 0 °C and 80 °C, preferably between 20 °C and 40 °C. When the acylation is complete, the product is isolated by extraction, filtration, chromatography or other methods known to those skilled in the art.

20

The general procedure (A) is further illustrated in the following example:

0102-0000-1020Example 9 (General Procedure (A))PEM

1H-Benzotriazole-5-carboxylic acid phenylamide



25 Benzotriazole-5-carboxylic acid (856 mg), HOAt (715 mg) and EDAC (1.00 g) were dissolved in DMF (17.5 mL) and the mixture was stirred at room temperature 1 hour. A 0.5 mL aliquot of this mixture was added to aniline (13.7 μL , 0.15 mmol) and the resulting mixture was vigorously shaken at room temperature for 16 hours. 1N hydrochloric acid (2 mL) and ethyl acetate (1 mL) were added and the mixture was vigorously shaken at room temperature for 2

hours. The organic phase was isolated and concentrated *in vacuo* to afford the title compound.


HPLC-MS (Method B): m/z: 239 (M+1); Rt = 3.93 min.

5

The compounds in the following examples were similarly made. Optionally, the compounds may be isolated by filtration or by chromatography.

0102-0000-1019 Example 10 (General Procedure (A)) PEM

10 1H-Benzotriazole-5-carboxylic acid (4-methoxyphenyl)amide




HPLC-MS (Method A): m/z: 269 (M+1) & 291 (M+23); Rt = 2.41 min

HPLC-MS (Method B): m/z: 239 (M+1); Rt = 3.93 min.

15 0102-0000-1021 Example 11 (General Procedure (A)) PEM


{4-[(1H-Benzotriazole-5-carbonyl)amino]phenyl}carbamic acid *tert*-butyl ester



HPLC-MS (Method B): m/z: 354 (M+1); Rt = 4.58 min.

20 0102-0000-1022 Example 12 (General Procedure (A)) PEM


1H-Benzotriazole-5-carboxylic acid (4-acetylamino)phenyl)amide



HPLC-MS (Method B): m/z: 296 (M+1); Rt = 3.32 min.

25 0102-0000-1023 Example 13 (General Procedure (A)) PEM


1H-Benzotriazole-5-carboxylic acid (3-fluorophenyl)amide



HPLC-MS (Method B): m/z: 257 (M+1); Rt = 4.33 min.

0102-0000-1024 Example 14 (General Procedure (A)) PEM


5 1H-Benzotriazole-5-carboxylic acid (2-chlorophenyl)amide



HPLC-MS (Method B): m/z: 273 (M+1); Rt = 4.18 min.

0102-0000-1025 Example 15 (General Procedure (A)) PEM


10 4-[(1H-Benzotriazole-5-carbonyl)amino]benzoic acid methyl ester



HPLC-MS (Method A): m/z: 297 (M+1); Rt : 2,60 min. HPLC-MS (Method B): m/z: 297 (M+1); Rt = 4.30 min.

0102-0000-1026 Example 16 (General Procedure (A)) PEM


15 1H-Benzotriazole-5-carboxylic acid (4-butylphenyl)amide



HPLC-MS (Method B): m/z: 295 (M+1); Rt = 5.80 min.


0102-0000-1027 Example 17 (General Procedure (A)) PEM

20 1H-Benzotriazole-5-carboxylic acid (1-phenylethyl)amide




HPLC-MS (Method B): m/z: 267 (M+1); Rt = 4.08 min.

0102-0000-1028 Example 18 (General Procedure (A)) PEM

1*H*-Benzotriazole-5-carboxylic acid benzylamideHPLC-MS (Method B): *m/z*: 253 (*M*+1); *Rt* = 3.88 min.


5

0102-0000-1029 Example 19 (General Procedure (A)) PEM

1*H*-Benzotriazole-5-carboxylic acid 4-chlorobenzylamideHPLC-MS (Method B): *m/z*: 287 (*M*+1); *Rt* = 4.40 min.


10

0102-0000-1030 Example 20 (General Procedure (A)) PEM

1*H*-Benzotriazole-5-carboxylic acid 2-chlorobenzylamideHPLC-MS (Method B): *m/z*: 287 (*M*+1); *Rt* = 4.25 min.


15

0102-0000-1031 Example 21 (General Procedure (A)) PEM

1*H*-Benzotriazole-5-carboxylic acid 4-methoxybenzylamideHPLC-MS (Method B): *m/z*: 283 (*M*+1); *Rt* = 3.93 min.


20

0102-0000-1032 Example 22 (General Procedure (A)) PEM

1*H*-Benzotriazole-5-carboxylic acid 3-methoxybenzylamideHPLC-MS (Method B): *m/z*: 283 (*M*+1); *Rt* = 3.97 min.

25


0102-0000-1033 Example 23 (General Procedure (A)) PEM
1*H*-Benzotriazole-5-carboxylic acid (1,2-diphenylethyl)amide



HPLC-MS (Method B): *m/z*: 343 (*M*+1); *R*_t = 5.05 min.

5


0102-0000-1034 Example 24 (General Procedure (A)) PEM
1*H*-Benzotriazole-5-carboxylic acid 3-bromobenzylamide



HPLC-MS (Method B): *m/z*: 331 (*M*+1); *R*_t = 4.45 min.

10


0102-0000-1035 Example 25 (General Procedure (A)) PEM
4-[[[(1*H*-Benzotriazole-5-carbonyl)amino]methyl]benzoic acid



HPLC-MS (Method B): *m/z*: 297 (*M*+1); *R*_t = 3.35 min.

15


0102-0000-1036 Example 26 (General Procedure (A)) PEM
1*H*-Benzotriazole-5-carboxylic acid phenethylamide



HPLC-MS (Method B): *m/z*: 267 (*M*+1); *R*_t = 4.08 min.


20

0102-0000-1037 Example 27 (General Procedure (A)) PEM
1*H*-Benzotriazole-5-carboxylic acid [2-(4-chlorophenyl)ethyl]amide




HPLC-MS (Method B): *m/z*: 301 (*M*+1); *R*_t = 4.50 min.


0102-0000-1038 Example 28 (General Procedure (A)) PEM

1*H*-Benzotriazole-5-carboxylic acid [2-(4-methoxyphenyl)ethyl]amide5 HPLC-MS (Method B): *m/z*: 297 (*M*+1); *R*_t = 4.15 min.


0102-0000-1039 Example 29 (General Procedure (A)) PEM

1*H*-Benzotriazole-5-carboxylic acid [2-(3-methoxyphenyl)ethyl]amide10 HPLC-MS (Method B): *m/z*: 297 (*M*+1); *R*_t = 4.13 min.


0102-0000-1040 Example 30 (General Procedure (A)) PEM

1*H*-Benzotriazole-5-carboxylic acid [2-(3-chlorophenyl)ethyl]amide15 HPLC-MS (Method B): *m/z*: 301 (*M*+1); *R*_t = 4.55 min.

0102-0000-1041 Example 31 (General Procedure (A)) PEM

1*H*-Benzotriazole-5-carboxylic acid (2,2-diphenylethyl)amide20 HPLC-MS (Method B): *m/z*: 343 (*M*+1); *R*_t = 5.00 min.


0102-0000-1042 Example 32 (General Procedure (A)) PEM

1*H*-Benzotriazole-5-carboxylic acid (3,4-dichlorophenyl)methylamide

HPLC-MS (Method B): m/z : 321 ($M+1$); R_t = 4.67 min.

0102-0000-1043 Example 33 (General Procedure (A)) PEM

1*H*-Benzotriazole-5-carboxylic acid methylphenylamide




5

HPLC-MS (Method B): m/z : 253 ($M+1$); R_t = 3.82 min.

0102-0000-1044 Example 34 (General Procedure (A)) PEM

1*H*-Benzotriazole-5-carboxylic acid benzylmethylamide




10

HPLC-MS (Method B): m/z : 267 ($M+1$); R_t = 4.05 min.

0102-0000-1045 Example 35 (General Procedure (A)) PEM

1*H*-Benzotriazole-5-carboxylic acid [2-(3-chloro-4-methoxyphenyl)ethyl]methylamide




15

HPLC-MS (Method B): m/z : 345 ($M+1$); R_t = 4.37 min.

0102-0000-1046 Example 36 (General Procedure (A)) PEM

1*H*-Benzotriazole-5-carboxylic acid methylphenethylamide




20

HPLC-MS (Method B): m/z : 281 ($M+1$); R_t = 4.15 min.

0102-0000-1047 Example 37 (General Procedure (A)) PEM


1*H*-Benzotriazole-5-carboxylic acid [2-(3,4-dimethoxyphenyl)ethyl]methylamide



HPLC-MS (Method B): m/z : 341 (M+1); R_t = 3.78 min;

0102-0000-1048 Example 38 (General Procedure (A)) PEM


5 1H-Benzotriazole-5-carboxylic acid (2-hydroxy-2-phenylethyl)methylamide



HPLC-MS (Method B): m/z : 297 (M+1); R_t = 3.48 min.

Example 39 (General procedure (A))


10 1H-Benzotriazole-5-carboxylic acid (3-bromophenyl)amide



HPLC-MS (Method A): m/z : 317 (M+1); R_t = 3.19 min.

Example 40 (General procedure (A))


15 1H-Benzotriazole-5-carboxylic acid (4-bromophenyl)amide



HPLC-MS (Method A): m/z : 317 (M+1); R_t = 3.18 min.

Example 41 (General procedure (A))


20 {4-[(1H-Benzotriazole-5-carbonyl)amino]benzoylamino}acetic acid



HPLC-MS (Method A): m/z : 340 (M+1); R_t = 1.71 min.

Example 42 (General procedure (A))


4-[(1H-Benzotriazole-5-carbonyl)amino]phenyl}acetic acid



5 HPLC-MS (Method A): m/z: 297 (M+1); Rt = 2.02 min.

Example 43 (General procedure (A))


3-{4-[(1H-Benzotriazole-5-carbonyl)amino]phenyl}acrylic acid



10 HPLC-MS (Method A): m/z: 309 (M+1); Rt = 3.19 min.

Example 44 (General procedure (A))


{3-[(1H-Benzotriazole-5-carbonyl)amino]phenyl}acetic acid



15 HPLC-MS (Method A): m/z: 297 (M+1); Rt = 2.10 min.

Example 45 (General procedure (A))


2-{4-[(1H-Benzotriazole-5-carbonyl)amino]phenoxy}-2-methylpropionic acid



20 HPLC-MS (Method A): m/z: 341 (M+1); Rt = 2.42 min.

Example 46 (General procedure (A))


3-{4-[(1H-Benzotriazole-5-carbonyl)amino]benzoylamino}propionic acid



HPLC-MS (Method A): m/z : 354 ($M+1$); R_t = 1.78 min.

Example 47 (General procedure (A))


- 5 3-{4-[(1H-Benzotriazole-5-carbonyl)amino]phenyl}propionic acid



HPLC-MS (Method A): m/z : 311 ($M+1$); R_t = 2.20 min.

Example 48 (General procedure (A))


- 10 1H-Benzotriazole-5-carboxylic acid (4-benzyloxyphenyl)amide



HPLC-MS (Method A): m/z : 345 ($M+1$); R_t = 3.60 min.

Example 49 (General procedure (A))


- 15 1H-Benzotriazole-5-carboxylic acid (3-chloro-4-methoxyphenyl)amide



HPLC-MS (Method A): m/z : 303 ($M+1$); R_t = 2.88 min.

Example 50 (General procedure (A))


- 20 1H-Benzotriazole-5-carboxylic acid (4-phenoxyphenyl)amide



HPLC-MS (Method A): m/z: 331 (M+1); Rt = 3.62 min.

Example 51 (General procedure (A))


- 5 1H-Benzotriazole-5-carboxylic acid (4-butoxyphenyl)amide



HPLC-MS (Method A): m/z: 311 (M+1); Rt = 3.59 min.

Example 52 (General procedure (A))


- 10 1H-Benzotriazole-5-carboxylic acid (3-bromo-4-trifluoromethoxyphenyl)amide



HPLC-MS (Method A): m/z: 402 (M+1); Rt = 3.93 min.

Example 53 (General procedure (A))


- 15 1H-Benzotriazole-5-carboxylic acid (3,5-dichloro-4-hydroxyphenyl)amide



HPLC-MS (Method A): m/z: 323 (M+1); Rt = 2.57 min.

Example 54 (General procedure (A))


- 20 4-(((1H-Benzotriazole-5-carbonyl)amino)methyl)benzoic acid



HPLC-MS (Method A): m/z: 297 (M+1); Rt = 1.86 min.

Example 55 (General procedure (A))


5 {4-[(1H-Benzotriazole-5-carbonyl)amino]phenylsulfanyl}acetic acid



HPLC-MS (Method A): m/z: 329 (M+1); Rt = 2.34 min.

Example 56


10 N-(1H-Benzotriazol-5-yl)acetamide



HPLC-MS (Method A): m/z: 177 (M+1); Rt = 0.84 min.

Example 57 (General Procedure (A))


15 1H-Benzotriazole-5-carboxylic acid 4-nitrobenzylamide



The following compound is prepared according to general procedure (N) as described below:


Example 58 (General procedure (N))

20 1H-Benzotriazole-5-carboxylic acid 4-chlorobenzylamide




HPLC-MS (Method B): m/z: 287 (M+1); Rt = 4.40 min.


Example 59 2-[(1H-Benzotriazol-5-ylimino)methyl]-4,6-dichlorophenol



5 Example 60 Diethyl 2-[(1H-benzotriazol-6-ylamino)methylidene]malonate




Example 61 N1-(1H-Benzotriazol-5-yl)-3-chlorobenzamide




10


Example 62 N1-(1H-Benzotriazol-5-yl)-3,4,5-trimethoxybenzamide



Example 63 N2-(1H-Benzotriazol-5-yl)-3-chlorobenzo[b]thiophene-2-carboxamide




Example 64 6-Bromo-1H-benzotriazole




5

Example 65 2-[(1H-Benzotriazol-5-ylimino)methyl]-4-bromophenol



10 General procedure (B) for preparation of compounds of general formula I₂:



wherein X, Y, A and R³ are as defined above and A is optionally substituted with up to four substituents R⁷, R⁸, R⁹, and R¹⁰ as defined above.

15


The chemistry is well known (eg Lohray et al., *J. Med. Chem.*, **1999**, 42, 2569-81) and is generally performed by reacting a carbonyl compound (aldehyde or ketone) with the heterocyclic ring (eg thiazolidine-2,4-dione (X = O; Y = S), rhodanine (X = Y = S) and

hydantoin (X = O; Y = NH) in the presence of a base, such as sodium acetate, potassium acetate, ammonium acetate, piperidinium benzoate or an amine (eg piperidine, triethylamine and the like) in a solvent (eg acetic acid, ethanol, methanol, DMSO, DMF, NMP, toluene, benzene) or in a mixture of two or more of these solvents. The reaction is performed at room temperature or at elevated temperature, most often at or near the boiling point of the mixture. Optionally, azeotropic removal of the formed water can be done.

This general procedure (B) is further illustrated in the following example:

Example 66 (General procedure (B))

10 5-(3-Phenoxybenzylidene)thiazolidine-2,4-dione




A solution of thiazolidine-2,4-dione (90%, 78 mg, 0.6 mmol) and ammonium acetate (92 mg, 1.2 mmol) in acetic acid (1 mL) was added to 3-phenoxybenzaldehyde (52 μ L, 0.6 mmol) and the resulting mixture was shaken at 115 °C for 16 hours. After cooling, the mixture was concentrated *in vacuo* to afford the title compound.

HPLC-MS (Method A): m/z: 298 (M+1); Rt = 4.54 min.

The compounds in the following examples were similarly prepared. Optionally, the compounds can be further purified by filtration and washing with water, ethanol and / or heptane instead of concentration *in vacuo*. Also optionally the compounds can be purified by washing with ethanol, water and/or heptane, or by chromatography, such as preparative HPLC.

Example 67 (General procedure (B))


25 5-(4-Dimethylaminobenzylidene)thiazolidine-2,4-dione



HPLC-MS (Method C): m/z: 249 (M+1); Rt = 4.90 min

Example 68 (General procedure (B))


30 5-Naphthalen-1-ylmethylenethiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 256 (M+1); Rt = 4,16 min.

Example 69 (General procedure (B))


5 5-Benzylidene-thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 206 (M+1); Rt = 4,87 min.

Example 70 (General procedure (B))


10 5-(4-Diethylaminobenzylidene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 277 (M+1); Rt = 4.73 min.

Example 71 (General procedure (B))


15 5-(4-Methoxy-benzylidene)-thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 263 (M+1); Rt = 4,90 min.

Example 72 (General procedure (B))


20 5-(4-Chloro-benzylidene)-thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 240 (M+1); Rt = 5,53 min.

Example 73 (General procedure (B))


5 5-(4-Nitro-benzylidene)-thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 251 (M+1); Rt = 4,87 min.

Example 74 (General procedure (B))


10 5-(4-Hydroxy-3-methoxy-benzylidene)-thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 252 (M+1); Rt = 4,07 min.

Example 75 (General procedure (B))


15 5-(4-Methylsulfanylbenzylidene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 252 (M+1); Rt = 5,43 min.

Example 76 (General procedure (B))

20 5-(2-Pentyloxybenzylidene)thiazolidine-2,4-dione




HPLC-MS (Method C): m/z : 292 ($M+1$); R_t = 4.75 min.

^1H NMR ($\text{DMSO}-d_6$): δ = 0.90 (3H, t), 1.39 (4H, m), 1.77 (2H, p), 4.08 (2H, t), 7.08 (1H, t), 7.14 (1H, d), 7.43 (2H, m), 8.03 (1H, s), 12.6 (1H, bs).

5

Example 77 (General procedure (B))

5-(3-Fluoro-4-methoxybenzylidene)thiazolidine-2,4-dione




HPLC-MS (Method A): m/z : 354 ($M+1$); R_t = 4.97 min.

10

Example 78 (General procedure (B))

5-(4-tert-Butylbenzylidene)thiazolidine-2,4-dione




HPLC-MS (Method A): m/z : 262 ($M+1$); R_t = 6.70 min.

15

Example 79 (General procedure (B))

N-[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)phenyl]acetamide




HPLC-MS (Method A): m/z : 263 ($M+1$); R_t = 3.90 min.

20

Example 80 (General procedure (B))

5-Biphenyl-4-ylmethylene-thiazolidine-2,4-dione




HPLC-MS (Method A): m/z: 282 (M+1); Rt = 4,52 min.

5

Example 81 (General procedure (B))

5-(4-Phenoxy-benzylidene)-thiazolidine-2,4-dione




HPLC-MS (Method A): m/z: 298 (M+1); Rt = 6,50 min.

10

Example 82 (General procedure (B))

5-(3-Benzyloxybenzylidene)thiazolidine-2,4-dione




HPLC-MS (Method A): m/z: 312 (M+1); Rt = 6,37 min.

15

Example 83 (General procedure (B))

5-(3-p-Tolyloxybenzylidene)thiazolidine-2,4-dione




HPLC-MS (Method A): m/z: 312 (M+1); Rt = 6,87 min.

20

Example 84 (General procedure (B))


5-Naphthalen-2-ylmethylene-thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 256 (M+1); Rt = 4.15 min.

Example 85 (General procedure (B))

5-Benzo[1,3]dioxol-5-ylmethylenethiazolidine-2,4-dione




5

HPLC-MS (Method A): m/z: 250 (M+1), Rt = 3.18 min.

Example 86 (General procedure (B))

5-(4-Chlorobenzylidene)-2-thioxothiazolidin-4-one




10

HPLC-MS (Method A): m/z: 256 (M+1); Rt = 4,51 min.

Example 87 (General procedure (B))

5-(4-Dimethylaminobenzylidene)-2-thioxothiazolidin-4-one




15

HPLC-MS (Method A): m/z: 265 (M+1); Rt = 5,66 min.

Example 88 (General procedure (B))

5-(4-Nitrobenzylidene)-2-thioxothiazolidin-4-one




20

HPLC-MS (Method A): m/z: 267 (M+1); Rt = 3,94 min.

Example 89 (General procedure (B))

5-(4-Methylsulfanylbenzylidene)-2-thioxothiazolidin-4-one




HPLC-MS (Method A): m/z: 268 (M+1); Rt = 6,39 min.

5

Example 90 (General procedure (B))

5-(3-Fluoro-4-methoxybenzylidene)-2-thioxothiazolidin-4-one




HPLC-MS (Method A): m/z: 270 (M+1); Rt = 5,52 min.

10

Example 91 (General procedure (B))

5-Naphthalen-2-ylmethylene-2-thioxothiazolidin-4-one




HPLC-MS (Method A): m/z: 272 (M+1); Rt = 6,75 min.

15

Example 92 (General procedure (B))

5-(4-Diethylaminobenzylidene)-2-thioxothiazolidin-4-one




HPLC-MS (Method A): m/z: 293 (M+1); Rt = 5,99 min.

20

Example 93 (General procedure (B))


5-Biphenyl-4-ylmethylene-2-thioxothiazolidin-4-one



HPLC-MS (Method A): m/z: 298 (M+1); Rt = 7,03 min.

Example 94 (General procedure (B))


- 5 5-(3-Phenoxybenzylidene)-2-thioxothiazolidin-4-one



HPLC-MS (Method A): m/z: 314 (M+1); Rt = 6,89 min.

Example 95 (General procedure (B))


- 10 5-(3-Benzoyloxybenzylidene)-2-thioxothiazolidin-4-one



HPLC-MS (Method A): m/z: 328 (M+1); Rt = 6,95 min.

Example 96 (General procedure (B))


- 15 5-(4-Benzoyloxybenzylidene)-2-thioxothiazolidin-4-one



HPLC-MS (Method A): m/z: 328 (M+1); RT = 6,89 min.

Example 97 (General procedure (B))


- 20 5-Naphthalen-1-ylmethylene-2-thioxothiazolidin-4-one



HPLC-MS (Method A): m/z: 272 (M+1); Rt = 6,43 min.

Example 98 (General procedure (B))


5-(3-Methoxybenzyl)thiazolidine-2,4-dione



5 HPLC-MS (Method A): m/z: 236 (M+1); Rt = 3,05 min.

Example 99 (General procedure (D))


4-[2-Chloro-4-(2,4-dioxothiazolidin-5-ylidenemethyl)phenoxy]butyric acid ethyl ester



10 HPLC-MS (Method A): m/z: 392 (M+23), Rt = 4.32 min.

Example 100 (General procedure (D))


4-[2-Bromo-4-(2,4-dioxothiazolidin-5-ylidenemethyl)-phenoxy]-butyric acid



15 HPLC-MS (Method A): m/z: 410 (M+23); Rt = 3,35 min.

Example 101 (General procedure (B))


5-(3-Bromobenzylidene)thiazolidine-2,4-dione



20 HPLC-MS (Method A): m/z: 285 (M+1); Rt = 4.01 min.

Example 102 (General procedure (B))


5-(4-Bromobenzylidene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 285 (M+1); Rt = 4.05 min.

Example 103 (General procedure (B))


5 5-(3-Chlorobenzylidene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 240 (M+1); Rt = 3.91 min.

Example 104 (General procedure (B))


10 5-Thiophen-2-ylmethylenethiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 212 (M+1); Rt = 3.09 min.

Example 105 (General procedure (B))


15 5-(4-Bromothiophen-2-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 291 (M+1); Rt = 3.85 min.

Example 106 (General procedure (B))


20 5-(3,5-Dichlorobenzylidene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 274 (M+1); Rt = 4.52 min.

Example 107 (General procedure (B))


5 5-(1-Methyl-1H-indol-3-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 259 (M+1); Rt = 3.55 min.

Example 108 (General procedure (B))


10 5-(1H-Indol-3-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 245 (M+1); Rt = 2.73 min.

Example 109 (General procedure (B))


15 5-Fluoren-9-ylidenethiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 280 (M+1); Rt = 4.34 min.

Example 110 (General procedure (B))

5-(1-Phenylethylidene)thiazolidine-2,4-dione




HPLC-MS (Method A): m/z: 220 (M+1); Rt = 3,38 min.

5

Example 111 (General procedure (B))

5-[1-(4-Methoxyphenyl)-ethylidene]-thiazolidine-2,4-dione




HPLC-MS (Method A): m/z: 250 (M+1); Rt = 3.55 min.

10

Example 112 (General procedure (B))

5-(1-Naphthalen-2-yl-ethylidene)-thiazolidine-2,4-dione




HPLC-MS (Method A): m/z: 270 (M+1); Rt = 4,30 min.

15

Example 113 (General procedure (B))

5-[1-(4-Bromophenyl)-ethylidene]-thiazolidine-2,4-dione




HPLC-MS (Method A): m/z: 300 (M+1); Rt = 4,18 min.

20

Example 114 (General procedure (B))

5-(2,2-Diphenylethylidene)-thiazolidine-2,4-dione




HPLC-MS (Method A): m/z: 296 (M+1); Rt = 4,49 min.

5

Example 115 (General procedure (B))

5-[1-(3-Methoxyphenyl)-ethylidene]-thiazolidine-2,4-dione




HPLC-MS (Method A): m/z: 250 (M+1); Rt = 3,60 min.

10

Example 116 (General procedure (B))

5-[1-(6-Methoxynaphthalen-2-yl)-ethylidene]-thiazolidine-2,4-dione




HPLC-MS (Method A): m/z: 300 (M+1); Rt = 4,26 min.

15

Example 117 (General procedure (B))

5-[1-(4-Phenoxyphenyl)-ethylidene]-thiazolidine-2,4-dione




HPLC-MS (Method A): m/z: 312 (M+1); Rt = 4,68 min.

20

Example 118 (General procedure (B))


5-[1-(3-Fluoro-4-methoxyphenyl)ethylidene]thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 268 (M+1); Rt = 3,58 min.

Example 119 (General procedure (B))


- 5 5-[1-(3-Bromophenyl)-ethylidene]-thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 300 (M+1); Rt = 4,13 min.

Example 120 (General procedure (B))


- 10 5-Anthracen-9-ylmethylenethiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 306 (M+1); Rt = 4,64 min.

Example 121 (General procedure (B))


- 15 5-(2-Methoxynaphthalen-1-ylmethylene)-thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 286 (M+1); Rt = 4,02 min.

Example 122 (General procedure (B))


- 20 5-(4-Methoxynaphthalen-1-ylmethylene)-thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 286 (M+1); Rt = 4,31 min.

Example 123 (General procedure (B))


- 5 5-(4-Dimethylaminonaphthalen-1-ylmethylene)-thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 299 (M+1); Rt = 4,22 min.

Example 124 (General procedure (B))


- 10 5-(4-Methylnaphthalen-1-ylmethylene)-thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 270 (M+1); Rt = 4,47 min.


Example 125 (General procedure (B))

- 15 5-Pyridin-2-ylmethylene-thiazolidine-2,4-dione



Example 126

- 5-Pyridin-2-ylmethyl-thiazolidine-2,4-dione




- 5-Pyridin-2-ylmethylene-thiazolidine-2,4-dione (5 g) in tetrahydrofuran (300 ml) was added 10% Pd/C (1 g) and the mixture was hydrogenated at ambient pressure for 16 hours. More 10% Pd/C (5 g) was added and the mixture was hydrogenated at 50 psi for 16 hours. After filtration and evaporation *in vacuo*, the residue was purified by column chromatography eluting with a mixture of ethyl acetate and heptane (1:1). This afforded the title compound (0.8 g, 16%) as a solid.

TLC: R_f = 0.30 (SiO₂; EtOAc: heptane 1:1)

10


Example 127 (General procedure (B))

5-(1H-Imidazol-4-ylmethylene)-thiazolidine-2,4-dione



15 Example 128 (General procedure (B))


5-(4-Benzyloxy-benzylidene)-thiazolidine-2,4-dione



HPLC-MS (Method A): m/z : 6,43 min ; 99 % (2A)


20 Example 129 (General procedure (B))

5-[4-(4-Fluorobenzyloxy)benzylidene]-2-thioxothiazolidin-4-one



Example 130 (General procedure (B))


5-(4-Butoxybenzylidene)-2-thioxothiazolidin-4-one



5

Example 131 (General procedure (B))


5-(3-Methoxybenzylidene)thiazolidine-2,4-dione



10 HPLC-MS (Method A): m/z: 236 (M+1); Rt = 4,97 min

Example 132 (General procedure (B))


15 5-(3-Methoxybenzylidene)imidazolidine-2,4-dione



HPLC-MS (Method A): m/z: 219 (M+1); Rt = 2.43 min.

Example 133 (General procedure (B))


20 5-(4-Methoxybenzylidene)imidazolidine-2,4-dione



HPLC-MS (Method A): m/z: 219 (M+1); Rt = 2.38 min.


Example 134 (General procedure (B))

5-(2,3-Dichlorobenzylidene)thiazolidine-2,4-dione



5 Example 135 (General procedure (B))


5-Benzofuran-7-ylmethylenethiazolidine-2,4-dione



HPLC-MS (Method C): m/z: 247 (M+1); Rt = 4,57 min.

10 Example 136 (General procedure (B))


5-Benzo[1,3]dioxol-4-ylmethylenethiazolidine-2,4-dione



HPLC-MS (Method C): m/z: 250 (M+1); Rt = 4,00 min.

15 Example 137 (General procedure (B))


5-(4-Methoxy-2,3-dimethylbenzylidene)thiazolidine-2,4-dione



HPLC-MS (Method C): m/z: 264 (M+1); Rt = 5,05 min.

20 Example 138 (General procedure (B))


5-(2-Benzyloxy-3-methoxybenzylidene)thiazolidine-2,4-dione



HPLC-MS (Method C): m/z: 342 (M+1); Rt = 5,14 min.

Example 139 (General procedure (B))


5 5-(2-Hydroxybenzylidene)thiazolidine-2,4-dione



HPLC-MS (Method C): m/z: 222 (M+1); Rt = 3,67 min.

Example 140 (General procedure (B))


10 5-(2,4-Dichlorobenzylidene)thiazolidine-2,4-dione



¹H-NMR (DMSO-*d*₆): 7.60 (2H, "s"), 7.78 (1H, s), 7.82 (1H, s).

Example 141 (General procedure (B))


15 5-(2-Chlorobenzylidene)thiazolidine-2,4-dione



¹H-NMR (DMSO-*d*₆): 7.40 (1H, t), 7.46 (1H, t), 7.57 (1H, d), 7.62 (1H, d), 7.74 (1H, s).

Example 142 (General procedure (B))


20 5-(2-Bromobenzylidene)thiazolidine-2,4-dione



¹H-NMR (DMSO-*d*₆): 7.33 (1H, t), 7.52 (1H, t), 7.60 (1H, d), 7.71 (1H, s), 7.77 (1H, d).

Example 143 (General procedure (B))


5-(2,4-Dimethoxybenzylidene)thiazolidine-2,4-dione



5 HPLC-MS (Method C): m/z: 266 (M+1) Rt = 4,40 min.

Example 144 (General procedure (B))


5-(2-Methoxybenzylidene)thiazolidine-2,4-dione



10 HPLC-MS (Method C): m/z: 236 (M+1); Rt = 4,17 min.

Example 145 (General procedure (B))


5-(2,6-Difluorobenzylidene)thiazolidine-2,4-dione



15 HPLC-MS (Method C): m/z: 242 (M+1); Rt = 4,30 min.

Example 146 (General procedure (B))


5-(2,4-Dimethylbenzylidene)thiazolidine-2,4-dione



20 HPLC-MS (Method C): m/z: 234 (M+1); Rt = 5,00 min.

Example 147 (General procedure (B))


5-(2,4,6-Trimethoxybenzylidene)thiazolidine-2,4-dione



HPLC-MS (Method C): m/z: 296 (M+1); Rt = 4,27 min.

Example 148 (General procedure (B))


5 5-(4-Hydroxy-2-methoxybenzylidene)thiazolidine-2,4-dione



HPLC-MS (Method C): m/z: 252 (M+1); Rt = 3,64 min.

Example 149 (General procedure (B))

10 5-(4-Hydroxynaphthalen-1-ylmethylene)thiazolidine-2,4-dione




¹H-NMR (DMSO-*d*₆): δ = 7.04 (1H, d), 7.57 (2H, m), 7.67 (1H, t), 8.11 (1H, d), 8.25 (1H, d), 8.39 (1H, s) 11.1 (1H, s), 12.5 (1H, bs). HPLC-MS (Method C): m/z: 272 (M+1); Rt = 3.44 min.

15

Example 150 (General procedure (B))

5-(2-Trifluoromethoxybenzylidene)thiazolidine-2,4-dione




HPLC-MS (Method C): m/z: 290 (M+1); Rt = 4,94 min.

20

Example 151 (General procedure (B))


5-Biphenyl-2-ylmethylenethiazolidine-2,4-dione



HPLC-MS (Method C): m/z: 282 (M+1); Rt = 5,17 min.

Example 152 (General procedure (B))


5 5-(2-Benzyloxybenzylidene)thiazolidine-2,4-dione



HPLC-MS (Method C): m/z: 312 (M+1); Rt = 5,40 min.

Example 153 (General procedure (B))


10 5-Adamantan-2-ylidenethiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 250 (M+1); Rt = 4,30 min.

Example 154 (General Procedure (B))


15 5-[3-(4-Nitrophenyl)allylidene]thiazolidine-2,4-dione



HPLC-MS (Method C): m/z: 277 (M+1); Rt = 3.63 min.

Example 155 (General Procedure (B))


20 5-[3-(2-Methoxyphenyl)allylidene]thiazolidine-2,4-dione



HPLC-MS (Method C): m/z : 262 ($M+1$); R_t = 3.81 min.

Example 156 (General Procedure (B))


5 5-[3-(4-Methoxyphenyl)allylidene]thiazolidine-2,4-dione



HPLC-MS (Method C): m/z : 262 ($M+1$); R_t = 3.67 min.


Example 157 (General procedure (B))

10 5-(4-Hydroxybenzylidene)thiazolidine-2,4-dione



Example 158 (General procedure (B))

5-(4-Dimethylaminobenzylidene)pyrimidine-2,4,6-trione




15

HPLC-MS (Method C): m/z = 260 ($M+1$) R_t = 2,16 min.

Example 159 (General procedure (B))


5-(9-Ethyl-9H-carbazol-2-ylmethylene)-pyrimidine-2,4,6-trione



HPLC-MS (Method C): $m/z = 334$ ($M+1$); $R_t = 3,55$ min.

Example 160 (General procedure (B))


5 5-(4-Hexyloxynaphthalen-1-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method C): $m/z = 356$ ($M+1$); $R_t = 5.75$ min.

Example 161 (General procedure (B))


10 5-(4-Decyloxynaphthalen-1-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method C): $m/z = 412$ ($M+1$); $R_t = 6.44$ min.

Example 162 (General procedure (B))


15 5-[4-(2-Aminoethoxy)-naphthalen-1-ylmethylene]-thiazolidine-2,4-dione



HPLC-MS (Method C): $m/z = 315$ ($M+1$); $R_t = 3,24$ min.

Example 163 (General procedure (B))


5 5-(2,4-Dimethyl-9H-carbazol-3-ylmethylene)-pyrimidine-2,4,6-trione



HPLC-MS (Method C): $m/z = 334$ ($M+1$); $R_t = 3,14$ min.


Example 164 (General procedure (B))

10 4-(4-Hydroxy-3-methoxybenzylidene)hydantoin




Example 165 (General procedure (B))

5-Benzylidenehydantoin



15

General procedure (C) for preparation of compounds of general formula I₂:




wherein X, Y, A, and R³ are as defined above and A is optionally substituted with up to four substituents R⁷, R⁸, R⁹, and R¹⁰ as defined above.

- 5 This general procedure (C) is quite similar to general procedure (B) and is further illustrated in the following example:

Example 166 (General procedure (C))

5-(3,4-Dibromobenzylidene)thiazolidine-2,4-dione



10

A mixture of thiazolidine-2,4-dione (90%, 65 mg, 0.5 mmol), 3,4-dibromobenzaldehyde (132 mg, 0.5 mmol), and piperidine (247 μ L, 2.5 mmol) was shaken in acetic acid (2 mL) at 110 °C for 16 hours. After cooling, the mixture was concentrated to dryness *in vacuo*.


- 15 The resulting crude product was shaken with water, centrifuged, and the supernatant was discarded. Subsequently the residue was shaken with ethanol, centrifuged, the supernatant was discarded and the residue was further evaporated to dryness to afford the title compound.

- 20 ¹H NMR (Acetone-*d*₆): δ _H 7.99 (d, 1H), 7.90 (d, 1H), 7.70 (s, 1H), 7.54 (d, 1H); HPLC-MS (Method A): *m/z*: 364 (M+1); *R*_t = 4.31 min.

- 25 The compounds in the following examples were similarly prepared. Optionally, the compounds can be further purified by filtration and washing with water instead of concentration *in vacuo*. Also optionally the compounds can be purified by washing with ethanol, water and/or heptane, or by preparative HPLC.

Example 167 (General procedure (C))

5-(4-Hydroxy-3-iodo-5-methoxybenzylidene)thiazolidine-2,4-dione




Mp = 256 °C; ^1H NMR (DMSO- d_6) δ = 12.5 (s,broad,1H), 10.5 (s,broad,1H), 7.69 (s,1H), 7.51 (d,1H), 7.19 (d,1H), 3.88 (s,3H), ^{13}C NMR (DMSO- d_6) δ_c = 168.0, 167.7, 149.0, 147.4, 133.0, 131.2, 126.7, 121.2, 113.5, 85.5, 56.5; HPLC-MS (Method A): m/z: 378 (M+1); Rt = 3.21 min.

5

Example 168 (General procedure (C))


5-(4-Hydroxy-2,6-dimethylbenzylidene)thiazolidine-2,4-dione



10 HPLC-MS (Method C): m/z: 250 (M+1); Rt.= 2.45 min.

Example 169 (General procedure (C))

4-[5-Bromo-6-(2,4-dioxothiazolidin-5-ylidenemethyl)-naphthalen-2-yloxy-methyl]-benzoic acid




15

HPLC-MS (Method C): m/z: 506 (M+23); Rt.= 4.27 min.

Example 170 (General procedure (C))

5-(4-Bromo-2,6-dichlorobenzylidene)thiazolidine-2,4-dione




20

HPLC-MS (Method C): m/z: 354 (M+1); Rt.= 4.36 min.

Example 171 (General procedure (C))

5-(6-Hydroxy-2-naphthylmethylene) thiazolidine-2,4-dione



Mp 310-314 °C, ¹H NMR (DMSO-*d*₆): δ_H = 12.5 (s,broad,1H), 8.06(d,1H), 7.90-7.78(m,2H), 7.86 (s,1H), 7.58 (dd,1H), 7.20 7.12 (m,2H). ¹³C NMR (DMSO-*d*₆): δ_C = 166.2, 165.8, 155.4, 133.3, 130.1, 129.1, 128.6, 125.4, 125.3, 125.1, 124.3, 120.0, 117.8, 106.8; HPLC-MS (Method A): m/z: 272 (M+1); Rt = 3.12 min.

Preparation of the starting material, 6-hydroxy-2-naphtalenecarbaldehyde:

6-Cyano-2-naphthalenecarbaldehyde (1.0 g, 5.9 mmol) was dissolved in dry hexane (15 mL) under nitrogen. The solution was cooled to -60 °C and a solution of diisobutyl aluminium hydride (DIBAH) (15 mL, 1M in hexane) was added dropwise. After the addition, the solution was left at room temperature overnight. Saturated ammonium chloride solution (20 mL) was added and the mixture was stirred at room temperature for 20 min, subsequently aqueous H₂SO₄ (10% solution, 15 mL) was added followed by water until all salt was dissolved. The resulting solution was extracted with ethyl acetate (3x), the combined organic phases were dried with MgSO₄, evaporated to dryness to afford 0.89 g of 6-hydroxy-2-naphtalenecarbaldehyde.

Mp.: 153.5-156.5 °C; HPLC-MS (Method A): m/z: 173 (M+1); Rt = 2.67 min; ¹H NMR (DMSO-*d*₆): δ_H = 10.32(s,1H), 8.95 (d,1H), 10.02 (s,1H), 8.42 (s,broad,1H), 8.01 (d,1H), 7.82-7.78 (m,2H), 7.23-7.18 (m,2H).

Alternative preparation of 6-hydroxy-2-naphtalenecarbaldehyde:


To a stirred cooled mixture of 6-bromo-2-hydroxynaphthalene (25.3 g, 0.113 mol) in THF (600 mL) at -78 °C was added n-BuLi (2.5 M, 100 mL, 0.250 mol) dropwise. The mixture turned yellow and the temperature rose to -64 °C. After ca 5 min a suspension appeared. After addition, the mixture was maintained at -78 °C. After 20 minutes, a solution of DMF (28.9 mL, 0.373 mol) in THF (100 mL) was added over 20 minutes. After addition, the mixture was allowed to warm slowly to room temperature. After 1 hour, the mixture was poured in ice/water (200 mL). To the mixture citric acid was added to a pH of 5. The mixture was stirred for 0.5 hour. Ethyl acetate (200 mL) was added and the organic layer was

separated and washed with brine (100 mL), dried over Na_2SO_4 and concentrated. To the residue was added heptane with 20% ethyl acetate (ca 50 mL) and the mixture was stirred for 1 hour. The mixture was filtered and the solid was washed with ethyl acetate and dried *in vacuo* to afford 16 g of the title compound.

5

Example 172 (General procedure (C))

5-(3-Iodo-4-methoxybenzylidene)thiazolidine-2,4-dione



^1H NMR ($\text{DMSO}-d_6$): δ_{H} 12.55 (s, broad, 1H), 8.02 (d, 1H), 7.72 (s, 1H), 7.61 (d, 1H), 7.18 (d, 1H), 3.88 (s, 3H); ^{13}C NMR ($\text{DMSO}-d_6$): δ_{C} 168.1, 167.7, 159.8, 141.5, 132.0, 130.8, 128.0, 122.1, 112.5, 87.5, 57.3. HPLC-MS (Method A): m/z : 362 ($\text{M}+1$); Rt = 4.08 min.

10

Preparation of the starting material, 3-iodo-4-methoxybenzaldehyde:

15 4-Methoxybenzaldehyde (0.5 g, 3.67 mmol) and silver trifluoroacetate (0.92 g, 4.19 mmol) were mixed in dichloromethane (25 mL). Iodine (1.19 g, 4.7 mmol) was added in small portions and the mixture was stirred overnight at room temperature under nitrogen. The mixture was subsequently filtered and the residue washed with DCM. The combined filtrates were treated with an aqueous sodium thiosulfate solution (1 M) until the colour disappeared.


20 Subsequent extraction with dichloromethane (3 x 20 mL) followed by drying with MgSO_4 and evaporation *in vacuo* afforded 0.94 g of 3-iodo-4-methoxybenzaldehyde.

Mp 104-107 °C; HPLC-MS (Method A): m/z : 263 ($\text{M}+1$); Rt = 3.56 min.; ^1H NMR (CDCl_3): δ_{H} = 8.80 (s, 1H), 8.31 (d, 1H), 7.85 (dd, 1H), 6.92 (d, 1H), 3.99 (s, 3H).

25

Example 173 (General procedure (C))

5-(1-Bromonaphthalen-2-ylmethylene)thiazolidine-2,4-dione




HPLC-MS (Method A): m/z : =336 ($\text{M}+1$); Rt = 4.46 min.

30

Example 174 (General procedure (C))


1-[5-(2,4-Dioxothiazolidin-5-ylidenemethyl)thiazol-2-yl]piperidine-4-carboxylic acid ethyl ester



^1H NMR ($\text{DMSO}-d_6$): δ_{H} = 7.88 (s,1H), 7.78 (s,1H), 4.10 (q,2H), 4.0-3.8 (m,2H), 3.40-3.18 (m,2H), 2.75-2.60 (m,1H), 2.04-1.88 (m,2H), 1.73-1.49 (m,2H), 1.08 (t,3H); HPLC-MS (Method A): m/z: 368 ($\text{M}+1$); Rt = 3.41 min.

Example 175 (General procedure (C))


5-(2-Phenyl-[1,2,3]triazol-4-ylmethylene) thiazolidine-2,4-dione



^1H NMR ($\text{DMSO}-d_6$): δ_{H} = 12.6 (s,broad,1H), 8.46 (s,1H), 8.08 (dd,2H), 7.82 (s,1H), 7.70-7.45 (m, 3H). HPLC-MS (Method A): m/z: 273 ($\text{M}+1$); Rt = 3.76 min.

Example 176 (General procedure (C))


5-(Quinolin-4-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 257 ($\text{M}+1$); Rt = 2.40 min.

Example 177 (General procedure (C))


5-(6-Methylpyridin-2-ylmethylene)thiazolidine-2,4-dione



^1H NMR ($\text{DMSO}-d_6$): δ_{H} = 12.35 (s, broad, 1H), 7.82 (t, 1H), 7.78 (s, 1H), 7.65 (d, 1H), 7.18 (d, 1H), 2.52 (s, 3H); HPLC-MS (Method A): m/z : 221 ($\text{M}+1$); Rt = 3.03 min.

5 Example 178 (General procedure (C))

5-(2,4-dioxothiazolidin-5-ylidenemethyl)-furan-2-ylmethylacetate




^1H NMR ($\text{DMSO}-d_6$): δ_{H} = 12.46 (s, broad, 1H), 7.58 (s, 1H), 7.05 (d, 1H), 6.74 (s, 1H), 5.13 (s, 2H), 2.10 (s, 3H). HPLC-MS (Method A): m/z : 208 ($\text{M}-\text{CH}_3\text{COO}$); Rt = 2.67 min.

10

Example 179 (General procedure (C))

5-(2,4-Dioxothiazolidin-5-ylidenemethyl)furan-2-sulfonic acid




HPLC-MS (Method A): m/z : 276 ($\text{M}+1$); Rt = 0.98 min.

15

Example 180 (General procedure (C))

5-(5-Benzyloxy-1H-pyrrolo[2,3-c]pyridin-3-ylmethylene)-thiazolidine-2,4-dione




HPLC-MS (Method A): m/z : 352 ($\text{M}+1$); Rt = 3.01 min.

20

Example 181 (General procedure (C))


5-(Quinolin-2-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 257 (M+1); Rt = 3.40 min.

Example 182 (General procedure (C))


- 5 5-(2,4-Dioxothiazolidin-5-ylidenemethyl)thiophene-2-carboxylic acid



HPLC-MS (Method A): m/z: 256 (M+1); Rt = 1.96 min.

Example 183 (General procedure (C))


- 10 5-(2-Phenyl-1H-imidazol-4-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 272 (M+1); Rt = 2.89 min.

Example 184 (General procedure (C))


- 15 5-(4-Imidazol-1-yl-benzylidene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 272 (M+1); Rt = 1.38 min.

Example 185 (General procedure (C))


- 20 5-(9-Ethyl-9H-carbazol-3-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 323 (M+1); Rt = 4.52 min.

Example 186 (General procedure (C))


5-(1,4-Dimethyl-9H-carbazol-3-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 323 (M+1); Rt = 4.35 min.

Example 187 (General procedure (C))


5-(2-Methyl-1H-indol-3-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 259 (M+1); Rt = 3.24 min.

Example 188 (General procedure (C))

5-(2-Ethylindol-3-ylmethylene)thiazolidine-2,4-dione



2-Methylindole (1.0 g, 7.6mmol) dissolved in diethyl ether (100 mL) under nitrogen was treated with n-Butyl lithium (2 M in pentane, 22.8 mmol) and potassium *tert*-butoxide (15.2 mmol) with stirring at RT for 30 min. The temperature was lowered to -70 C and methyl iodide (15.2 mmol) was added and the resulting mixture was stirred at -70 for 2 h. Then 5 drops of water was added and the mixture allowed to warm up to RT. Subsequently, the mixture was poured into water (300 mL), pH was adjusted to 6 by means of 1N hydrochloric acid and the mixture was extracted with diethyl ether. The organic phase was dried with Na₂SO₄ and evaporated to dryness. The residue was purified by column chromatography on silica gel using heptane/ether(4/1) as eluent. This afforded 720 mg (69 %) of 2-ethylindole.

¹H NMR (DMSO-*d*₆): δ = 10.85 (1H,s); 7.39 (1H,d); 7.25 (1H,d); 6.98(1H,t); 6.90(1H,t); 6.10 (1H,s); 2.71 (2H,q); 1.28 (3H,t).

2-Ethylindole (0.5 g, 3.4mmol) dissolved in DMF (2 mL) was added to a cold (0 °C) premixed (30 minutes) mixture of DMF (1.15 mL) and phosphorous oxychloride (0.64 g, 4.16 mmol). After addition of 2-ethylindole, the mixture was heated to 40 °C for 1 h, water (5 mL) was added and the pH adjusted to 5 by means of 1 N sodium hydroxide. The mixture was subsequently extracted with diethyl ether, the organic phase isolated, dried with MgSO₄ and evaporated to dryness affording 2-ethylindole-3-carbaldehyde (300 mg).


HPLC-MS (Method C): m/z:174 (M+1); Rt. =2.47 min.

2-Ethylindole-3-carbaldehyde (170 mg) was treated with thiazolidine-2,4-dione using the general procedure (C) to afford the title compound (50 mg).

HPLC-MS (Method C):m/z: 273 (M+1); Rt.= 3.26 min.

Example 189 (General procedure (C))


5-[2-(4-Bromophenylsulfanyl)-1-methyl-1H-indol-3-ylmethylene]thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 447 (M+1); Rt = 5.25 min.

Example 190 (General procedure (C))


5-[2-(2,4-Dichlorobenzyloxy)-naphthalen-1-ylmethylene]thiazolidine-2,4-dione



HPLC-MS (Method A): (anyone 1) m/z : 430 ($M+1$); R_t = 5.47 min.

Example 191 (General procedure (C))


- 5 5-{4-[3-(4-Bromophenyl)-3-oxopropenyl]-benzylidene}thiazolidine-2,4-dione



HPLC-MS (Method A): m/z : 416 ($M+1$); R_t = 5.02 min.

Example 192 (General procedure (C))


- 10 5-(4-Pyridin-2-ylbenzylidene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z : 283 ($M+1$), R_t = 2.97 min.

Example 193 (General procedure (C))


- 15 5-(3,4-Bisbenzyloxybenzylidene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z : 418 ($M+1$); R_t = 5.13 min.

Example 194 (General procedure (C))


- 20 5-[4-(4-Nitrobenzyloxy)-benzylidene]thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 357 (M+1); Rt = 4.45 min.

Example 195 (General procedure (C))


5 5-(2-Phenyl-1H-indol-3-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 321 (M+1); Rt = 3.93 min.

Example 196 (General procedure (C))


10 5-(5-Benzyloxy-1H-indol-3-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 351 (M+1); Rt = 4.18 min.

Example 197 (General procedure (C))


15 5-(4-Hydroxybenzylidene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z: 222 (M+1); Rt = 2.42 min.

Example 198 (General procedure (C))

5-(1-Methyl-1H-indol-2-ylmethylene)thiazolidine-2,4-dione




^1H NMR ($\text{DMSO}-d_6$): $\delta_{\text{H}} = 12.60$ (s, broad, 1H), 7.85 (s, 1H), 7.68 (dd, 1H), 7.55 (dd, 1H), 7.38 (dt, 1H), 7.11 (dt, 1H) 6.84 (s, 1H), 3.88 (s, 3H); HPLC-MS (Method A): m/z : 259 (M+1); $R_t = 4.00$ min.

5

Example 199 (General procedure (C))

5-(5-Nitro-1H-indol-3-ylmethylene)thiazolidine-2,4-dione




.Mp 330-333 °C, ^1H NMR ($\text{DMSO}-d_6$): $\delta_{\text{H}} = 12.62$ (s, broad, 1H), 8.95 (d, 1H), 8.20 (s, 1H), 8.12 (dd, 1H), 7.98 (s, broad, 1H), 7.68 (d, 1H); HPLC-MS (Method A): m/z : 290 (M+1); $R_t = 3.18$ min.

10

Example 200 (General procedure (C))

5-(6-Methoxynaphthalen-2-ylmethylene)thiazolidine-2,4-dione




HPLC-MS (Method A): m/z : 286 (M+1); $R_t = 4.27$ min.

15

Example 201 (General procedure (C))

5-(3-Bromo-4-methoxybenzylidene)thiazolidine-2,4-dione




HPLC-MS (Method A): m/z : 314 (M+1), $R_t = 3.96$ min.

20

Example 202 (General procedure (C))


3-((2-Cyanoethyl)-[4-(2,4-dioxothiazolidin-5-ylidenemethyl)phenyl]amino)propionitrile



HPLC-MS (Method A): m/z : 327 ($M+1$); R_t = 2.90 min.

Example 203 (General procedure (C))


- 5 3-(2,4-Dioxothiazolidin-5-ylidenemethyl)indole-6-carboxylic acid methyl ester



HPLC-MS (Method A): m/z : 303 ($M+1$); R_t = 3.22-3.90 min.

Example 204

- 10 3-(2,4-Dioxothiazolidin-5-ylidenemethyl)indole-6-carboxylic acid pentyl ester.




3-(2,4-Dioxothiazolidin-5-ylidenemethyl)indole-6-carboxylic acid methyl ester (example 203, 59 mg; 0.195mmol) was stirred in pentanol (20 mL) at 145 °C for 16 hours. The mixture was evaporated to dryness affording the title compound (69 mg).

15

HPLC-MS (Method C): m/z : 359 ($M+1$); R_t = 4.25 min.

Example 205 (General procedure (C))

- 3-(2,4-Dioxothiazolidin-5-ylidenemethyl)indole-7-carboxylic acid




20

HPLC-MS (Method A): m/z : 289 ($M+1$); R_t = 2.67 min.

Example 206 (General procedure (C))


- 5-(1-Benzylindol-3-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z : 335 ($M+1$); R_t = 4.55 min.

Example 207 (General procedure (C))


5 5-(1-Benzenesulfonylindol-3-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z : = 385 ($M+1$); R_t = 4.59 min.

Example 208 (General procedure (C))


10 5-(4-[1,2,3]Thiadiazol-4-ylbenzylidene)thiazolidine-2,4-dione



HPLC-MS (Method A): m/z : 290 ($M+1$); R_t = 3.45 min.

Example 209 (General procedure (C))


15 5-[4-(4-Nitrobenzyloxy)-benzylidene]thiazolidine-2,4-dione



HPLC-MS (Method A): m/z : 357 ($M+1$); R_t = 4.42 min.

Example 210 (General procedure (C))

3-(2,4-Dioxothiazolidin-5-ylidenemethyl)indole-1-carboxylic acid ethyl ester




HPLC-MS (Method A): m/z: 317 (M+1); Rt = 4.35 min.

5

Example 211 (General procedure (C))

5-[2-(4-Pentylbenzoyl)-benzofuran-5-ylmethylene]thiazolidine-2,4-dione




HPLC-MS (Method A): m/z: 420 (M+1); Rt = 5.92 min.

10

Example 212 (General procedure (C))

5-[1-(2-Fluorobenzyl)-4-nitroindol-3-ylmethylene]thiazolidine-2,4-dione




HPLC-MS (Method A): (Anyone 1) m/z: 398 (M+1); Rt = 4.42 min.

15

Example 213 (General procedure (C))

5-(4-Benzyloxyindol-3-ylmethylene)thiazolidine-2,4-dione




HPLC-MS (Method A): m/z: 351 (M+1); Rt = 3.95 min.

20

Example 214 (General procedure (C))

5-(4-Isobutylbenzylidene)-thiazolidine-2,4-dione




HPLC-MS (Method A): m/z: 262 (M+1); Rt = 4.97 min.

5

Example 215 (General procedure (C))

Trifluoromethanesulfonic acid 4-(2,4-dioxothiazolidin-5-ylidenemethyl)naphthalen-1-yl ester



HPLC-MS (Method A): m/z: 404 (M+1); Rt = 4.96 min.

10

Preparation of starting material:


4-Hydroxy-1-naphthaldehyde (10 g, 58 mmol) was dissolved in pyridin (50 ml) and the mixture was cooled to 0-5 °C. With stirring, trifluoromethanesulfonic acid anhydride (11.7 ml, 70 mmol) was added drop-wise. After addition was complete, the mixture was allowed to warm up to room temperature, and diethyl ether (200 ml) was added. The mixture was washed with water (2 x 250 ml), hydrochloric acid (3N, 200 ml), and saturated aqueous sodium chloride (100 ml). After drying (MgSO₄), filtration and concentration in vacuo, the residue was purified by column chromatography on silica gel eluting with a mixture of ethyl acetate and heptane (1:4). This afforded 8.35 g (47%) trifluoromethanesulfonic acid 4-formylnaphthalen-1-yl ester, mp 44-46.6 °C.

15

20

Example 216 (General procedure (C))

5-(4-Nitroindol-3-ylmethylene)-thiazolidine-2,4-dione




HPLC-MS (Method A): m/z: 290 (M+1); Rt = 3.14 min.

25


Example 217 (General procedure (C))

5-(3,5-Dibromo-4-hydroxy-benzylidene)thiazolidine-2,4-dione



- 5 ^1H NMR ($\text{DMSO}-d_6$): $\delta_{\text{H}} = 12.65$ (broad, 1H), 10.85 (broad, 1H), 7.78 (s, 2H), 7.70 (s, 1H);
HPLC-MS (Method A): m/z : 380 ($M+1$); $R_t = 3.56$ min.

Example 218 (General procedure (C))



- 10 HPLC-MS (Method A): m/z : 385 ($M+1$); $R_t = 5.08$ min.


General procedure for preparation of starting materials for examples 218 - 221:

- Indole-3-carbaldehyde (3.8 g, 26 mmol) was stirred with potassium hydroxide (1.7 g) in acetone (200 mL) at RT until a solution was obtained indicating full conversion to the indole potassium salt. Subsequently the solution was evaporated to dryness *in vacuo*. The residue was dissolved in acetone to give a solution containing 2.6 mmol/20 mL.

- 20 mL portions of this solution were mixed with equimolar amounts of arylmethylbromides in acetone (10 mL). The mixtures were stirred at RT for 4 days and subsequently evaporated to dryness and checked by HPLC-MS. The crude products, 1-benzylated indole-3-carbaldehydes, were used for the reaction with thiazolidine-2,4-dione using the general procedure C.

Example 219 (General procedure (C))


- 25 4-[3-(2,4-Dioxothiazolidin-5-ylidenemethyl)indol-1-ylmethyl]benzoic acid methyl ester



HPLC-MS (Method A): m/z : 393 ($M+1$); R_t = 4.60 min.

Example 220 (General procedure (C))


- 5 5-[1-(9,10-Dioxo-9,10-dihydroanthracen-2-ylmethyl)-1*H*-indol-3-ylmethylene]thiazolidine-2,4-dione



HPLC-MS (Method A): m/z : 465 ($M+1$); R_t = 5.02 min.

- 10 Example 221 (General procedure (C))


4'-[3-(2,4-Dioxothiazolidin-5-ylidenemethyl)indol-1-ylmethyl]biphenyl-2-carbonitrile



HPLC-MS (Method A): m/z : 458 ($M+23$); R_t = 4.81 min.

- 15 Example 222 (General procedure (C))

3-[3-(2,4-Dioxothiazolidin-5-ylidenemethyl)-2-methylindol-1-ylmethyl]benzonitrile.



2-Methylindole-3-carbaldehyde (200 mg, 1.26 mmol) was added to a slurry of 3-bromomethylbenzenecarbonitrile (1.26 mmol) followed by sodium hydride, 60%, (1.26 mmol) in DMF (2 mL). The mixture was shaken for 16 hours, evaporated to dryness and washed


with water and ethanol. The residue was treated with thiazolidine-2,4-dione following the general procedure C to afford the title compound (100 mg).

HPLC-MS (Method C): m/z: 374 (M+1); Rt. = 3.95 min.

5

Example 223 (General procedure (C))

5-(1-Benzyl-2-methylindol-3-ylmethylene)thiazolidine-2,4-dione.




10 This compound was prepared in analogy with the compound described in example 222 from benzyl bromide and 2-methylindole-3-carbaldehyde, followed by reaction with thiazolidine-2,4-dione resulting in 50 mg of the title compound.

HPLC-MS (Method C): m/z: 349 (M+1); Rt. = 4.19 min.

15 Example 224

4-[3-(2,4-Dioxothiazolidin-5-ylidenemethyl)-2-methylindol-1-ylmethyl]benzoic acid methyl ester




20 This compound was prepared in analogy with the compound described in example 222 from 4-(bromomethyl)benzoic acid methyl ester and 2-methylindole-3-carbaldehyde, followed by reaction with thiazolidine-2,4-dione.

HPLC-MS (Method C): m/z: 407 (M+1); Rt.= 4.19 min.

Example 225 (General procedure (C))

5-(2-Chloro-1-methyl-1H-indol-3-ylmethylene)thiazolidine-2,4-dione




HPLC-MS (Method A): m/z: 293 (M+1); Rt = 4.10 min.

5

Example 226 (General procedure (C))

5-(4-Hydroxy-3,5-diiodo-benzylidene)-thiazolidine-2,4-dione




HPLC-MS (Method A): m/z: 474 (M+1); Rt = 6.61 min.

10

Example 227 (General procedure (C))

5-(4-Hydroxy-3-iodobenzylidene)thiazolidine-2,4-dione




HPLC-MS (Method C): m/z: 348 (M+1); Rt. = 3.13 min

15 ¹H-NMR: (DMSO-*d*₆): 11.5 (1H,broad); 7.95(1H,d); 7.65(1H,s); 7.45 (1H,dd); 7.01(1H,dd); 3.4 (1H,broad).

Example 228 (General procedure (C))


20 5-(2,3,6-Trichlorobenzylidene)thiazolidine-2,4-dione



H PLC-MS (Method C): m/z: 309 (M+1); Rt.= 4.07 min

5 Example 229 (General procedure (C))

5-(2,6-Dichlorobenzylidene)thiazolidine-2,4-dione




Mp. 152-154°C.

HPLC-MS (Method C): m/z: 274 (M+1), Rt.= 3.70 min

10 ¹H-NMR: (DMSO-*d*₆): 12.8 (1H, broad); 7.72 (1H,s); 7.60 (2H,d); 7.50 (1H,t).

Example 230 (General procedure (C))

5-[1-(2,6-Dichloro-4-trifluoromethylphenyl)-2,5-dimethyl-1*H*-pyrrol-3-ylmethylene]thiazolidine-2,4-dione




15

HPLC-MS (Method C): m/z: 436 (M+1); Rt. 4.81 min

Example 231 (General procedure (C))

5-[1-(3,5-Dichlorophenyl)-5-(4-methanesulfonylphenyl)-2-methyl-1*H*-pyrrol-3-ylmethylene]-thiazolidine-2,4-dione


20



HPLC-MS (Method C): m/z : 508 ($M+1$); R_t . = 4.31 min

Example 232 (General procedure (C))


- 5 5-[1-(2,5-Dimethoxyphenyl)-5-(4-methanesulfonylphenyl)-2-methyl-1*H*-pyrrol-3-ylmethylene]-thiazolidine-2,4-dione



HPLC-MS (Method C): m/z : 499 ($M+1$); R_t . = 3.70 min

- 10 Example 233 (General procedure (C))


4-[3-(2,4-Dioxothiazolidin-5-ylidenemethyl)-2,5-dimethylpyrrol-1-yl]benzoic acid



HPLC-MS (Method C): m/z : 342 ($M+1$); R_t . = 3.19 min

- 15 Example 234 (General procedure (C))


5-(4-Hydroxy-2,6-dimethoxybenzylidene)thiazolidine-2,4-dione



HPLC-MS (Method C): m/z : 282 ($M+1$); R_t . = 2.56, mp =331-333 °C

Example 235 (General procedure (C))

5-(2,6-Dimethylbenzylidene)thiazolidine-2,4-dione




M.p: 104-105 °C

5 HPLC-MS (Method C): m/z: 234 (M+1); Rt.= 3.58 min,

Example 236 (General procedure (C))

5-(2,6-Dimethoxybenzylidene)thiazolidine-2,4-dione




10 Mp: 241-242 °C

HPLC-MS (Method C): m/z: 266 (M+1); Rt.= 3.25 min;

Example 237 (General procedure (C))

5-[4-(2-Fluoro-6-nitrobenzyloxy)-2,6-dimethoxybenzylidene]thiazolidine-2,4-dione




15

Mp: 255-256 °C

HPLC-MS (Method C): m/z: 435 (M+1), Rt 4.13 min,

Example 238 (General procedure (C))


20 5-Benzofuran-2-ylmethylenethiazolidine-2,4-dione



HPLC-MS (Method C): m/z:246 (M+1); Rt.= 3.65 min, mp = 265-266 °C .

Example 239 (General procedure (C))


5-[3-(4-Dimethylaminophenyl)allylidene]thiazolidine-2,4-dione



- 5 HPLC-MS (Method C): m/z : 276 ($M+1$); R_t = 3.63, mp = 259-263 °C
 $^1\text{H-NMR}$: ($\text{DMSO}-d_6$) δ = 12.3 (1H, broad); 7.46 (2H, d); 7.39 (1H, d); 7.11 (1H, d); 6.69 (2H, d); 6.59 (1H, dd); 2.98 (3H, s).

Example 240 (General procedure (C))

- 10 5-(2-Methyl-3-phenylallylidene)thiazolidine-2,4-dione




Mp: 203-210 °C

HPLC-MS (Method C): m/z : 246 ($M+1$); R_t = 3.79 min.

- 15 Example 241 (General procedure (C))

5-(2-Chloro-3-phenylallylidene)thiazolidine-2,4-dione




Mp: 251-254 °C

HPLC-MS (Method C): m/z : 266 ($M+1$); R_t = 3.90 min

20

Example 242 (General procedure (C))

5-(2-Oxo-1,2-dihydroquinolin-3-ylmethylene)thiazolidine-2,4-dione




- 25 Mp: 338-347 °C

HPLC-MS (Method C): m/z: 273 (M+1); Rt. = 2.59 min.

Example 243 (General procedure (C))


5-(2,4,6-Tribromo-3-hydroxybenzylidene)thiazolidine-2,4-dione.



HPLC-MS (Method C): m/z: 459 (M+1); Rt. = 3.65 min.

Example 244 (General procedure (C))


5-(5-Bromo-2-methylindol-3-ylmethylene)thiazolidine-2,4-dione.



HPLC-MS (Method C): m/z: 339 (M+1); Rt = 3.37min.

Example 245 (General procedure (C))


5-(7-Bromo-2-methylindol-3-ylmethylene)thiazolidine-2,4-dione.



HPLC-MS (Method C): m/z: 319 (M+1); Rt = 3.48min.

Example 246 (General procedure (C))


5-(6-Bromoindol-3-ylmethylene)thiazolidine-2,4-dione.



HPLC-MS (Method C): m/z: 325 (M+1); Rt = 3.54 min.

Example 247 (General procedure (C))


5-(8-Methyl-2-oxo-1,2-dihydroquinolin-3-ylmethylene)thiazolidine-2,4-dione.



HPLC-MS (Method C): m/z: 287 (M+1); Rt = 2.86 min.

5 Example 248 (General procedure (C))


5-(6-Methoxy-2-oxo-1,2-dihydroquinolin-3-ylmethylene)thiazolidine-2,4-dione.



HPLC-MS (Method C): m/z: 303 (M+1); Rt = 2.65 min.

10 Example 249 (General procedure (C))


5-Quinolin-3-ylmethylenethiazolidine-2,4-dione.



HPLC-MS (Method C): m/z: 257 (M+1); Rt = 2.77 min.

15 Example 250 (General procedure (C))


5-(8-Hydroxyquinolin-2-ylmethylene)thiazolidine-2,4-dione.



HPLC-MS (Method C): m/z: 273 (M+1); Rt = 3.44 min.

20 Example 251 (General procedure (C))

5-Quinolin-8-ylmethylenethiazolidine-2,4-dione.



HPLC-MS (Method C): m/z: 257 (M+1); Rt = 3.15 min.

Example 252 (General procedure (C))


5-(1-Bromo-6-methoxynaphthalen-2-ylmethylene)thiazolidine-2,4-dione.



5 HPLC-MS (Method C): m/z: 366 (M+1); Rt = 4.44 min.

Example 253 (General procedure (C))


5-(6-Methyl-2-oxo-1,2-dihydroquinolin-3-ylmethylene)thiazolidine-2,4-dione.



10 HPLC-MS (Method C): m/z: 287 (M+1); Rt. = 2.89 min.

Example 254 (General procedure (D))


5-(2,6-Dichloro-4-dibenzylaminobenzylidene)thiazolidine-2,4-dione.



15 HPLC-MS (Method C): m/z: 469 (M+1); Rt = 5.35 min.

Example 255 (General Procedure (C))

7-(2,4-Dioxothiazolidin-5-ylidenemethyl)-4-methoxybenzofuran-2-carboxylic acid



20 HPLC-MS (Method C): m/z: 320 (M+1); Rt = 2.71 min.

Preparation of the intermediate, 7-formyl-4-methoxybenzofuran-2-carboxylic acid:

A mixture of 2-hydroxy-6-methoxybenzaldehyde (6.4 g, 42 mmol), ethyl bromoacetate (14.2 mL, 128 mmol) and potassium carbonate (26 g, 185 mmol) was heated to 130 °C. After 3 h the mixture was cooled to room temperature and acetone (100 mL) was added, the mixture was subsequently filtered and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel eluting with a mixture of ethyl acetate and heptane (1:4). This afforded 7.5 g (55%) of ethyl 4-methoxybenzofuran-2-carboxylate.


A solution of ethyl 4-methoxybenzofuran-2-carboxylate (6.9 g, 31.3 mmol) in dichloromethane (70 ml) was cooled to 0 °C and a solution of titanium tetrachloride (13.08 g, 69 mmol) was added drop wise. After 10 minutes dichloromethoxymethane (3.958 g, 34 mmol) was added over 10 minutes. After addition, the mixture was warmed to room temperature for 18 hours and the mixture poured into hydrochloric acid (2N, 100 mL). The mixture was stirred for 0.5 hour and then extracted with a mixture of ethyl acetate and toluene (1:1). The organic phase was dried over Na₂SO₄ and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel eluting with a mixture of ethyl acetate and heptane (1:4). This afforded 5.8 g (80%) of ethyl 7-formyl-4-methoxybenzofuran-2-carboxylate.

7-formyl-4-methoxybenzofuran-2-carboxylate (5.0 g, 21.5 mmol) and sodium carbonate (43 mmol) in water (100 mL) was refluxed until a clear solution appeared (about 0.5 hour). The solution was filtered and acidified to pH = 1 with hydrochloric acid (2 N), the resulting product was filtered off and washed with ethyl acetate and ethanol and dried to afford 3.5 g (74%) of 7-formyl-4-methoxybenzofuran-2-carboxylic acid as a solid.

¹H NMR (DMSO-*d*₆): δ = 10.20 (s, 1H) ; 8.07 (d, 1H) ; 7.70 (s, 1H) ; 7.17 (d, 1H) ; 4.08 (s, 3H).

Example 256 (General Procedure (C))

5-(4-Methoxybenzofuran-7-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method C): m/z: 267 (M+1); Rt = 3.30 min.

Preparation of the intermediate, 4-methoxybenzofuran-7-carbaldehyde:


5 A mixture of 7-formyl-4-methoxybenzofuran-2-carboxylic acid (3.0 g, 13.6 mmol) and Cu (0.6 g, 9.44 mmol) in quinoline (6 mL) was refluxed. After 0.5 h the mixture was cooled to room temperature and water (100 mL) and hydrochloric acid (10 N, 20 mL) were added. The mixture was extracted with a mixture of ethyl acetate and toluene (1:1), filtered through celite and the organic layer separated and washed with a sodium carbonate solution, dried over Na₂SO₄ and concentrated *in vacuo* to afford 1.5 g crude product. Column chromatography
10 SiO₂, EtOAc/heptanes=1/4 gave 1.1 g (46%) of 4-methoxybenzofuran-7-carbaldehyde as a solid.

¹H NMR (CDCl₃): δ: 10.30 (s,1H) ; 7.85 (d,1H) ; 7.75 (d,1H) ; 6.98 (d,1H) ; 6.87 (d,1H) ; 4.10 (s,3H). HPLC-MS (Method C) :m/z: 177 (M+1); Rt. = 7.65 min.

15

Example 257 (General Procedure (C))

5-(4-Hydroxybenzofuran-7-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method C): m/z: = 262 (M+1); Rt 2.45 min.

20


Preparation of the intermediate, 4-hydroxybenzofuran-7-carbaldehyde

A mixture of 4-methoxybenzofuran-7-carbaldehyde (1.6 g, 9.1 mmol) and pyridine hydrochloride (4.8 g, 41.7mmol) in quinoline (8 mL) was refluxed. After 8 h the mixture was
25 cooled to room temperature and poured into water (100 mL) and hydrochloric acid (2 N) was added to pH = 2. The mixture was extracted with a mixture of ethyl acetate and toluene (1:1), washed with a sodium carbonate solution, dried with Na₂SO₄ and concentrated *in vacuo* to afford 0.8 g crude product. This was purified by column chromatography on silica gel, eluting with a mixture of ethyl acetate and heptane (1:3). This afforded 250 mg of 4-
30 hydroxybenzofuran-7-carbaldehyde as a solid.

^1H NMR ($\text{DMSO}-d_6$): δ = 11.35 (s, broad, 1H) ; 10.15 (s, 1H) ; 8.05 (d, 1H) ; 7.75 (d, 1H) ; 7.10 (d, 1H); 6.83 (d, 1H). HPLC-MS (Method C): m/z : 163 ($M+1$); R_t = 6.36 min.

Example 258 (General Procedure (C))

5 5-(5-Bromo-2,3-dihydrobenzofuran-7-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method C): m/z : 328 ($M+1$); R_t = 3.66 min.

Preparation of the intermediate, 5-bromo-2,3-dihydrobenzofuran-7-carbaldehyde:


- 10 To a cooled (15°C) stirred mixture dihydrobenzofuran (50.9 g, 0.424 mol) in acetic acid (500 mL), a solution of bromine (65.5 mL, 1.27 mol) in acetic acid (200 mL) was added drop wise over 1 hour. After stirring for 18 hours, a mixture of $\text{Na}_2\text{S}_2\text{O}_5$ (150 g) in water (250 mL) was added carefully, and the mixture was concentrated *in vacuo*. Water (200 mL) was added and the mixture was extracted with ethyl acetate containing 10% heptane, dried over Na_2SO_4 and
- 15 concentrated *in vacuo* to give crude 5,7-dibromo-2,3-dihydrobenzofuran which was used as such for the following reaction steps. To a cooled solution (-78°C) of crude 5,7-dibromo-2,3-dihydrobenzofuran (50.7 g, 0.182 mol) in THF (375 mL) a solution of $n\text{-BuLi}$ (2.5 M, 80 mL, 0.200 mol) in hexane was added. After addition, the mixture was stirred for 20 min. DMF (16 mL) was then added drop wise at -78°C . After addition, the mixture was stirred at room
- 20 temperature for 3 h and then the mixture was poured into a mixture of ice water, (500 mL) and hydrochloric acid (10 N, 40 mL) and extracted with toluene, dried over Na_2SO_4 and concentrated *in vacuo*. Column chromatography on silica gel eluting with a mixture of ethyl acetate and heptane (1:4) afforded 23 g of 5-bromo-2,3-dihydrobenzofuran-7-carbaldehyde as a solid.

25

^1H NMR (CDCl_3): δ : 10.18 (s, 1H) ; 7.75 (d, 1H) ; 7.55 (d, 1H) ; 4.80 (t, 2H) ; 3.28 (t, 2H).

Example 259 (General Procedure (C))

5-(4-Cyclohexylbenzylidene)thiazolidine-2,4-dione



HPLC-MS (Method C): m/z: 288 (M+1); Rt = 5.03 min.

Preparation of the intermediate, 4-cyclohexylbenzaldehyde:

5

This compound was synthesized according to a modified literature procedure (*J. Org. Chem.*, **37**, No.24, (1972), 3972-3973).


Cyclohexylbenzene (112.5 g, 0.702 mol) and hexamethylenetetramine (99.3 g, 0.708 mol) were mixed in TFA (375 mL). The mixture was stirred under nitrogen at 90 °C for 3 days.

10 After cooling to room temperature the red-brown mixture was poured into ice-water (3600 ml) and stirred for 1 hour. The solution was neutralized with Na₂CO₃ (2 M solution in water) and extracted with dichloromethane (2.5 L). The organic phase was dried (Na₂SO₄) and the solvent was removed *in vacuo*. The remaining red-brown oil was purified by fractional distillation to afford the title compound (51 g, 39%).

15 ¹H NMR (CDCl₃): δ 9.96 (s, 1H), 7.80 (d, 2H), 7.35 (d, 2H), 2.58 (m, 1H), 1.94-1.70 (m, 5H), 1.51-1.17 (m, 5H)


Other ligands of the invention include

20 3',5'-Dichloro-4'-(2,4-dioxothiazolidin-5-ylidenemethyl)biphenyl-4-carboxylic acid:



Example 260 (General procedure (C))


5-(1-Bromo-6-hydroxynaphthalen-2-ylmethylene)-thiazolidine-2,4-dione



HPLC-MS (Method C): $m/z = 350$ ($M+1$); $R_t = 3.45$ min.

Example 261 (General procedure (C))


5 5-[4-(2-Bromoethoxy)-naphthalen-1-ylmethylene]-thiazolidine-2,4-dione



HPLC-MS (Method C): $m/z = 380$ ($M+1$); $R_t = 3.52$ min.

Example 262 (General procedure (C))


10 5-(2-Methyl-5-nitro-1H-indol-3-ylmethylene)-thiazolidine-2,4-dione



HPLC-MS (Method C): $m/z = 304$ ($M+1$); $R_t = 2.95$ min.

Example 263 (General procedure (C))


15 5-(4-Naphthalen-2-yl-thiazol-2-ylmethylene)-thiazolidine-2,4-dione



HPLC-MS (Method C): $m/z = 339$ (M+1); $R_t = 4.498$ min.

Example 264 (General procedure (C))


5 5-[4-(4-Methoxy-naphthalen-1-yl)-thiazol-2-ylmethylene]-thiazolidine-2,4-dione



HPLC-MS (Method C): $m/z = 369$ (M+1); $R_t = 4.456$ min.

Example 265 (General procedure (C))


10 5-(2-Pyridin-4-yl-1H-indol-3-ylmethylene)-thiazolidine-2,4-dione



HPLC-MS (Method C): $m/z = 322$ (M+1); $R_t = 2.307$ min.

Example 266 (General procedure (C))


5-[5-(4-Chlorophenyl)-1H-pyrazol-4-ylmethylene]-thiazolidine-2,4-dione

HPLC-MS (Method C): m/z = 306 (M+1); R_t = 3.60 min.

5

Example 267 (General procedure (C))


5-[5-(2,5-Dimethylphenyl)-1H-pyrazol-4-ylmethylene]-thiazolidine-2,4-dione

HPLC-MS (Method C): m/z = 300 (M+1); R_t = 3.063 min.

10

Example 268 (General procedure (C))


5-(2-Phenyl-benzo[d]imidazo[2,1-b]thiazol-3-ylmethylene)-thiazolidine-2,4-dione

HPLC-MS (Method C): m/z = 378 (M+1); R_t = 3.90 min.

15

Example 269 (General procedure (C))


N-{4-[2-(2,4-Dioxothiazolidin-5-ylidenemethyl)-phenoxy]-phenyl}-acetamide

HPLC-MS (Method C): m/z = 355 (M+1); Rt. 3.33 min.

5

Example 270 (General procedure (C))


5-(2-Phenyl-imidazo[1,2-a]pyridin-3-ylmethylene)-thiazolidine-2,4-dione

HPLC-MS (Method C): m/z = 322 (M+1); Rt. = 2.78 min.

10

Example 271 (General procedure (C))


5-(2-Naphthalen-2-yl-imidazo[1,2-a]pyridin-3-ylmethylene)-thiazolidine-2,4-dione

HPLC-MS (Method C): m/z = 372 (M+1); Rt. = 2.78 min.

15

Example 272 (General procedure (C))


5-[6-Bromo-2-(3-methoxyphenyl)-imidazo[1,2-a]pyridin-3-ylmethylene]-thiazolidine-2,4-dione



HPLC-MS (Method C): $m/z = 431$ ($M+1$); $R_t = 3.30$ min.

5 Example 273 (General procedure (C))


5-(1,2,3,4-Tetrahydrophenanthren-9-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method C): $m/z = 310$ ($M+1$); $R_t = 4.97$ min.

10 Example 274 (General procedure (C))


5-(3,5,5,8,8-Pentamethyl-5,6,7,8-tetrahydro-naphthalen-2-ylmethylene)thiazolidine-2,4-dione



HPLC-MS (Method C): $m/z = 330$ ($M+1$); $R_t = 5.33$ min.

Example 275 (General procedure (C))

5-[6-(2,4-Dichloro-phenyl)-imidazo[2,1-b]thiazol-5-ylmethylene]-thiazolidine-2,4-dione

HPLC-MS (Method C): m/z = 396 (M+1); Rt. = 3.82 min.

5

Example 276 (General procedure (C))


5-(5-Bromobenzofuran-7-ylmethylene)-thiazolidine-2,4-dione

HPLC-MS (Method C): m/z = 324 (M+1); Rt. = 3.82 min.

10

Example 277 (General procedure (C))

4-[3-(2,4-Dioxothiazolidin-5-ylidenemethyl)-1,4-dimethylcarbazol-9-ylmethyl]-benzoic acid

HPLC-MS (Method C): m/z = 457 (M+1); Rt = 4,23 min.

15


Preparation of intermediary aldehyde:

1,4 Dimethylcarbazol-3-carbaldehyde (0.68 g, 3.08 mmol) was dissolved in dry DMF (15 mL), NaH (diethyl ether washed) (0.162 g, 6.7 mol) was slowly added under nitrogen and the mixture was stirred for 1 hour at room temperature. 4-Bromomethylbenzoic acid (0.73 g, 3.4 mmol) was slowly added and the resulting slurry was heated to 40 °C for 16 hours. Water (5 mL) and hydrochloric acid (6N, 3 mL) were added. After stirring for 20 min at room temperature, the precipitate was filtered off and washed twice with acetone to afford after drying 0.38 g (34%) of 4-(3-formyl-1,4-dimethylcarbazol-9-ylmethyl)benzoic acid.

10 HPLC-MS (Method C) : m/z = 358 (M+1), RT. = 4.15 min.

Example 278 (General procedure (C))


4-[7-(2,4-Dioxothiazolidin-5-ylidenemethyl)-benzofuran-5-yl]-benzoic acid



15 Starting aldehyde commercially available (Syncom BV, NL)
HPLC-MS (Method C): m/z = 366 (M+1); Rt. = 3.37 min.

Example 279 (General procedure (C))


4-[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)-2-nitrophenoxy]-benzoic acid methyl ester



20 HPLC-MS (Method C): m/z = 401 (M+1); Rt. = 4.08 min.

Example 280 (General procedure (C))


3',5'-Dichloro-4'-(2,4-dioxothiazolidin-5-ylidenemethyl)-biphenyl-4-carboxylic acid



5 Starting aldehyde commercially available (Syncom BV, NL)

HPLC-MS (Method C): $m/z = 394$ ($M+1$); $R_t = 3.71$ min.

Example 281 (General procedure (C))




10

HPLC-MS (Method C): $m/z = 232$ ($M+1$); $R_t = 3.6$ min.

Example 282

5-(2-Methyl-1H-indol-3-ylmethyl)-thiazolidine-2,4-dione



15


5-(2-Methyl-1H-indol-3-ylmethylene)thiazolidine-2,4-dione (prepared as described in example 187, 1.5 g, 5.8 mmol) was dissolved in pyridine (20 mL) and THF (50 mL), LiBH_4 (2 M in THF, 23.2 mmol) was slowly added with a syringe under cooling on ice. The mixture was heated to 85 °C for 2 days. After cooling, the mixture was acidified with concentrated

hydrochloric acid to pH 1. The aqueous layer was extracted 3 times with ethyl acetate, dried with MgSO_4 treated with activated carbon, filtered and the resulting filtrate was evaporated *in vacuo* to give 1.3 g (88%) of the title compound.

- 5 HPLC-MS (Method C): $m/z = 261$ ($M+1$); $R_t = 3.00$ min.

Example 283

4-[4-(2,4-Dioxothiazolidin-5-ylmethyl)naphthalen-1-yloxy]butyric acid



- 10 4-[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)naphthalen-1-yloxy]butyric acid (4.98 g, 13.9 mmol, prepared as described in example 469) was dissolved in dry THF (50 mL) and added dry pyridine (50 mL) and, in portions, lithium borohydride (2.0 M, in THF, 14 mL). The resulting slurry was refluxed under nitrogen for 16 hours, added (after cooling) more lithium borohydride (2.0 M, in THF, 7 mL). The resulting mixture was refluxed under nitrogen for 16
- 15 hours. The mixture was cooled and added more lithium borohydride (2.0 M, in THF, 5 mL). The resulting mixture was refluxed under nitrogen for 16 hours. After cooling to 5 °C, the mixture was added water (300 mL) and hydrochloric acid (150 mL). The solid was isolated by filtration, washed with water (3 x 500 mL) and dried. Recrystallization from acetonitrile (500 mL) afforded 2.5 g of the title compound.

20


$^1\text{H-NMR}$ ($\text{DMSO}-d_6$, selected peaks): $\delta = 3.42$ (1H, dd), 3.90 (1H, dd), 4.16 (2H, "t"), 4.95 (1H, dd), 6.92 (1H, d), 7.31 (1H, d), 7.54 (1H, t), 7.62 (1H, t), 8.02 (1H, d), 8.23 (1H, d), 12.1 (1H, bs), 12.2 (1H, bs).

HPLC-MS (Method C): $m/z = 382$ ($M+23$); $R_t = 3.23$ min.

25

Example 284

5-Naphthalen-1-ylmethylthiazolidine-2,4-dione












5-Naphthalen-1-ylmethylene-2,4-dione (1.08 g, 4.2 mmol, prepared as described in example 68) was dissolved in dry THF (15 mL) and added dry pyridine (15 mL) and, in portions, lithium borohydride (2.0 M, in THF, 4.6 mL). The resulting mixture was refluxed under nitrogen for 16 hours. After cooling to 5 °C, the mixture was added water (100 mL), and, in portions, concentrated hydrochloric acid (40 mL). More water (100 mL) was added, and the mixture was extracted with ethyl acetate (200 mL). The organic phase was washed with water (3 x 100 mL), dried and concentrated *in vacuo*. The residue was dissolved in ethyl acetate (50 mL) added activated carbon, filtered and concentrated *in vacuo* and dried to afford 0.82 g (75%) of the title compound.









¹H-NMR (DMSO-*d*₆): δ = 3.54 (1H, dd), 3.98 (1H, dd), 5.00 (1H, dd), 7.4-7.6 (4H, m), 7.87 (1H, d), 7.96 (1H, d), 8.11 (1H, d), 12.2 (1H, bs).









HPLC-MS (Method C): *m/z* = 258 (M+1); *R*_t = 3.638 min.









15 The following preferred compounds of the invention may be prepared according to procedures similar to those described in the three examples above:









| | | |
|-------------|--|--|
| Example 285 | | |
| Example 286 | | |
| Example 287 | | |









| | | |
|-------------|--|--|
| Example 288 |  | |
| Example 289 |  | |
| Example 290 |  | |
| Example 291 |  | |
| Example 292 |  | |
| Example 293 |  | |
| Example 294 |  | |
| Example 295 |  | |
| Example 296 |  | |









| | | |
|-------------|--|--|
| Example 297 |  | |
| Example 298 |  | |
| Example 299 |  | |
| Example 300 |  | |
| Example 301 |  | |
| Example 302 |  | |
| Example 303 |  | |
| Example 304 |  | |









| | | |
|-------------|--|--|
| Example 305 |  | |
| Example 306 |  | |
| Example 307 |  | |
| Example 308 |  | |
| Example 309 |  | |
| Example 310 |  | |
| Example 311 |  | |
| Example 312 |  | |








| | | |
|-------------|---|--|
| Example 313 |  | |
| Example 314 |  | |
| Example 315 |  | |
| Example 316 |  | |
| Example 317 |  | |
| Example 318 |  | |
| Example 319 |  | |
| Example 320 |  | |








| | | |
|-------------|---|--|
| Example 321 |  <chem>CCCC(=O)c1ccc(cc1)Oc2cc(ccc2)SC(=O)NC(=O)c3ccccc3</chem> | |
| Example 322 |  <chem>O=C1NC(=O)SC1=NC(=O)SC(=O)Nc2ccc(cc2)Cc3ccccc3SC(=O)NC(=O)c4ccccc4</chem> | |
| Example 323 |  <chem>CCCCCCCCOCc1ccc(cc1)Cc2ccccc2SC(=O)NC(=O)c3ccccc3</chem> | |
| Example 324 |  <chem>O=C1NC(=O)SC1=NC(=O)SC(=O)Nc2ccc(cc2)Cc3ccc4ccccc4c3OC(=O)O</chem> | |
| Example 325 |  <chem>O=C1NC(=O)SC1=NC(=O)SC(=O)Nc2ccc(cc2)Cc3ccc4ccccc4c3OC(=O)O</chem> | |
| Example 326 |  <chem>O=C1NC(=O)SC1=NC(=O)SC(=O)Nc2ccc(cc2)Cc3ccc4ccccc4c3OC(=O)O</chem> | |
| Example 327 |  <chem>CCOC(=O)CCCCOCc1ccc2ccccc2c1Cc3ccccc3SC(=O)NC(=O)c4ccccc4</chem> | |
| Example 328 |  <chem>O=C1NC(=O)SC1=NC(=O)SC(=O)Nc2ccc(cc2)Cc3ccc4ccccc4c3OC(=O)O</chem> | |








| | | |
|-------------|--|--|
| | | |
| Example 329 |  | |
| Example 330 |  | |
| Example 331 |  | |
| Example 332 |  | |
| Example 333 |  | |
| Example 334 |  | |
| Example 335 |  | |
| Example 336 |  | |







| | | |
|-------------|--|--|
| Example 337 |  | |
| Example 338 |  | |
| Example 339 |  | |
| Example 340 |  | |
| Example 341 |  | |
| Example 342 |  | |
| Example 343 |  | |
| Example 344 |  | |

| | | |
|-------------|--|--|
| Example 345 |  | |
| Example 346 |  | |
| Example 347 |  | |
| Example 348 |  | |
| Example 349 |  | |
| Example 350 |  | |
| Example 351 |  | |
| Example 352 |  | |

| | | |
|-------------|--|--|
| Example 353 |  | |
| Example 354 |  | |
| Example 355 |  | |
| Example 356 |  | |
| Example 357 |  | |
| Example 358 |  | |
| Example 359 |  | |

| | | |
|-------------|--|--|
| Example 360 |  | |
| Example 361 |  | |
| Example 362 |  | |
| Example 363 |  | |
| Example 364 |  | |
| Example 365 |  | |
| Example 366 |  | |


| | | |
|-------------|--|--|
| Example 367 |  | |
| Example 368 |  | |
| Example 369 |  | |
| Example 370 |  | |
| Example 371 |  | |
| Example 372 |  | |
| Example 373 |  | |

| | | |
|-------------|---|--|
| Example 374 |  | |
| Example 375 |  | |
| Example 376 |  | |
| Example 377 |  | |
| Example 378 |  | |
| Example 379 |  | |

The following compounds are commercially available and may be prepared using general procedures (B) and / or (C).


Example 380

- 5 5-(5-Bromo-1H-indol-3-ylmethylene)thiazolidine-2,4-dione




Example 381

5-Pyridin-4-ylmethylenethiazolidine-2,4-dione




Example 382

5 5-(3-Bromo-4-methoxybenzylidene)thiazolidine-2,4-dione




Example 383

5-(3-Nitrobenzylidene)thiazolidine-2,4-dione




Example 384

5-Cyclohexylidene-1,3-thiazolidine-2,4-dione




Example 385

5-(3,4-Dihydroxybenzylidene)thiazolidine-2,4-dione




Example 386

5-(3-Ethoxy-4-hydroxybenzylidene)thiazolidine-2,4-dione




Example 387

5-(4-Hydroxy-3-methoxy-5-nitrobenzylidene)thiazolidine-2,4-dione




Example 388

5-(3-Ethoxy-4-hydroxybenzylidene)thiazolidine-2,4-dione




Example 389

5-(4-Hydroxy-3,5-dimethoxybenzylidene)thiazolidine-2,4-dione




15 Example 390

5-(3-Bromo-5-ethoxy-4-hydroxybenzylidene)thiazolidine-2,4-dione




Example 391


5-(3-Ethoxy-4-hydroxy-5-nitrobenzylidene)thiazolidine-2,4-dione



5 Example 392




Example 393




10

Example 394




Example 395




15


Example 396




5 Example 397




Example 398




Example 399




15 Example 400




Example 401



5 Example 402




Example 403




10

Example 404




Example 405

15 5-(3-Hydroxy-5-methyl-phenylamino)-thiazolidine-2,4-dione




Example 406




5

Example 407




Example 408




10


Example 409




Example 410




Example 411




10 Example 412




Example 413




Example 414




Example 415




10 Example 416



Example 417




Example 418




5


Example 419



10 Example 420




Example 421




15

Example 422




Example 423




5


Example 424



10 Example 425




Example 426




15

Example 427




Example 428




5

Example 429




Example 430




10

Example 431

5-(4-Diethylamino-2-methoxy-benzylidene)-imidazolidine-2,4-dione




Example 432




5

Example 433




Example 434




10

Example 435




Example 436




5

Example 437




Example 438




10

Example 439




Example 440




5

Example 441




Example 442




10

Example 443




Example 444




5


Example 445




10 Example 446



Example 447




Example 448




5

Example 449




Example 450




10

Example 451




Example 452




5

Example 453



Example 454


10 5-(4-Diethylamino-benzylidene)-2-imino-thiazolidin-4-one




Example 455




Example 456



5 Example 457





Example 458



10

Example 459



General procedure (D) for preparation of compounds of general formula**I₃:**

5

wherein X, Y, and R³ are as defined above,

n is 1 or 3-20,

E is arylene or heterarylene (including up to four optional substituents, R¹³, R¹⁴, R¹⁵, and R^{15A} as defined above),

- 10 R' is a standard carboxylic acid protecting group, such as C₁-C₆-alkyl or benzyl and Lea is a leaving group, such as chloro, bromo, iodo, methanesulfonyloxy, toluenesulfonyloxy or the like.

- 15 Step 1 is an alkylation of a phenol moiety. The reaction is preformed by reacting R¹⁰-C(=O)-E-OH with an ω-bromo-alkane-carboxylic acid ester (or a synthetic equivalent) in the presence of a base such as sodium or potassium carbonate, sodium or potassium hydroxide, sodium hydride, sodium or potassium alkoxide in a solvent, such as DMF, NMP, DMSO, acetone, acetonitrile, ethyl acetate or isopropyl acetate. The reaction is performed at 20 – 160 °C, usually at room temperature, but when the phenol moiety has one or more
- 20 substituents heating to 50 °C or more can be beneficial, especially when the substituents are in the ortho position relatively to the phenol. This will readily be recognised by those skilled in the art.

Step 2 is a hydrolysis of the product from step 1.


Step 3 is similar to general procedure (B) and (C).

This general procedure (D) is further illustrated in the following examples:

5

Example 460 (General procedure (D))

4-[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)phenoxy]butyric acid



Step 1:

- 10 A mixture of 4-hydroxybenzaldehyde (9.21 g, 75 mmol), potassium carbonate (56 g, 410 mmol) and 4-bromobutyric acid ethyl ester (12.9 mL, 90 mmol) in *N,N*-dimethylformamide (250 mL) was stirred vigorously for 16 hours at room temperature. The mixture was filtered and concentrated *in vacuo* to afford 19.6 g (100%) of 4-(4-formylphenoxy)butyric acid ethyl ester as an oil. ¹H-NMR (DMSO-*d*₆): δ 1.21 (3H, t), 2.05 (2H, p), 2.49 (2H, t), 4.12 (4H, m),
15 7.13 (2H, d), 7.87 (2H, d), 9.90 (1H, s). HPLC-MS (Method A): m/z = 237 (M+1); R_t = 3.46 min.

Step 2:

- 4-(4-Formylphenoxy)butyric acid ethyl ester (19.6 g, 75 mmol) was dissolved in methanol
20 (250 mL) and 1N sodium hydroxide (100 mL) was added and the resulting mixture was stirred at room temperature for 16 hours. The organic solvent was evaporated *in vacuo* (40 °C, 120 mBar) and the residue was acidified with 1N hydrochloric acid (110 mL). The mixture was filtered and washed with water and dried *in vacuo* to afford 14.3 g (91%) 4-(4-formylphenoxy)butyric acid as a solid. ¹H-NMR (DMSO-*d*₆): δ 1.99 (2H, p), 2.42 (2H, t), 4.13
25 (2H, t), 7.14 (2H, d), 7.88 (2H, d), 9.90 (1H, s), 12.2 (1H, bs). HPLC-MS (Method A): m/z = 209 (M+1); R_t = 2.19 min.

Step 3:

- Thiazolidine-2,4-dione (3.55 g, 27.6 mmol), 4-(4-formylphenoxy)butyric acid (5.74 g, 27.6
30 mmol), anhydrous sodium acetate (11.3 g, 138 mmol) and acetic acid (100 mL) was refluxed for 16 h. After cooling, the mixture was filtered and washed with acetic acid and water. Drying


in vacuo afforded 2.74 g (32%) of 4-[4-(2,4-dioxothiazolidin-5-ylidenemethyl)phenoxy]butyric acid as a solid.

¹H-NMR (DMSO-*d*₆): δ 1.97 (2H, p), 2.40 (2H, t), 4.07 (2H, t), 7.08 (2H, d), 7.56 (2H, d), 7.77 (1H, s), 12.2 (1H, bs), 12.5 (1H, bs); HPLC-MS (Method A): *m/z*: 308 (M+1); *R*_t = 2.89 min.

5

Example 461 (General procedure (D))

[3-(2,4-Dioxothiazolidin-5-ylidenemethyl)phenoxy]acetic acid



Step 3:


10 Thiazolidine-2,4-dione (3.9 g, 33 mmol), 3-formylphenoxyacetic acid (6.0 g, 33 mmol), anhydrous sodium acetate (13.6 g, 165 mmol) and acetic acid (100 mL) was refluxed for 16 h. After cooling, the mixture was filtered and washed with acetic acid and water. Drying in vacuo afforded 5.13 g (56%) of [3-(2,4-dioxothiazolidin-5-ylidenemethyl)phenoxy]acetic acid as a solid.

15 ¹H-NMR (DMSO-*d*₆): δ 4.69 (2H, s), 6.95 (1H, dd), 7.09 (1H, t), 7.15 (1H, d), 7.39 (1H, t), 7.53 (1H, s); HPLC-MS (Method A): *m/z* = 280 (M+1) (poor ionisation); *R*_t = 2.49 min.

The compounds in the following examples were similarly prepared.

20 Example 462 (General procedure (D))


3-[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)phenyl]acrylic acid



¹H-NMR (DMSO-*d*₆): δ 6.63 (1H, d), 7.59-7.64 (3H, m), 7.77 (1H, s), 7.83 (2H, m).

25 Example 463 (General procedure (D))

[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)phenoxy]acetic acid




Triethylamine salt: $^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 4.27 (2H, s), 6.90 (2H, d), 7.26 (1H, s), 7.40 (2H, d).

5


Example 464 (General procedure (D))

4-(2,4-Dioxothiazolidin-5-ylidenemethyl)benzoic acid



10 Example 465 (General procedure (D))


3-(2,4-Dioxothiazolidin-5-ylidenemethyl)benzoic acid



$^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 7.57 (1H, s), 7.60 (1H, t), 7.79 (1H, dt), 7.92 (1H, dt), 8.14 (1H, t).

15 Example 466 (General procedure (D))

4-[2-Chloro-4-(2,4-dioxothiazolidin-5-ylidenemethyl)phenoxy]butyric acid




$^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 2.00 (2H, p), 2.45 (2H, t), 4.17 (2H, t), 7.31 (1H, d), 7.54 (1H, dd), 7.69 (1H, d), 7.74 (1H, s), 12.2 (1H, bs), 12.6 (1H, bs). HPLC-MS (Method A): m/z : 364 ($M+23$); R_t = 3.19 min.

20

Example 467 (General procedure (D))

4-[2-Bromo-4-(2,4-dioxothiazolidin-5-ylidenemethyl)phenoxy]butyric acid




$^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 1.99 (2H, p), 2.46 (2H, t), 4.17 (2H, t), 7.28 (1H, d), 7.57 (1H, dd), 7.25 (1H, s), 7.85 (1H, d), 12.2 (1H, bs), 12.6 (1H, bs). HPLC-MS (Method A): m/z : 410 ($M+23$); R_t = 3.35 min.

5

Example 468 (General procedure (D))

4-[2-Bromo-4-(4-oxo-2-thioxothiazolidin-5-ylidenemethyl)phenoxy]butyric acid




$^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 1.99 (2H, p), 2.45 (2H, t), 4.18 (2H, t), 7.28 (1H, d), 7.55 (1H, dd), 7.60 (1H, s), 7.86 (1H, d), 12.2 (1H, bs), 13.8 (1H, bs). HPLC-MS (Method A): m/z : 424 ($M+23$); R_t = 3.84 min.

10

HPLC-MS (Method A): m/z : 424 ($M+23$); R_t = 3.84 min

Example 469 (General procedure (D))

15 4-[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)naphthalen-1-yloxy]butyric acid




$^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 2.12 (2H, p), 2.5 (below DMSO), 4.28 (2H, t), 7.12 (1H, d), 7.6-7.7 (3H, m), 8.12 (1H, d), 8.31 (1H, d), 8.39 (1H, s), 12.2 (1H, bs), 12.6 (1H, bs). HPLC-MS (Method A): m/z : 380 ($M+23$); R_t = 3.76 min.

20

Example 470 (General procedure (D))

5-[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)naphthalen-1-yloxy]pentanoic acid




HPLC-MS (Method A): m/z: 394 (M+23); Rt = 3.62 min.

¹H-NMR (DMSO-*d*₆): δ 1.78 (2H, m), 1.90 (2H, m), 2.38 (2H, t), 4.27 (2H, t), 7.16 (1H, d), 7.6-7.75 (3H, m), 8.13 (1H, d), 8.28 (1H, d), 8.39 (1H, s), 12.1 (1H, bs), 12.6 (1H, bs).

5

Example 471

5-[2-Bromo-4-(2,4-dioxothiazolidin-5-ylidenemethyl)naphthalen-1-yloxy]pentanoic acid.




- 10 5-[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)-naphthalen-1-yloxy]pentanoic acid (example 470, 185 mg, 0.5 mmol) was treated with an equimolar amount of bromine in acetic acid (10 mL). Stirring at RT for 14 days followed by evaporation to dryness afforded a mixture of the brominated compound and unchanged starting material. Purification by preparative HPLC on a C18 column using acetonitrile and water as eluent afforded 8 mg of the title compound.

15

HPLC-MS (Method C): m/z: 473 (M+23), Rt. = 3.77 min

Example 472

4-[2-Bromo-4-(2,4-dioxothiazolidin-5-ylidenemethyl)naphthalen-1-yloxy]butyric acid.




20

Starting with 4-[4-(2,4-dioxothiazolidin-5-ylidenemethyl)-naphthalen-1-yloxy]-butyric acid (example 469, 0.5 mmol) using the same method as in example 471 afforded 66 mg of the title compound.

- 25 HPLC-MS (Method C): m/z: 459 (M+23) ; Rt. = 3.59 min.

Example 473 (General procedure (D))


[2-Bromo-4-(2,4-dioxothiazolidin-5-ylidenemethyl)phenoxy]acetic acid



- 5 $^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 4.90 (2H, s), 7.12 (1H, d), 7.52 (1H, dd), 7.65 (1H, s) 7.84 (1H, d). HPLC-MS (Method A): m/z : not observed; R_t = 2.89 min.

Example 474 (General procedure (D))

4-[3-(2,4-Dioxothiazolidin-5-ylidenemethyl)phenoxy]butyric acid




10

- $^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 1.98 (2H, p), 2.42 (2H, t), 4.04 (2H, t), 7.05 (1H, dd), 7.15 (2H, m), 7.45 (1H, t), 7.77 (1H, s), 12.1 (1H, bs), 12.6 (1H, bs). HPLC-MS (Method A): m/z : 330 ($M+23$); R_t = 3.05 min.

15 Example 475 (General procedure (D))


[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)-3-methoxyphenoxy]acetic acid

HPLC-MS (Method B): m/z : 310 ($M+1$); R_t = 3,43 min.

20

Example 476 (General procedure (D))


[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)naphthalen-1-yloxy]acetic acid



HPLC-MS (Method A): m/z : 330 ($M+1$); R_t = 3.25 min.

Example 477 (General procedure (D))


5 8-(2,4-Dioxothiazolidin-5-ylidenemethyl)naphthalene-1-carboxylic acid



HPLC-MS (Method A): m/z : 299 ($M+1$); R_t = 2.49 min.

Example 478 (General procedure (D))

10 [3-(2,4-Dioxothiazolidin-5-ylidenemethyl)indol-1-yl]acetic acid



HPLC-MS (Method A): m/z : 303 ($M+1$); R_t = 2.90 min.

Preparation of starting material:

- 15 3-Formylindol (10 g, 69 mmol) was dissolved in *N,N*-dimethylformamide (100 mL) and under an atmosphere of nitrogen and with external cooling, keeping the temperature below 15 °C, sodium hydride (60% in mineral oil, 3.0 g, 76 mmol) was added in portions. Then a solution of ethyl bromoacetate (8.4 mL, 76 mmol) in *N,N*-dimethylformamide (15 mL) was added dropwise over 30 minutes and the resulting mixture was stirred at room temperature for 16
- 20 hours. The mixture was concentrated *in vacuo* and the residue was partitioned between water (300 mL) and ethyl acetate (2 x 150 mL). The combined organic extracts were washed with a saturated aqueous solution of ammonium chloride (100 mL), dried ($MgSO_4$) and

concentrated in vacuo to afford 15.9 g (quant.) of (3-formylindol-1-yl)acetic acid ethyl ester as an oil.


¹H-NMR (CDCl₃): δ_H = 1.30 (3H, t), 4.23 (2H, q), 4.90 (2H, s), 7.3 (3H, m), 7.77 (1H, s), 8.32 (1H, d), 10.0 (1H, s).

(3-Formylindol-1-yl)acetic acid ethyl ester (15.9 g 69 mmol) was dissolved in 1,4-dioxane (100 mL) and 1N sodium hydroxide (10 mL) was added and the resulting mixture was stirred at room temperature for 4 days. Water (500 mL) was added and the mixture was washed with diethyl ether (150 mL). The aqueous phase was acidified with 5N hydrochloric acid and extracted with ethyl acetate (250 + 150 mL). The combined organic extracts were dried (MgSO₄) and concentrated in vacuo to afford 10.3 g (73%) of (3-formylindol-1-yl)acetic acid as a solid.

¹H-NMR (DMSO-*d*₆): δ_H = 5.20 (2H, s), 7.3 (2H, m), 7.55 (1H, d), 8.12 (1H, d), 8.30 (1H, s), 9.95 (1H, s), 13.3 (1H, bs).

Example 479 (General procedure (D))

3-[3-(2,4-Dioxothiazolidin-5-ylidenemethyl)indol-1-yl]propionic acid



HPLC-MS (Method A): m/z: 317 (M+1); Rt = 3.08 min.

Preparation of starting material:


A mixture of 3-formylindol (10 g, 69 mmol), ethyl 3-bromopropionate (10.5 mL, 83 mmol) and potassium carbonate (28.5 g, 207 mmol) and acetonitrile (100 mL) was stirred vigorously at reflux temperature for 2 days. After cooling, the mixture was filtered and the filtrate was concentrated *in vacuo* to afford 17.5 g (quant.) of 3-(3-formylindol-1-yl)propionic acid ethyl ester as a solid.

¹H-NMR (DMSO-*d*₆): δ_H = 1.10 (3H, t), 2.94 (2H, t), 4.02 (2H, q), 4.55 (2H, t), 7.3 (2H, m), 7.67 (1H, d), 8.12 (1H, d), 8.30 (1H, s), 9.90 (1H, s).

- 3-(3-Formylindol-1-yl)propionic acid ethyl ester (17.5 g 69 mmol) was hydrolysed as
5 described above to afford 12.5 g (83%) of 3-(3-formylindol-1-yl)propionic acid as a solid.

¹H-NMR (DMSO-*d*₆): δ_H = 2.87 (2H, t), 4.50 (2H, t), 7.3 (2H, m), 7.68 (1H, d), 8.12 (1H, d), 8.31 (1H, s), 9.95 (1H, s), 12.5 (1H, bs).


- 10 Example 480 (General procedure (D))
{5-[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)benzylidene]-4-oxo-2-thioxothiazolidin-3-yl}acetic acid



HPLC-MS (Method A): m/z: 429 (M+23); Rt = 3.89 min.

15 .

- Example 481 (General procedure (D))
6-(2,4-Dioxothiazolidin-5-ylidenemethyl)naphthalen-2-yloxyoctanoic acid



- 20 HPLC-MS (Method C): m/z: 436 (M+23); Rt.= 4.36 min


- The intermediate aldehyde for this compound was prepared by a slightly modified procedure:
6-Hydroxynaphthalene-2-carbaldehyde (1.0 g, 5.8 mmol) was dissolved in DMF (10 mL) and
sodium hydride 60% (278 mg) was added and the mixture stirred at RT for 15 min. 8-
25 Bromooctanoic acid (0.37 g, 1.7 mmol) was converted to the sodium salt by addition of
sodium hydride 60% and added to an aliquot (2.5 mL) of the above naphtholate solution and
the resulting mixture was stirred at RT for 16 hours. Aqueous acetic acid (10 %) was added
and the mixture was extracted 3 times with diethyl ether. The combined organic phases were

dried with MgSO_4 and evaporated to dryness affording 300 mg of 8-(6-formylnaphthalen-2-yloxy)octanoic acid.

HPLC-MS (Method C): m/z 315 ($M+1$); R_t = 4.24 min.

5 Example 482 (General procedure (D))

12-[6-(2,4-Dioxothiazolidin-5-ylidenemethyl)naphthalen-2-yloxy]dodecanoic acid.




HPLC-MS (Method C): m/z : 492 ($M+23$); R_t = 5.3 min.

10 The intermediate aldehyde was prepared similarly as described in example 481.

Example 483 (General procedure (D))

11-[6-(2,4-Dioxothiazolidin-5-ylidenemethyl)naphthalen-2-yloxy]undecanoic acid.




15

HPLC-MS (Method C): m/z : 478 ($M+23$); R_t = 5.17 min.

The intermediate aldehyde was prepared similarly as described in example 481.

20 Example 484 (General procedure (D))

15-[6-(2,4-Dioxothiazolidin-5-ylidenemethyl)naphthalen-2-yloxy]pentadecanoic acid.




HPLC-MS (Method C): m/z : 534 ($M+23$); R_t = 6.07 min.

25 The intermediate aldehyde was prepared similarly as described in example 481.

Example 485 (General procedure (D))


6-[6-(2,4-Dioxothiazolidin-5-ylidenemethyl)naphthalen-2-yloxy]hexanoic acid.



HPLC-MS (Method C): m/z : 408 ($M+23$); R_t . = 3.71 min.

Example 486 (General procedure (D))


- 5 4-[6-(2,4-Dioxothiazolidin-5-ylidenemethyl)naphthalen-2-yloxy]butyric acid.



HPLC-MS (Method C): m/z : 380 ($M+23$); R_t . = 3.23 min.

Example 487 (General procedure (D))


- 10 6-[6-(2,4-Dioxothiazolidin-5-ylidenemethyl)naphthalen-2-yloxy]hexanoic acid ethyl ester.



HPLC-MS (Method C): m/z : 436 ($M+23$); R_t . = 4.64 min.

Example 488 (General procedure (D))


- 15 4-[6-(2,4-Dioxothiazolidin-5-ylidenemethyl)naphthalen-2-yloxy]butyric acid ethyl ester.



HPLC-MS (Method C): m/z : 408 ($M+23$); R_t . = 4.28 min.

Example 489 (General procedure (D))


- 20 2-[5-[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)naphthalen-1-yloxy]pentyl]malonic acid



HPLC-MS (Method C): $m/z = 444$ ($M+1$); $R_t = 3,84$ min.

Example 490 (General procedure (D))


- 5 2-{5-[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)naphthalen-1-yloxy]pentyl}malonic acid diethyl ester



HPLC-MS (Method C): $m/z = 500$ ($M+1$); $R_t = 5.18$ min. .

- 10 **Example 491 (General procedure (D))**


4-[4-(2,4,6-Trioxotetrahydropyrimidin-5-ylidenemethyl)naphthalen-1-yloxy]butyric acid



HPLC-MS (Method C): $m/z = 369$ ($M+1$); $R_t = 2,68$ min.

- 15 **Example 492**

N-(3-Aminopropyl)-4-[4-(2,4-dioxothiazolidin-5-ylidenemethyl)-naphthalen-1-yloxy]-butyramide



To a mixture of 4-[4-(2,4-dioxothiazolidin-5-ylidenemethyl)naphthalen-1-yloxy]butyric acid (example 469, 5.9 g, 16.5 mmol) and 1-hydroxybenzotriazole (3.35 g, 24.8 mmol) in DMF (60 mL) was added 1-ethyl-3-(3'-dimethylaminopropyl)carbodiimide hydrochloride (4.75 g, 24.8 mmol) and the resulting mixture was stirred at room temperature for 2 hours. *N*-(3-amino-propylcarbamic acid *tert*-butyl ester (3.45 g, 19.8 mmol) was added and the resulting mixture was stirred at room temperature for 16 hours. The mixture was concentrated *in vacuo* and ethyl acetate and dichloromethane were added to the residue. The mixture was filtered, washed with water and dried *in vacuo* to afford 4.98 g (59%) of (3-{4-[4-(2,4-dioxothiazolidin-5-ylidenemethyl)naphthalen-1-yloxy]butyrylamino}propyl)carbamic acid *tert*-butyl ester.


HPLC-MS (Method C): *m/z*: 515 (*M*+1); *R*_t = 3.79 min.

(3-{4-[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)naphthalen-1-yloxy]butyrylamino}-propyl)carbamic acid *tert*-butyl ester (4.9 g, 9.5 mmol) was added dichloromethane (50 mL) and trifluoroacetic acid (50 mL) and the resulting mixture was stirred at room temperature for 45 minutes. The mixture was concentrated *in vacuo* and co-evaporated with toluene. To the residue was added ethyl acetate (100 mL) and the mixture was filtered and dried *in vacuo* to afford the title compound as the trifluoroacetic acid salt.


HPLC-MS (Method C): *m/z*: 414 (*M*+1); *R*_t = 2.27 min.

Compounds of the invention includes:


Example 493




Example 494




Example 495




Example 496




Example 497




Example 498




Example 499




Example 500




Example 501



Example 502




Example 503



Example 504 (Prepared analogously to General Procedure (D))

5 2-{5-[4-(2,4-Thiazolidindion-5-ylidenemethyl)naphthalen-1-yloxy]pentyl}malonic acid



A solution of 4-hydroxy-1-naphtaldehyde (1.0 g, 5.81 mmol), 2-(5-bromopentyl)malonic acid diethyl ester (2.07 g, 6.68 mmol) and potassium carbonate (4.01 g, 29 mmol) in DMF (50 mL) was stirred at 100° C for 3 hours. The mixture was cooled and the salt was filtered off. The solvent was then removed under reduced pressure to afford 2.9 g of crude 2-[5-(4-formylnaphtalen-1-yloxy)pentyl]malonic acid diethyl ester which was used for the next reaction without further purification.

HPLC-MS (Method C): m/z: 401 (M+1); Rt = 5.16 min. ¹H-NMR (DMSO-*d*₆): δ = 1.18 (t, 6 H), 1.39 (m, 2 H), 1.55 (m, 2 H), 1.87 (m, 4 H), 3.48 (t, 1 H), 4.13 (m, 4 H), 4.27 (t, 2 H), 7.17 (d, 1 H), 7.64 (t, 1 H), 7.75 (t, 1 H), 8.13 (d, 1 H), 8.29 (d, 1 H), 9.24 (d, 1 H), 10.19 (s, 1 H).

1.4 g (3.5 mmol) of crude 2-[5-(4-formylnaphtalen-1-yloxy)pentyl]malonic acid diethyl ester was treated with aqueous sodium hydroxide (1N, 8.75 mL, 8.75 mmol) and methanol (50 mL). The solution was stirred at 70° C for 5 hours and the mixture was concentrated under reduced pressure. Hydrochloric acid (6 N) was added until pH <2. The resulting slurry was stirred until it solidified. The crystals were filtered off, washed with water and then dried *in vacuo* to afford 1.1 g (92%) of 2-[5-(4-formylnaphtalen-1-yloxy)pentyl]malonic acid. The product was used in the next step without further purification.

HPLC-MS (Method C): m/z: 345 (M+1); Rt = 3.52 min. ¹H-NMR (DMSO-*d*₆): δ = 1.40 (m, 2 H), 1.55 (m, 2 H), 1.80 (m, 2 H), 1.90 (m, 2 H), 3.24 (t, 1 H), 4.29 (t, 2 H), 7.19 (d, 1 H), 7.64 (t, 1 H), 7.75 (t, 1 H), 8.14 (d, 1 H), 8.30 (d, 1 H), 9.23 (d, 1 H), 10.18 (s, 1 H), 12.69 (s, 2 H).

To a solution of 2-[5-(4-formylnaphtalen-1-yloxy)pentyl]malonic acid (0.36 g, 1.05 mmol) in acetic acid (10 mL) was added 2,4-thiazolidindione (0.16 g, 1.36 mmol) and piperidine (0.52 mL, 5.25 mmol). The solution was heated to 105 °C for 24 hours. After cooling to room temperature, the solvents were removed *in vacuo*. Water was added to the residue. The


precipitate was filtered off and washed with water. Recrystallisation from acetonitrile afforded 200 mg (43%) of the title compound as a solid.

HPLC-MS (Method C): m/z : 422 ($M-CO_2+Na$); R_t = 4.08 min. 1H -NMR($DMSO-d_6$): δ = 1,41


- 5 (m, 2 H), 1.55 (m, 4 H), 1.88 (m, 2 H), 2.23 (t, 1 H), 4.24 (t, 2 H), 7.61-7.74 (m, 3 H), 8.12 (d, 1 H), 8.28 (d, 1 H), 8.38 (s, 1 H), 12.00 (s, 1 H), 12.59 (s, 2 H).

- 10 The following compounds are commercially available and may be prepared according to general procedure (D):


Example 505




- 15 Example 506




Example 507



Example 508




Example 509




5

Example 510



10 Example 511




15

The following salicylic acid derivatives do all bind to the His B10 Zn²⁺ site of the insulin hexamer:


Example 512

Salicylic acid




5 Example 513

Thiosalicylic acid (or: 2-Mercaptobenzoic acid)




Example 514

10 2-Hydroxy-5-nitrobenzoic acid



Example 515


3-Nitrosalicylic acid



15


Example 516

5,5'-Methylenedisalicylic acid



Example 517


2-Amino-5-trifluoromethylbenzoic acid



5


Example 518

2-Amino-4-chlorobenzoic acid




10 Example 519

2-Amino-5-methoxybenzoic acid




Example 520

15




Example 521




20


Example 522




Example 523



Example 524




10 Example 525




Example 526

5-Iodosalicylic acid




Example 527

5-Chlorosalicylic acid



Example 528


1-Hydroxy-2-naphthoic acid



5

Example 529


3,5-Dihydroxy-2-naphthoic acid



10

Example 530


3-Hydroxy-2-naphthoic acid



15


Example 531

3,7-Dihydroxy-2-naphthoic acid




Example 532

20 2-Hydroxybenzo[a]carbazole-3-carboxylic acid



Example 533


7-Bromo-3-hydroxy-2-naphthoic acid



This compound was prepared according to Murphy *et al.*, *J. Med. Chem.* **1990**, 33, 171-8.
HPLC-MS (Method A): m/z: 267 (M+1); Rt: = 3.78 min.

Example 534


10 1,6-Dibromo-2-hydroxynaphthalene-3-carboxylic acid



This compound was prepared according to Murphy *et al.*, *J. Med. Chem.* **1990**, 33, 171-8.
HPLC-MS (Method A): m/z: 346 (M+1); Rt: = 4,19 min.

15 Example 535

7-Formyl-3-hydroxynaphthalene-2-carboxylic Acid



A solution of 7-bromo-3-hydroxynaphthalene-2-carboxylic acid (15.0 g, 56.2 mmol) (example 533) in tetrahydrofuran (100 mL) was added to a solution of lithium hydride (893 mg, 112 mmol) in tetrahydrofuran (350 mL). After 30 minutes stirring at room temperature, the resulting solution was heated to 50 °C for 2 minutes and then allowed to cool to ambient temperature over a period of 30 minutes. The mixture was cooled to -78 °C, and butyllithium (1.6 M in hexanes, 53 mL, 85 mmol) was added over a period of 15 minutes. *N,N*-

20


Dimethylformamide (8.7 mL, 8.2 g, 112 mmol) was added after 90 minutes additional stirring. The cooling was discontinued, and the reaction mixture was stirred at room temperature for 17 hours before it was poured into 1 N hydrochloric acid (aq.) (750 mL). The organic solvents were evaporated in vacuo, and the resulting precipitate was filtered off and rinsed with water (3 x 100 mL) to yield the crude product (16.2 g). Purification on silica gel (dichloromethane / methanol / acetic acid = 90:9:1) furnished the title compound as a solid.

¹H-NMR (DMSO-*d*₆): δ 11.95 (1H, bs), 10.02 (1H, s), 8.61 (1H, s), 8.54 (1H, s), 7.80 (2H, bs), 7.24 (1H, s); HPLC-MS (Method (A)): m/z: 217 (M+1); Rt = 2.49 min.

10


Example 536

3-Hydroxy-7-methoxy-2-naphthoic acid




15 Example 537

4-Amino-2-hydroxybenzoic acid




Example 538

20 5-Acetylamino-2-hydroxybenzoic acid



Example 539


2-Hydroxy-5-methoxybenzoic acid



The following compounds were prepared as described below:

Example 540

5 4-Bromo-3-hydroxynaphthalene-2-carboxylic acid




3-Hydroxynaphthalene-2-carboxylic acid (3.0 g, 15.9 mmol) was suspended in acetic acid (40 mL) and with vigorous stirring a solution of bromine (817 μ L, 15.9 mmol) in acetic acid (10 mL) was added drop wise during 30 minutes. The suspension was stirred at room
10 temperature for 1 hour, filtered and washed with water. Drying in vacuo afforded 3.74 g (88%) of 4-bromo-3-hydroxynaphthalene-2-carboxylic acid as a solid.

$^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 7.49 (1H, t), 7.75 (1H, t), 8.07 (2H, "t"), 8.64 (1H, s). The substitution pattern was confirmed by a COSY experiment, showing connectivities between the 3 (4
hydrogen) "triplets". HPLC-MS (Method A): m/z : 267 ($M+1$); R_t = 3.73 min.

15

Example 541

3-Hydroxy-4-iodonaphthalene-2-carboxylic acid




3-Hydroxynaphthalene-2-carboxylic acid (0.5 g, 2.7 mmol) was suspended in acetic acid (5
20 mL) and with stirring iodine monochloride (135 μ L, 2.7 mmol) was added. The suspension was stirred at room temperature for 1 hour, filtered and washed with water. Drying afforded 0.72 g (85%) of 4-iodo-3-hydroxynaphthalene-2-carboxylic acid as a solid.

$^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 7.47 (1H, t), 7.73 (1H, t), 7.98 (1H, d), 8.05 (1H, d), 8.66 (1H, s).

HPLC-MS (Method A): m/z : 315 ($M+1$); R_t = 3.94 min.

Example 542

2-Hydroxy-5-[(4-methoxyphenylamino)methyl]benzoic acid



5

p-Anisidine (1.3 g, 10.6 mmol) was dissolved in methanol (20 mL) and 5-formylsalicylic acid (1.75 g, 10.6 mmol) was added and the resulting mixture was stirred at room temperature for 16 hours. The solid formed was isolated by filtration, re-dissolved in N-methyl pyrrolidone (20 mL) and methanol (2 mL). To the mixture was added sodium cyanoborohydride (1.2 g) and the mixture was heated to 70 °C for 3 hours. To the cooled mixture was added ethyl acetate (100 mL) and the mixture was extracted with water (100 mL) and saturated aqueous ammonium chloride (100 mL). The combined aqueous phases were concentrated *in vacuo* and a 2 g aliquot was purified by SepPac chromatography eluting with mixtures of acetonitrile and water containing 0.1% trifluoroacetic acid to afford the title compound.


15

HPLC-MS (Method A): m/z: 274 (M+1); Rt = 1.77 min.

¹H-NMR (methanol-*d*₄): δ 3.82 (3H, s), 4.45 (2H, s), 6.96 (1H, d), 7.03 (2H, d), 7.23 (2H, d), 7.45 (1H, dd), 7.92 (1H, d).

20 Example 543

2-Hydroxy-5-(4-methoxyphenylsulfamoyl)benzoic acid



25


A solution of 5-chlorosulfonylsalicylic acid (0.96 g, 4.1 mmol) in dichloromethane (20 mL) and triethylamine (1.69 mL, 12.2 mmol) was added p-anisidine (0.49 g, 4.1 mmol) and the resulting mixture was stirred at room temperature for 16 hours. The mixture was added dichloromethane (50 mL) and was washed with water (2 x 100 mL). Drying (MgSO₄) of the organic phase and concentration *in vacuo* afforded 0.57 g crude product. Purification by

column chromatography on silica gel eluting first with ethyl acetate:heptane (1:1) then with methanol afforded 0.1 g of *the title compound*.

HPLC-MS (Method A): m/z: 346 (M+23); Rt = 2.89 min.

- 5 ¹H-NMR (DMSO-*d*₆): δ 3.67 (3H, s), 6.62 (1H, d), 6.77 (2H, d), 6.96 (2H, d), 7.40 (1H, dd), 8.05 (1H, d), 9.6 (1H, bs).

General procedure (E) for preparation of compounds of general formula I₄:




wherein Lea is a leaving group such as Cl, Br, I or OSO₂CF₃, R is hydrogen or C₁-C₆-alkyl, optionally the two R-groups may together form a 5-8 membered ring, a cyclic boronic acid ester, and J is as defined above.

15 An analogous chemical transformation has previously been described in the literature (Bumagin et al., *Tetrahedron*, **1997**, 53, 14437-14450). The reaction is generally known as the Suzuki coupling reaction and is generally performed by reacting an aryl halide or triflate with an arylboronic acid or a heteroarylboronic acid in the presence of a palladium catalyst
20 and a base such as sodium acetate, sodium carbonate or sodium hydroxide. The solvent can be water, acetone, DMF, NMP, HMPA, methanol, ethanol toluene or a mixture of two or more of these solvents. The reaction is performed at room temperature or at elevated temperature.

The general procedure (E) is further illustrated in the following example:

25 **Example 544 (General Procedure (E))**

7-(4-Acetylphenyl)-3-hydroxynaphthalene-2-carboxylic Acid




To 7-bromo-3-hydroxynaphthalene-2-carboxylic acid (100 mg, 0.37 mmol) (example 533) was added a solution of 4-acetylphenylboronic acid (92 mg, 0.56 mmol) in acetone (2.2 mL) followed by a solution of sodium carbonate (198 mg, 1.87 mmol) in water (3.3 mL). A suspension of palladium(II) acetate (4 mg, 0.02 mmol) in acetone (0.5 mL) was filtered and added to the above solution. The mixture was purged with N₂ and stirred vigorously for 24 hours at room temperature. The reaction mixture was poured into 1 N hydrochloric acid (aq.) (60 mL) and the precipitate was filtered off and rinsed with water (3 x 40 mL). The crude product was dissolved in acetone (25 mL) and dried with magnesium sulfate (1 h). Filtration followed by concentration furnished the title compound as a solid (92 mg).

¹H-NMR (DMSO-*d*₆): δ 12.60 (1H, bs), 8.64 (1H, s), 8.42 (1H, s), 8.08 (2H, d), 7.97 (2H, d), 7.92 (2H, m), 7.33 (1H, s), 2.63 (3H, s); HPLC-MS (Method (A)): *m/z*: 307 (M+1); *R*_t = 3.84 min.

The compounds in the following examples were prepared in a similar fashion. Optionally, the compounds can be further purified by recrystallization from e.g. ethanol or by chromatography.

Example 545 (General Procedure (E))


3-Hydroxy-7-(3-methoxyphenyl)naphthalene-2-carboxylic acid



HPLC-MS (Method (A)): *m/z*: 295 (M+1); *R*_t = 4.60 min.

Example 546 (General Procedure (E))


3-Hydroxy-7-phenylnaphthalene-2-carboxylic acid



HPLC-MS (Method (A)): *m/z*: 265 (M+1); *R*_t = 4.6 min.

Example 547 (General Procedure (E))


3-Hydroxy-7-p-tolynaphthalene-2-carboxylic acid



HPLC-MS (Method (A)): m/z: 279 (M+1); Rt = 4.95 min.

Example 548 (General Procedure (E))


5 7-(4-Formylphenyl)-3-hydroxynaphthalene-2-carboxylic acid



HPLC-MS (Method (A)): m/z: 293 (M+1); Rt = 4.4 min.

Example 549 (General Procedure (E))


10 6-Hydroxy-[1,2]binaphthalenyl-7-carboxylic acid



HPLC-MS (Method (A)): m/z: 315 (M+1); Rt = 5.17 min.

Example 550 (General Procedure (E))


15 7-(4-Carboxy-phenyl)-3-hydroxynaphthalene-2-carboxylic acid



HPLC-MS (Method (A)): m/z: 309 (M+1); Rt = 3.60 min.

Example 551 (General Procedure (E))


20 7-Benzofuran-2-yl-3-hydroxynaphthalene-2-carboxylic acid



HPLC-MS (Method (A)): m/z: 305 (M+1); Rt = 4.97 min.

Example 552 (General Procedure (E))


5 3-Hydroxy-7-(4-methoxyphenyl)-naphthalene-2-carboxylic acid



HPLC-MS (Method (A)): m/z: 295 (M+1); Rt = 4.68 min.

Example 553 (General Procedure (E))


10 7-(3-Ethoxyphenyl)-3-hydroxynaphthalene-2-carboxylic acid



HPLC-MS (Method (A)): m/z: 309 (M+1); Rt = 4.89 min.

Example 554 (General Procedure (E))


15 7-Benzo[1,3]dioxol-5-yl-3-hydroxynaphthalene-2-carboxylic acid



HPLC-MS (Method (A)): m/z: 309 (M+1); Rt = 5.61 min.


Example 555 (General Procedure (E))

20 7-Biphenyl-3-yl-3-hydroxynaphthalene-2-carboxylic acid



HPLC-MS (Method (A)): m/z: 341 (M+1); Rt = 5.45 min.

General procedure (F) for preparation of compounds of general formula I₅:




wherein R³⁰ is hydrogen or C₁-C₆-alkyl and T is as defined above

This general procedure (F) is further illustrated in the following example:

Example 556 (General procedure (F))

3-Hydroxy-7-[(4-(2-propyl)phenylamino)methyl]naphthalene-2-carboxylic Acid




7-Formyl-3-hydroxynaphthalene-2-carboxylic acid (40 mg, 0.19 mmol) (example 535) was suspended in methanol (300 μ L). Acetic acid (16 μ L, 17 mg, 0.28 mmol) and 4-(2-propyl)aniline (40 μ L, 40 mg, 0.30 mmol) were added consecutively, and the resulting mixture was stirred vigorously at room temperature for 2 hours. Sodium cyanoborohydride (1.0 M in tetrahydrofuran, 300 μ L, 0.3 mmol) was added, and the stirring was continued for another 17 hours. The reaction mixture was poured into 6 N hydrochloric acid (aq.) (6 mL), and the precipitate was filtered off and rinsed with water (3 x 2 mL) to yield the title compound (40 mg) as its hydrochloride salt. No further purification was necessary.

¹H-NMR (DMSO-*d*₆): δ 10.95 (1H, bs), 8.45 (1H, s), 7.96 (1H, s), 7.78 (1H, d), 7.62 (1H, d), 7.32 (1H, s), 7.13 (2H, bd), 6.98 (2H, bd), 4.48 (2H, s), 2.79 (1H, sept), 1.14 (6H, d); HPLC-MS (Method (A)): m/z: 336 (M+1); Rt = 3.92 min.

The compounds in the following examples were made using this general procedure (F).

Example 557 (General procedure (F))


7-[[[4-Bromophenyl]amino]methyl]-3-hydroxynaphthalene-2-carboxylic Acid



HPLC-MS (Method C): m/z: 372 (M+1); Rt = 4.31min.

Example 558 (General procedure (F))


7-[[[3,5-Dichlorophenyl]amino]methyl]-3-hydroxynaphthalene-2-carboxylic Acid



HPLC-MS (Method C): m/z: 362 (M+1); Rt = 4.75 min.

Example 559 (General procedure (F))


7-[[[Benzothiazol-6-yl]amino]methyl]-3-hydroxynaphthalene-2-carboxylic Acid



HPLC-MS (Method C): m/z: 351 (M+1); Rt = 3.43 min.

Example 560 (General procedure (F))


3-Hydroxy-7-[[[quinolin-6-yl]amino]methyl]naphthalene-2-carboxylic Acid



HPLC-MS (Method C): m/z: 345 (M+1); Rt = 2.26 min.

Example 561 (General procedure (F))


3-Hydroxy-7-[[[4-methoxyphenyl]amino]methyl]naphthalene-2-carboxylic Acid



HPLC-MS (Method C): m/z : 324 ($M+1$); R_t = 2.57 min.

Example 562 (General procedure (F))


- 5 7-[[[(2,3-Dihydrobenzofuran-5-ylmethyl)amino]methyl]-3-hydroxynaphthalene-2-carboxylic Acid



HPLC-MS (Method C): m/z : 350 ($M+1$); R_t = 2.22 min.

- 10 Example 563 (General procedure (F))


7-[[[(4-Chlorobenzyl)amino]methyl]-3-hydroxynaphthalene-2-carboxylic Acid



HPLC-MS (Method C): m/z : 342 ($M+1$); R_t = 2.45 min.

- 15 Example 564 (General procedure (F))


3-Hydroxy-7-[[[(naphthalen-1-ylmethyl)amino]methyl]naphthalene-2-carboxylic Acid



HPLC-MS (Method C): m/z : 357 ($M+1$); R_t = 2.63 min.

- 20 Example 565 (General procedure (F))


7-[[[(Biphenyl-2-ylmethyl)amino]methyl]-3-hydroxynaphthalene-2-carboxylic Acid



HPLC-MS (Method C): m/z : 384 (M+1); R_t = 2.90 min.

Example 566 (General procedure (F))


3-Hydroxy-7-[[[4-phenoxybenzyl]amino]methyl]naphthalene-2-carboxylic Acid



HPLC-MS (Method C): m/z : 400 (M+1); R_t = 3.15 min.


Example 567 (General procedure (F))

3-Hydroxy-7-[[[4-methoxybenzyl]amino]methyl]naphthalene-2-carboxylic Acid



HPLC-MS (Method C): m/z : 338 (M+1); R_t = 2.32 min.

General procedure (G) for preparation of compounds of general formula I₆:




wherein J is as defined above and the moiety $(C_1-C_6\text{-alkanoyl})_2O$ is an anhydride.

The general procedure (G) is illustrated by the following example:

20 Example 568 (General procedure (G))

N-Acetyl-3-hydroxy-7-[(4-(2-propyl)phenylamino)methyl]naphthalene-2-carboxylic Acid




3-Hydroxy-7-[(4-(2-propyl)phenylamino)methyl]naphthalene-2-carboxylic acid (25 mg, 0.07 mmol) (example 556) was suspended in tetrahydrofuran (200 μ L). A solution of sodium hydrogencarbonate (23 mg, 0.27 mmol) in water (200 μ L) was added followed by acetic anhydride (14 μ L, 15 mg, 0.15 mmol). The reaction mixture was stirred vigorously for 65 hours at room temperature before 6 N hydrochloric acid (4 mL) was added. The precipitate was filtered off and rinsed with water (3 x 1 mL) to yield the title compound (21 mg). No further purification was necessary.

¹H-NMR (DMSO-*d*₆): δ 10.96 (1H, bs), 8.48 (1H, s), 7.73 (1H, s), 7.72 (1H, d), 7.41 (1H, dd), 7.28 (1H, s), 7.23 (2H, d), 7.18 (2H, d), 4.96 (2H, s), 2.85 (1H, sept), 1.86 (3H, s), 1.15 (6H, d); HPLC-MS (Method (A)): *m/z*: 378 (M+1); *R*_t = 3.90 min.

The compounds in the following examples were prepared in a similar fashion.

15 Example 569 (General procedure (G))


N-Acetyl-7-[[[4-(bromophenyl)amino]methyl]-3-hydroxynaphthalene-2-carboxylic Acid



HPLC-MS (Method C): *m/z*: 414 (M+1); *R*_t = 3.76 min.

20 Example 570 (General procedure (G))

N-Acetyl-7-[[[2,3-dihydrobenzofuran-5-ylmethyl]amino]methyl]-3-hydroxynaphthalene-2-carboxylic Acid




HPLC-MS (Method C): *m/z*: 392 (M+1); *R*_t = 3.26 min.

25

Example 571 (General procedure (G))

N-Acetyl-7-[[[4-(chlorobenzyl)amino]methyl]-3-hydroxynaphthalene-2-carboxylic Acid




HPLC-MS (Method C): m/z : 384 ($M+1$); R_t = 3.67 min.

Compounds of the invention may also include tetrazoles:

Example 572

5 5-(3-(Naphthalen-2-yloxymethyl)-phenyl)-1*H*-tetrazole



To a mixture of 2-naphthol (10 g, 0.07 mol) and potassium carbonate (10 g, 0.073 mol) in acetone (150 mL), alpha-bromo-m-tolunitril (13.6 g, 0.07 mol) was added in portions. The reaction mixture was stirred at reflux temperature for 2.5 hours. The cooled reaction mixture
10 was filtered and evaporated in vacuo affording an oily residue (19 g) which was dissolved in diethyl ether (150 mL) and stirred with a mixture of active carbon and $MgSO_4$ for 16 hours. The mixture was filtered and evaporated in vacuo affording crude 18.0 g (100 %) of 3-(naphthalen-2-yloxymethyl)-benzonitrile as a solid.
12 g of the above benzonitrile was recrystallised from ethanol (150 mL) affording 8.3 g (69
15 %) of 3-(naphthalen-2-yloxymethyl)-benzonitrile as a solid.

M.p. 60 - 61 °C.

Calculated for $C_{18}H_{13}NO$:

C, 83.37 %; H, 5.05 %; N, 5.40 %; Found

20 C, 83.51 %; H, 5.03 %; N, 5.38 %.

To a mixture of sodium azide (1.46 g, 22.5 mmol) and ammonium chloride (1.28 g, 24.0 mmol) in dry dimethylformamide (20 mL) under an atmosphere of nitrogen, 3-(naphthalen-2-yloxymethyl)-benzonitrile (3.9 g, 15 mmol) was added and the reaction mixture was stirred at
25 125 °C for 4 hours. The cooled reaction mixture was poured on to ice water (300 mL) and acidified to pH = 1 with 1 N hydrochloric acid. The precipitate was filtered off and washed with water, dried at 100 °C for 4 hours affording 4.2 g (93 %) of the title compound.

M.p. 200 - 202 °C.

30 Calculated for $C_{18}H_{14}N_4O$:


C, 71.51 %; H, 4.67 %; N, 18.54 %; Found

C, 72.11 %; H, 4.65 %; N, 17.43 %.

¹H NMR (400 MHz, DMSO-d₆) δ_H 5.36 (s, 2H), 7.29 (dd, 1H), 7.36 (dt, 1H), 7.47 (m, 2H), 7.66 (t, 1H), 7.74 (d, 1H), 7.84 (m, 3H), 8.02 (d, 1H), 8.22 (s, 1H).

5 Example 573

N-(3-(Tetrazol-5-yl)phenyl)-2-naphtoic acid amide



2-Naphtoic acid (10 g, 58 mmol) was dissolved in dichloromethane (100 mL) and N,N-dimethylformamide (0.2 mL) was added followed by thionyl chloride (5.1 ml, 70 mmol). The mixture was heated at reflux temperature for 2 hours. After cooling to room temperature, the mixture was added dropwise to a mixture of 3-aminobenzonitril (6.90 g, 58 mmol) and triethylamine (10 mL) in dichloromethane (75 mL). The resulting mixture was stirred at room temperature for 30 minutes. Water (50 mL) was added and the volatiles was exaporated in vacuo. The resulting mixture was filtered and the filter cake was washed with water followed by heptane (2 x 25 mL). Drying in vacuo at 50 °C for 16 hours afforded 15.0 g (95 %) of N-(3-cyanophenyl)-2-naphtoic acid amide.

M.p. 138-140 °C

The above naphthoic acid amide (10 g, 37 mmol) was dissolved in N,N-dimethylformamide (200 mL) and sodium azide (2.63 g, 40 mmol) and ammonium chloride (2.16 g, 40 mmol) were added and the mixture heated at 125 °C for 6 hours. Sodium azide (1.2 g) and ammonium chloride (0.98 g) were added and the mixture heated at 125 °C for 16 hours. After cooling, the mixture was poured into water (1.5 l) and stirred at room temperature for 30 minutes. The solid formed was filtered off, washed with water and dried in vacuo at 50 °C for 3 days affording 9.69 g (84 %) of the title compound as a solid which could be further purified by treatment with ethanol at reflux temperature.

¹H NMR (200 MHz, DMSO-d₆): δ_H 7.58-7.70 (m, 3H), 7.77 (d, 1H), 8.04-8.13 (m, 5H), 8.65 (d, 1H), 10.7 (s, 1H).


Calculated for $C_{18}H_{13}N_5O$, 0.75 H_2O :

C, 65.74 %; H, 4.44 %; N, 21.30 %. Found:

C, 65.58 %; H, 4.50 %; N, 21.05 %.

5 Example 574

5-[3-(Biphenyl-4-yloxymethyl)phenyl]-1H-tetrazole



To a solution of 4-phenylphenol (10.0 g, 59 mmol) in dry N,N-dimethyl-formamide (45 mL) kept under an atmosphere of nitrogen, sodium hydride (2.82 g, 71 mmol, 60 % dispersion in oil) was added in portions and the reaction mixture was stirred until gas evolution ceased. A solution of m-cyanobenzyl bromide (13 g, 65 mmol) in dry N,N-dimethylformamide (45 mL) was added dropwise and the reaction mixture was stirred at room temperature for 18 hours. The reaction mixture was poured on to ice water (150 mL). The precipitate was filtered of and washed with 50 % ethanol (3 x 50 mL), ethanol (2 x 50 mL), diethyl ether (80 mL), and dried in vacuo at 50 °C for 18 hours affording crude 17.39 g of 3-(biphenyl-4-yloxymethyl)-benzonitrile as a solid.

1H NMR (200 MHz, $CDCl_3$) δ_H 5.14 (s, 2H), 7.05 (m, 2H), 7.30 - 7.78 (m, 11H).


20

To a mixture of sodium azide (2.96 g, 45.6 mmol) and ammonium chloride (2.44 g, 45.6 mmol) in dry N,N-dimethylformamide (100 mL) under an atmosphere of nitrogen, 3-(biphenyl-4-yloxymethyl)-benzonitrile (10.0 g, 35.0 mmol) was added and the reaction mixture was stirred at 125 °C for 18 hours. The cooled reaction mixture was poured on to a mixture of 1N hydrochloric acid (60 mL) and ice water (500 mL). The precipitate was filtered off and washed with water (3 x 100 mL), 50 % ethanol (3 x 100 mL), ethanol (50 mL), diethyl ether (50 mL), ethanol (80 mL), and dried in vacuo at 50 °C for 18 hours affording 8.02 g (70 %) of the title compound.

30 1H NMR (200 MHz, $DMSO-d_6$) δ_H 5.31 (s, 2H), 7.19 (m, 2H), 7.34 (m, 1H), 7.47 (m, 2H), 7.69 (m, 6H), 8.05 (dt, 1H), 8.24 (s, 1H).

Example 575

5-(3-Phenoxymethyl)-phenyl-tetrazole



- 5 3-Bromomethylbenzonitrile (5.00 g, 25.5 mmol) was dissolved in N,N-dimethylformamide (50 mL), phenol (2.40 g, 25.5 mmol) and potassium carbonate (10.6 g, 77 mmol) were added. The mixture was stirred at room temperature for 16 hours. The mixture was poured into water (400 mL) and extracted with ethyl acetate (2 x 200 mL). The combined organic extracts were washed with water (2 x 100 mL), dried (MgSO₄) and evaporated in vacuo to afford 5.19 g (97 %) 3-(phenoxymethyl)benzonitrile as an oil.
- 10


TLC: R_f = 0.38 (Ethyl acetate/heptane = 1:4)

- The above benzonitrile (5.19 g, 24.8 mmol) was dissolved in N,N-dimethylformamide (100 mL) and sodium azide (1.93 g, 30 mmol) and ammonium chloride (1.59 g, 30 mmol) were added and the mixture was heated at 140 °C for 16 hours. After cooling, the mixture was poured into water (800 mL). The aqueous mixture was washed with ethyl acetate (200 mL). The pH of the aqueous phase was adjusted to 1 with 5 N hydrochloric acid and stirred at room temperature for 30 minutes. Filtration, washing with water and drying in vacuo at 50 °C afforded 2.06 g (33 %) of the title compound as a solid.
- 15
- 20

¹H NMR (200 MHz, CDCl₃ + DMSO-*d*₆) δ_H 5.05 (s, 2H), 6.88 (m, 3H), 7.21 (m, 2H), 7.51 (m, 2H), 7.96 (dt, 1H), 8.14 (s, 1H).

Example 576

- 25 5-[3-(Biphenyl-4-ylmethoxy)phenyl]-1H-tetrazole



To a solution of 3-cyanophenol (5.0 g, 40.72 mmol) in dry N,N-dimethylformamide (100 mL) kept under an atmosphere of nitrogen, sodium hydride (2 g, 48.86 mmol, 60 % dispersion in oil) was added in portions and the reaction mixture was stirred until gas evolution ceased. p-


Phenylbenzyl chloride (9.26 g, 44.79 mmol) and potassium iodide (0.2 g, 1.21 mmol) were added and the reaction mixture was stirred at room temperature for 60 hours. The reaction mixture was poured on to a mixture of saturated sodium carbonate (100 mL) and ice water (300 mL). The precipitate was filtered off and washed with water (3 x 100 mL), n-hexane (2 x 80 mL) and dried in vacuo at 50 °C for 18 hours affording 11.34 g (98 %) of 3-(biphenyl-4-ylmethoxy)-benzonitrile as a solid.

To a mixture of sodium azide (2.37 g, 36.45 mmol) and ammonium chloride (1.95 g, 36.45 mmol) in dry N,N-dimethylformamide (100 mL) under an atmosphere of nitrogen, 3-(biphenyl-4-ylmethoxy)-benzonitrile (8.0 g, 28.04 mmol) was added and the reaction mixture was stirred at 125 °C for 18 hours. To the cooled reaction mixture water (100 mL) was added and the reaction mixture stirred for 0.75 hour. The precipitate was filtered off and washed with water, 96 % ethanol (2 x 50 mL), and dried in vacuo at 50°C for 18 hours affording 5.13 g (56 %) of the title compound.

¹H NMR (200 MHz, DMSO-d₆) δ_H 5.29 (s, 2H), 7.31 (dd, 1H), 7.37 - 7.77 (m, 12H).

Example 577


20 5-[4-(Biphenyl-4-ylmethoxy)-3-methoxyphenyl]-1H-tetrazol



This compound was made similarly as described in example 576.


Example 578

25



Example 579


5-(2-Naphthylmethyl)-1H-tetrazole



This compound was prepared similarly as described in example 572, step 2.

Example 580


5 5-(1-Naphthylmethyl)-1H-tetrazole



This compound was prepared similarly as described in example 572, step 2.

Example 581

10 5-[4-(Biphenyl-4-yloxymethyl)phenyl]-1H-tetrazole



A solution of alpha-bromo-p-tolunitrile (5.00 g, 25.5 mmol), 4-phenylphenol (4.56 g, 26.8 mmol), and potassium carbonate (10.6 g, 76.5 mmol) in N,N-dimethylformamide (75 mL) was stirred vigorously for 16 hours at room temperature. Water (75 mL) was added and the mixture was stirred at room temperature for 1 hour. The precipitate was filtered off and washed thoroughly with water. Drying in vacuo over night at 50 °C afforded 7.09 g (97 %) of 4-(biphenyl-4-yloxymethyl)benzonitrile as a solid.


The above benzonitrile (3.00 g, 10.5 mmol) was dissolved in N,N-dimethylformamide (50 mL), and sodium azide (1.03 g, 15.8 mmol) and ammonium chloride (0.84 g, 15.8 mmol) were added and the mixture was stirred 16 hours at 125 °C. The mixture was cooled to room temperature and water (50 mL) was added. The suspension was stirred overnight, filtered, washed with water and dried in vacuo at 50 °C for 3 days to give crude 3.07 g (89 %) of the title compound. From the mother liquor crystals were collected and washed with water, dried by suction to give 0.18 g (5 %) of the title compound as a solid.

^1H NMR (200 MHz, DMSO- d_6): δ_{H} 5.21 (s, 2H), 7.12 (d, 2H), 7.30 (t, 1H), 7.42 (t, 2H), 7.56-7.63 (m, 6H), 8.03 (d, 2H).

Calculated for $\text{C}_{20}\text{H}_{16}\text{N}_4\text{O}$, $2\text{H}_2\text{O}$:

5 C, 65.92 %; H, 5.53 %; N, 15.37 %. Found:
C, 65.65 %; H, 5.01 %; N, 14.92 %.


Example 582



10


This compound was prepared similarly as described in example 576.

Example 583




15

Example 584




20

Example 585



Example 586


5-(3-(Biphenyl-4-yloxymethyl)-benzyl)-1H-tetrazole



5

Example 587


5-(1-Naphthyl)-1H-tetrazole



10 This compound was prepared similarly as described in example 572, step 2.

Example 588


5-[3-Methoxy-4-(4-methylsulfonylbenzyloxy)phenyl]-1H-tetrazole



15 This compound was made similarly as described in example 576.

Example 589


5-(2-Naphthyl)-1H-tetrazole



This compound was prepared similarly as described in example 572, step 2.


Example 590

5 2-Amino-N-(1H-tetrazol-5-yl)-benzamide



Example 591

5-(4-Hydroxy-3-methoxyphenyl)-1H-tetrazole




10

This compound was prepared similarly as described in example 572, step 2.

Example 592

4-(2H-Tetrazol-5-ylmethoxy)benzoic acid



15

To a mixture of methyl 4-hydroxybenzoate (30.0 g, 0.20 mol), sodium iodide (30.0 g, 0.20 mol) and potassium carbonate (27.6 g, 0.20 mol) in acetone (2000 mL) was added chloroacetonitrile (14.9 g, 0.20 mol). The mixture was stirred at RT for 3 days. Water was added and the mixture was acidified with 1N hydrochloric acid and the mixture was extracted with diethyl ether. The combined organic layers were dried over Na₂SO₄ and concentrated *in vacuo*. The residue was dissolved in acetone and chloroacetonitrile (6.04 g, 0.08 mol),

sodium iodide (12.0 g, 0.08 mol) and potassium carbonate (11.1 g, 0.08 mol) were added and the mixture was stirred for 16 hours at RT and at 60 °C. More chloroacetonitrile was added until the conversion was 97%. Water was added and the mixture was acidified with 1N hydrochloric acid and the mixture was extracted with diethyl ether. The combined organic layers were dried over Na₂SO₄ and concentrated *in vacuo* to afford methyl 4-cyanomethoxybenzoate in quantitative yield. This compound was used without further purification in the following step.


A mixture of methyl 4-cyanomethoxybenzoate (53.5 g, 0.20 mol), sodium azide (16.9 g, 0.26 mol) and ammonium chloride (13.9 g, 0.26 mol) in DMF 1000 (mL) was refluxed overnight under N₂. After cooling, the mixture was concentrated *in vacuo*. The residue was suspended in cold water and extracted with ethyl acetate. The combined organic phases were washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*, to afford methyl 4-(2H-tetrazol-5-ylmethoxy)benzoate. This compound was used as such in the following step.

Methyl 4-(2H-Tetrazol-5-ylmethoxy)-benzoate was refluxed in 3N sodium hydroxide. The reaction was followed by TLC (DCM:MeOH = 9:1). The reaction mixture was cooled, acidified and the product filtered off. The impure product was washed with DCM, dissolved in MeOH, filtered and purified by column chromatography on silica gel (DCM:MeOH = 9:1). The resulting product was recrystallised from DCM:MeOH=95:5. This was repeated until the product was pure. This afforded 13.82 g (30 %) of the title compound.

¹H-NMR (DMSO-*d*₆): 4.70 (2H, s), 7.48 (2H, d), 7.73 (2H, d), 13 (1H, bs).

Example 593

4-(2H-Tetrazol-5-ylmethylsulfanyl)benzoic acid




To a solution of sodium hydroxide (10.4 g, 0.26 mol) in degassed water (600 mL) was added 4-mercaptobenzoic acid (20.0 g, 0.13 mol). This solution was stirred for 30 minutes. To a solution of potassium carbonate (9.0 g, 65 mmol) in degassed water (400 mL) was added chloroacetonitrile (9.8 g, (0.13 mol) portion-wise. These two solutions were mixed and stirred for 48 hours at RT under N₂. The mixture was filtered and washed with heptane. The

aqueous phase was acidified with 3N hydrochloric acid and the product was filtered off, washed with water and dried, affording 4-cyanomethylsulfanylbenzoic acid (27.2 g, 88%). This compound was used without further purification in the following step.

- 5 A mixture of 4-cyanomethylsulfanylbenzoic acid (27.2 g, 0.14 mol), sodium azide (11.8 g, 0.18 mol) and ammonium chloride (9.7 g, 0.18 mol) in DMF (1000 mL) was refluxed overnight under N₂. The mixture was concentrated *in vacuo*. The residue was suspended in cold water and extracted with diethyl ether. The combined organic phases were washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*. Water was added and the precipitate
10 was filtered off. The aqueous layer was concentrated *in vacuo*, water was added and the precipitate filtered off. The combined impure products were purified by column chromatography using DCM:MeOH = 9:1 as eluent, affording the title compound (5.2 g, 16%).
- 15 ¹H-NMR (DMSO-*d*₆): 5.58 (2H, s), 7.15 (2H, d), 7.93 (2H, d), 12.7 (1H, bs).

Example 594

3-(2*H*-Tetrazol-5-yl)-9*H*-carbazole



- 20 3-Bromo-9*H*-carbazole was prepared as described by Smith *et al.* in *Tetrahedron* **1992**, *48*, 7479-7488.
- A solution of 3-bromo-9*H*-carbazole (23.08 g, 0.094 mol) and cuprous cyanide (9.33 g, 0.103 mol) in *N*-methyl-pyrrolidone (300 ml) was heated at 200 °C for 5 h. The cooled reaction mixture was poured on to water (600 ml) and the precipitate was filtered off and washed with
25 ethyl acetate (3 x 50 ml). The filtrate was extracted with ethyl acetate (3 x 250 ml) and the combined ethyl acetate extracts were washed with water (150 ml), brine (150 ml), dried (MgSO₄) and concentrated *in vacuo*. The residue was crystallised from heptanes and recrystallised from acetonitrile (70 ml) affording 7.16 g (40 %) of 3-cyano-9*H*-carbazole as a solid. M.p. 180 - 181 °C.


3-Cyano-9H-carbazole (5.77 g, 30 mmol) was dissolved in *N,N*-dimethylformamide (150 ml), and sodium azide (9.85 g, 152 mmol), ammonium chloride (8.04 g, 150 mmol) and lithium chloride (1.93 g, 46 mmol) were added and the mixture was stirred for 20 h at 125 °C. To the reaction mixture was added an additional portion of sodium azide (9.85 g, 152 mmol) and ammonium chloride (8.04 g, 150 mmol) and the reaction mixture was stirred for an additional 24 h at 125 °C. The cooled reaction mixture was poured on to water (500 ml). The suspension was stirred for 0.5 h, and the precipitate was filtered off and washed with water (3 x 200 ml) and dried *in vacuo* at 50 °C. The dried crude product was suspended in diethyl ether (500 ml) and stirred for 2 h, filtered off and washed with diethyl ether (2 x 200 ml) and dried *in vacuo* at 50 °C affording 5.79 g (82 %) of the title compound as a solid.

¹H-NMR (DMSO-*d*₆): δ 11.78 (1H, bs), 8.93 (1H, d), 8.23 (1H, d), 8.14 (1H, dd), 7.72 (1H, d), 7.60 (1H, d), 7.49 (1H, t), 7.28 (1H, t); HPLC-MS (Method C): *m/z*: 236 (M+1); *R*_t = 2.77 min.

- 15 The following commercially available tetrazoles do all bind to the His B10 Zn²⁺ site of the insulin hexamer:

Example 595


5-(3-Tolyl)-1*H*-tetrazole



20

Example 596


5-(2-Bromophenyl)tetrazole



25


Example 597

5-(4-Ethoxalylamino-3-nitrophenyl)tetrazole




Example 598

5




Example 599


10



Example 600




15 Example 601



Example 602


Tetrazole



5

Example 603


5-Methyltetrazole



10

Example 604


5-Benzyl-2H-tetrazole



15


Example 605

4-(2H-Tetrazol-5-yl)benzoic acid




Example 606

20 5-Phenyl-2H-tetrazole



Example 607


5-(4-Chlorophenylsulfanylmethyl)-2H-tetrazole



5


Example 608

5-(3-Benzyloxyphenyl)-2H-tetrazole




10 Example 609

2-Phenyl-6-(1H-tetrazol-5-yl)-chromen-4-one




Example 610


15



Example 611




Example 612




5

Example 613



10


Example 614




15

Example 615


5-(4-Bromo-phenyl)-1H-tetrazole




Example 616




Example 617



Example 618




15 Example 619




Example 620


20




Example 621




Example 622




Example 623




15 Example 624




Example 625




Example 626




Example 627




15 Example 628



Example 629




Example 630




5

Example 631




10


Example 632



15 Example 633




Example 634




5

Example 635




10


Example 636



15 Example 637




Example 638




5

Example 639




10


Example 640



15 Example 641




Example 642




5

Example 643




10

Example 644




Example 645




5 Example 646

5-(2,6-Dichlorobenzyl)-2H-tetrazole



General procedure (H) for preparation of compounds of general formula I₇:



wherein K, M, and T are as defined above.


The reaction is generally known as a reductive alkylation reaction and is generally performed by stirring an aldehyde with an amine at low pH (by addition of an acid, such as acetic acid or formic acid) in a solvent such as THF, DMF, NMP, methanol, ethanol, DMSO, dichloromethane, 1,2-dichloroethane, trimethyl orthoformate, triethyl orthoformate, or a mixture of two or more of these. As reducing agent sodium cyano borohydride or sodium triacetoxy borohydride may be used. The reaction is performed between 20°C and 120°C, preferably at room temperature.

When the reductive alkylation is complete, the product is isolated by extraction, filtration, chromatography or other methods known to those skilled in the art.

The general procedure (H) is further illustrated in the following example 647:

Example 647 (General procedure (H))


Biphenyl-4-ylmethyl-[3-(2H-tetrazol-5-yl)phenyl]amine



- 5 A solution of 5-(3-aminophenyl)-2H-tetrazole (example 874, 48 mg, 0.3 mmol) in DMF (250 μ L) was mixed with a solution of 4-biphenylaldehyde (54 mg, 0.3 mmol) in DMF (250 μ L) and acetic acid glacial (250 μ L) was added to the mixture followed by a solution of sodium cyano borohydride (15 mg, 0.24 mmol) in methanol (250 μ L). The resulting mixture was shaken at room temperature for 2 hours. Water (2 mL) was added to the mixture and the resulting mixture was shaken at room temperature for 16 hours. The mixture was
- 10 centrifugated (6000 rpm, 10 minutes) and the supernatant was removed by a pipette. The residue was washed with water (3 mL), centrifugated (6000 rpm, 10 minutes) and the supernatant was removed by a pipette. The residue was dried *in vacuo* at 40 °C for 16 hours to afford the title compound as a solid.
- 15 HPLC-MS (Method C): m/z: 328 (M+1), 350 (M+23); Rt = 4.09 min.

Example 648 (General procedure (H))


Benzyl-[3-(2H-tetrazol-5-yl)phenyl]amine



- 20 HPLC-MS (Method D): m/z: 252 (M+1); Rt = 3,74 min.

Example 649 (General procedure (H))


(4-Methoxybenzyl)-[3-(2H-tetrazol-5-yl)phenyl]amine



- 25 HPLC-MS (Method D): m/z: 282,2 (M+1); Rt = 3,57min.

Example 650 (General procedure (H))

4-[[3-(2*H*-Tetrazol-5-yl)phenylamino]methyl]phenol




HPLC-MS (Method D): m/z : 268,4 ($M+1$); R_t = 2,64 min.

5

Example 651 (General procedure (H))

(4-Nitrobenzyl)-[3-(2*H*-tetrazol-5-yl)phenyl]amine




HPLC-MS (Method D): m/z : 297,4 ($M+1$); R_t = 3,94 min.

10

Example 652 (General procedure (H))

(4-Chlorobenzyl)-[3-(2*H*-tetrazol-5-yl)phenyl]amine




HPLC-MS (Method D): m/z : 287,2 ($M+1$); R_t = 4,30 min.

15

Example 653 (General procedure (H))

(2-Chlorobenzyl)-[3-(2*H*-tetrazol-5-yl)phenyl]amine




HPLC-MS (Method D): m/z : 286 ($M+1$); R_t = 4,40 min.

20

Example 654 (General procedure (H))


(4-Bromobenzyl)-[3-(2*H*-tetrazol-5-yl)phenyl]amine



HPLC-MS (Method D): m/z:332 (M+1); Rt = 4,50 min.

Example 655 (General procedure (H))


5 (3-Benzyloxybenzyl)-[3-(2*H*-tetrazol-5-yl)phenyl]amine



HPLC-MS (Method D): m/z: 358 (M+1); Rt = 4,94 min.

Example 656 (General procedure (H))


10 Naphthalen-1-ylmethyl-[3-(2*H*-tetrazol-5-yl)phenyl]amine



HPLC-MS (Method D): m/z: 302 (M+1); Rt = 4,70 min.

Example 657 (General procedure (H))


15 Naphthalen-2-ylmethyl-[3-(2*H*-tetrazol-5-yl)phenyl]amine



HPLC-MS (Method D): m/z: 302 (M+1); Rt = 4,60 min.


Example 658 (General procedure (H))

20 4-[[3-(2*H*-Tetrazol-5-yl)phenylamino]methyl]benzoic acid




HPLC-MS (Method D): m/z: 296 (M+1); Rt = 3,24 min.

Example 659 (General procedure (H))

[3-(2*H*-Tetrazol-5-yl)-phenyl]-[3-(3-trifluoromethyl-phenoxy)benzyl]amine


- 5 HPLC-MS (Method D): m/z : 412 ($M+1$); R_t = 5,54 min.

Example 660 (General procedure (H))

(3-Phenoxybenzyl)-[3-(2*H*-tetrazol-5-yl)phenyl]amine


- 10 HPLC-MS (Method D): m/z : 344 ($M+1$); R_t = 5,04 min.

Example 661 (General procedure (H))

(4-Phenoxy-benzyl)-[3-(2*H*-tetrazol-5-yl)phenyl]amine

- 15 HPLC-MS (Method D): m/z : 344 ($M+1$); R_t = 5,00 min.


Example 662 (General procedure (H))

(4-[[3-(2*H*-Tetrazol-5-yl)phenylamino]methyl]phenoxy)acetic acid

- 20 HPLC-MS (Method D): m/z : 326 ($M+1$); R_t = 3,10 min.

Example 663 (General procedure (H))


(4-Benzyloxybenzyl)-[3-(2*H*-tetrazol-5-yl)phenyl]amine



HPLC-MS (Method D): m/z: 358 (M+1); Rt = 4,97 min.

Example 664 (General procedure (H))


- 5 3-(4-([3-(2H-Tetrazol-5-yl)phenylamino]methyl)phenyl)acrylic acid



HPLC-MS (Method D): m/z: 322 (M+1); Rt = 3,60 min.

Example 665 (General procedure (H))


- 10 Dimethyl-(4-([3-(2H-tetrazol-5-yl)phenylamino]methyl)naphthalen-1-yl)amine



HPLC-MS (Method D): m/z: 345 (M+1); Rt = 3,07 min.

Example 666 (General procedure (H))


- 15 (4'-Methoxybiphenyl-4-ylmethyl)-[3-(2H-tetrazol-5-yl)phenyl]amine



HPLC-MS (Method D): m/z: 358 (M+1); Rt = 4,97 min.

Example 667 (General procedure (H))


- 20 (2'-Chlorobiphenyl-4-ylmethyl)-[3-(2H-tetrazol-5-yl)phenyl]amine



HPLC-MS (Method D): m/z : 362 ($M+1$); R_t = 5,27 min.

Example 668 (General procedure (H))

5 Benzyl-[4-(2*H*-tetrazol-5-yl)phenyl]amine




For preparation of starting material, see example 875.

HPLC-MS (Method D): m/z : 252 ($M+1$); R_t = 3,97 min.

10 Example 669 (General procedure (H))


(4-Methoxybenzyl)-[4-(2*H*-tetrazol-5-yl)phenyl]amine



HPLC-MS (Method D): m/z : 282 ($M+1$); R_t = 3,94 min.

15 Example 670 (General procedure (H))


4-[[4-(2*H*-Tetrazol-5-yl)phenylamino]methyl]phenol



HPLC-MS (Method D): m/z : 268 ($M+1$); R_t = 3,14 min.

20 Example 671 (General procedure (H))


(4-Nitrobenzyl)-[4-(2*H*-tetrazol-5-yl)phenyl]amine



HPLC-MS (Method D): m/z: (M+1); Rt = 3,94 min.

Example 672 (General procedure (H))


5 (4-Chlorobenzyl)-[4-(2H-tetrazol-5-yl)phenyl]amine



HPLC-MS (Method D): m/z: (M+1); Rt = 4,47 min.

Example 673 (General procedure (H))


10 (2-Chlorobenzyl)-[4-(2H-tetrazol-5-yl)phenyl]amine



HPLC-MS (Method D): m/z: 286 (M+1); Rt = 4,37 min.

Example 674 (General procedure (H))


15 (4-Bromobenzyl)-[4-(2H-tetrazol-5-yl)phenyl]amine



HPLC-MS (Method D): m/z: 331 (M+1); Rt = 4,57 min.

Example 675 (General procedure (H))


20 (3-Benzyloxybenzyl)-[4-(2H-tetrazol-5-yl)phenyl]amine



HPLC-MS (Method D): m/z: 358 (M+1); Rt = 5,07min.

Example 676 (General procedure (H))

Naphthalen-1-ylmethyl-[4-(2*H*-tetrazol-5-yl)phenyl]amine




5

HPLC-MS (Method D): m/z: 302 (M+1); Rt = 4,70 min.

Example 677 (General procedure (H))

Naphthalen-2-ylmethyl-[4-(2*H*-tetrazol-5-yl)phenyl]amine




10

HPLC-MS (Method D): m/z: 302 (M+1); Rt = 4,70 min.

Example 678 (General procedure (H))

Biphenyl-4-ylmethyl-[4-(2*H*-tetrazol-5-yl)phenyl]amine




15

HPLC-MS (Method D): m/z: 328 (M+1); Rt = 5,07 min.

Example 679 (General procedure (H))

4-[[4-(2*H*-Tetrazol-5-yl)phenylamino]methyl]benzoic acid




20

HPLC-MS (Method D): m/z: 296 (M+1); Rt = 3,34 min.

Example 680 (General procedure (H))

[4-(2*H*-Tetrazol-5-yl)phenyl]-[3-(3-trifluoromethylphenoxy)benzyl]amine




HPLC-MS (Method D): m/z : 412 ($M+1$); R_t = 5,54 min.

5

Example 681 (General procedure (H))

(3-Phenoxybenzyl)-[4-(2*H*-tetrazol-5-yl)phenyl]amine




HPLC-MS (Method D): m/z : 344 ($M+1$); R_t = 5,07 min.

10

Example 682 (General procedure (H))

(4-Phenoxybenzyl)-[4-(2*H*-tetrazol-5-yl)-phenyl]-amine




HPLC-MS (Method D): m/z : 344 ($M+1$); R_t = 5,03 min.

15

Example 683 (General procedure (H))

3-[[4-(2*H*-Tetrazol-5-yl)phenylamino]methyl]benzoic acid




HPLC-MS (Method D): m/z : 286 ($M+1$); R_t = 3,47 min.

20

Example 684 (General procedure (H))


(4-[[4-(2*H*-Tetrazol-5-yl)phenylamino]methyl]phenoxy)acetic acid



HPLC-MS (Method D): m/z: 326 (M+1); Rt = 3,40 min.

Example 685 (General procedure (H))


5 (4-Benzyloxybenzyl)-[4-(2H-tetrazol-5-yl)phenyl]amine



HPLC-MS (Method D): m/z: 358 (M+1); Rt = 5,14 min.

Example 686 (General procedure (H))


10 3-(4-[[4-(2H-Tetrazol-5-yl)phenylamino]methyl]phenyl)acrylic acid



HPLC-MS (Method D): m/z: 322 (M+1); Rt = 3,66 min.

Example 687 (General procedure (H))


15 Dimethyl-(4-[[4-(2H-tetrazol-5-yl)phenylamino]methyl]naphthalen-1-yl)amine



HPLC-MS (Method D): m/z: 345 (M+1); Rt = 3,10 min.


Example 688 (General procedure (H))

20 (4'-Methoxybiphenyl-4-ylmethyl)-[4-(2H-tetrazol-5-yl)phenyl]amine




HPLC-MS (Method D): m/z: 358 (M+1); Rt = 5,04 min.

Example 689 (General procedure (H))

(2'-Chlorobiphenyl-4-ylmethyl)-[4-(2*H*-tetrazol-5-yl)-phenyl]-amine

5 HPLC-MS (Method D): m/z : 362 ($M+1$); R_t = 5,30 min.


General procedure (I) for preparation of compounds of general formula I₈:

wherein K, M and T are as defined above.

10 This procedure is very similar to general procedure (A), the only difference being the carboxylic acid is containing a tetrazole moiety. When the acylation is complete, the product is isolated by extraction, filtration, chromatography or other methods known to those skilled in the art.

15 The general procedure (I) is further illustrated in the following example 690:

Example 690 (General procedure (I))

4-[4-(2*H*-Tetrazol-5-yl)benzoylamino]benzoic acid

20 To a solution of 4-(2*H*-tetrazol-5-yl)benzoic acid (example 605, 4 mmol) and HOAt (4.2 mmol) in DMF (6 mL) was added 1-ethyl-3-(3'-dimethylaminopropyl)carbodiimide hydrochloride (4.2 mmol) and the resulting mixture was stirred at room temperature for 1 hour. An aliquot of this HOAt-ester solution (0.45 mL) was mixed with 0.25 mL of a solution of


4-aminobenzoic acid (1.2 mmol in 1 mL DMF). (Anilines as hydrochlorides can also be utilised, a slight excess of triethylamine was added to the hydrochloride suspension in DMF prior to mixing with the HOAt-ester.) The resulting mixture was shaken for 3 days at room temperature. 1N hydrochloric acid (2 mL) was added and the mixture was shaken for 16 hours at room temperature. The solid was isolated by centrifugation (alternatively by filtration or extraction) and was washed with water (3 mL). Drying *in vacuo* at 40 °C for 2 days afforded the title compound.

HPLC-MS (Method D): m/z: 310 (M+1); Rt = 2.83 min.

10

Example 691 (General procedure (I))

3-[4-(2H-Tetrazol-5-yl)benzoylamino]benzoic acid




HPLC-MS (Method D): m/z: 310 (M+1); Rt = 2.89 min.

15

Example 692 (General procedure (I))

3-{4-[4-(2H-Tetrazol-5-yl)benzoylamino]phenyl}acrylic acid




HPLC-MS (Method D): m/z: 336 (M+1); Rt = 3.10 min.

20

Example 693 (General procedure (I))

3-{4-[4-(2H-Tetrazol-5-yl)benzoylamino]phenyl}propionic acid




HPLC-MS (Method D): m/z: 338 (M+1); Rt = 2.97 min.

25

Example 694 (General procedure (I))

3-Methoxy-4-[4-(2*H*-tetrazol-5-yl)benzoylamino]benzoic acid




HPLC-MS (Method D): *m/z*: 340 (*M*+1); *R*_t = 3.03 min.

5

Example 695 (General procedure (I))

N-(4-Benzyloxyphenyl)-4-(2*H*-tetrazol-5-yl)benzamide




HPLC-MS (Method D): *m/z*: 372 (*M*+1); *R*_t = 4.47 min.

10

Example 696 (General procedure (I))

N-(4-Phenoxyphenyl)-4-(2*H*-tetrazol-5-yl)benzamide




HPLC-MS (Method D): *m/z*: 358 (*M*+1); *R*_t = 4.50 min.

15

Example 697 (General procedure (I))

N-(9*H*-Fluoren-2-yl)-4-(2*H*-tetrazol-5-yl)benzamide




HPLC-MS (Method D): *m/z*: 354 (*M*+1); *R*_t = 4.60 min.

20

Example 698 (General procedure (I))


N-(9-Ethyl-9*H*-carbazol-2-yl)-4-(2*H*-tetrazol-5-yl)benzamide



HPLC-MS (Method D): m/z : 383 ($M+1$); R_t = 4.60 min.

Example 699 (General procedure (I))


5 *N*-Phenyl-4-(2*H*-tetrazol-5-yl)benzamide



HPLC-MS (Method D): m/z : 266 ($M+1$); R_t = 3.23 min.

Example 700 (General procedure (I))

10 4-[4-(2*H*-Tetrazol-5-ylmethoxy)benzoylamino]benzoic acid




The starting material was prepared as described in example 592.

HPLC-MS (Method D): m/z : 340 ($M+1$); R_t = 2.83 min.

15 Example 701 (General procedure (I))


3-[4-(2*H*-Tetrazol-5-ylmethoxy)benzoylamino]benzoic acid



HPLC-MS (Method D): m/z : 340 ($M+1$); R_t = 2.90 min.

20 Example 702 (General procedure (I))


3-{4-[4-(2*H*-Tetrazol-5-ylmethoxy)benzoylamino]phenyl}acrylic acid



HPLC-MS (Method D): m/z: 366 (M+1); Rt = 3.07 min.

Example 703 (General procedure (I))


- 5 3-{4-[4-(2H-Tetrazol-5-ylmethoxy)benzoylamino]phenyl}propionic acid



HPLC-MS (Method D): m/z: 368 (M+1); Rt = 2.97 min.

Example 704 (General procedure (I))


- 10 3-Methoxy-4-[4-(2H-tetrazol-5-ylmethoxy)benzoylamino]benzoic acid



HPLC-MS (Method D): m/z: 370 (M+1); Rt = 3.07 min.

Example 705 (General procedure (I))


- 15 N-(4-Benzyloxyphenyl)-4-(2H-tetrazol-5-ylmethoxy)benzamide



HPLC-MS (Method D): m/z: 402 (M+1); Rt = 4.43 min.

Example 706 (General procedure (I))


- 20 N-(4-Phenoxyphenyl)-4-(2H-tetrazol-5-ylmethoxy)benzamide



HPLC-MS (Method D): m/z: 388 (M+1); Rt = 4.50 min.

Example 707 (General procedure (I))

N-(9H-Fluoren-2-yl)-4-(2*H*-tetrazol-5-ylmethoxy)benzamide




5

HPLC-MS (Method D): m/z: 384 (M+1); Rt = 4.57 min.

Example 708 (General procedure (I))

N-(9-Ethyl-9H-carbazol-2-yl)-4-(2*H*-tetrazol-5-ylmethoxy)benzamide




10

HPLC-MS (Method D): m/z: 413 (M+1); Rt = 4.57 min.

Example 709 (General procedure (I))

N-Phenyl-4-(2*H*-tetrazol-5-ylmethoxy)benzamide




15

HPLC-MS (Method D): m/z: 296 (M+1); Rt = 3.23 min.

Example 710 (General procedure (I))

4-[4-(2*H*-Tetrazol-5-ylmethylsulfanyl)benzoylamino]benzoic acid




20

The starting material was prepared as described in example 593.

HPLC-MS (Method D): m/z: 356 (M+1); Rt = 2.93 min.

Example 711 (General procedure (I))

3-[4-(2H-Tetrazol-5-ylmethylsulfanyl)benzoylamino]benzoic acid




HPLC-MS (Method D): m/z: 356 (M+1); Rt = 3.00 min.

5

Example 712 (General procedure (I))

3-[4-[4-(2H-Tetrazol-5-ylmethylsulfanyl)benzoylamino]phenyl]acrylic acid




HPLC-MS (Method D): m/z: 382 (M+1); Rt = 3.26 min.

10

Example 713 (General procedure (I))

3-[4-[4-(2H-Tetrazol-5-ylmethylsulfanyl)benzoylamino]phenyl]propionic acid




HPLC-MS (Method D): m/z: 384 (M+1); Rt = 3.10 min.

15

Example 714 (General procedure (I))

3-Methoxy-4-[4-(2H-tetrazol-5-ylmethylsulfanyl)benzoylamino]benzoic acid




HPLC-MS (Method D): m/z: 386 (M+1); Rt = 3.20 min.

20

Example 715 (General procedure (I))


N-(4-Benzyloxyphenyl)-4-(2H-tetrazol-5-ylmethylsulfanyl)benzamide



HPLC-MS (Method D): m/z : 418 ($M+1$); R_t = 4.57 min.

Example 716 (General procedure (I))


- 5 *N*-(4-Phenoxyphenyl)-4-(2*H*-tetrazol-5-ylmethylsulfanyl)benzamide



HPLC-MS (Method D): m/z : 404 ($M+1$); R_t = 4.60 min.

Example 717 (General procedure (I))


- 10 *N*-(9*H*-Fluoren-2-yl)-4-(2*H*-tetrazol-5-ylmethylsulfanyl)benzamide



HPLC-MS (Method D): m/z : 400 ($M+1$); R_t = 4.67 min.

Example 718 (General procedure (I))


- 15 *N*-(9-Ethyl-9*H*-carbazol-2-yl)-4-(2*H*-tetrazol-5-ylmethylsulfanyl)benzamide



HPLC-MS (Method D): m/z : 429 ($M+1$); R_t = 4.67 min.


Example 719 (General procedure (I))

- 20 *N*-Phenyl-4-(2*H*-tetrazol-5-ylmethylsulfanyl)benzamide



HPLC-MS (Method D): m/z : 312 ($M+1$); R_t = 3.40 min.


General procedure (J) for solution phase preparation of amides of general formula I₉:



wherein T is as defined above.

This general procedure (J) is further illustrated in the following example.

- 10 **Example 720 (General procedure (J)).**
 9-(3-Chlorobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole



- 15 3-(2H-Tetrazol-5-yl)-9H-carbazole (example 594, 17 g, 72.26 mmol) was dissolved in *N,N*-dimethylformamide (150 mL). Triphenylmethyl chloride (21.153 g, 75.88 mmol) and triethylamine (20.14 mL, 14.62 g, 144.50 mmol) were added consecutively. The reaction mixture was stirred for 18 hours at room temperature, poured into water (1.5 L) and stirred for an additional 1 hour. The crude product was filtered off and dissolved in dichloromethane (500 mL). The organic phase was washed with water (2 x 250 mL) and dried with magnesium sulfate (1 h). Filtration followed by concentration yielded a solid which was
- 20 triturated in heptanes (200 mL). Filtration furnished 3-[2-(triphenylmethyl)-2H-tetrazol-5-yl]-9H-carbazole (31.5 g) which was used without further purification.
- ¹H-NMR (CDCl₃): δ 8.87 (1H, d), 8.28 (1H, bs), 8.22 (1H, dd), 8.13 (1H, d), 7.49 (1H, d), 7.47-7.19 (18H, m); HPLC-MS (Method C): m/z : 243 (triphenylmethyl); R_t = 5.72 min.

3-[2-(Triphenylmethyl)-2*H*-tetrazol-5-yl]-9*H*-carbazole (200 mg, 0.42 mmol) was dissolved in methyl sulfoxide (1.5 mL). Sodium hydride (34 mg, 60 %, 0.85 mmol) was added, and the resulting suspension was stirred for 30 min at room temperature. 3-Chlorobenzyl chloride (85 μ L, 108 mg, 0.67 mmol) was added, and the stirring was continued at 40 °C for 18 hours.

- 5 The reaction mixture was cooled to ambient temperature and poured into 0.1 N hydrochloric acid (aq.) (15 mL). The precipitated solid was filtered off and washed with water (3 x 10 mL) to furnish 9-(3-chlorobenzyl)-3-[2-(triphenylmethyl)-2*H*-tetrazol-5-yl]-9*H*-carbazole, which was dissolved in a mixture of tetrahydrofuran and 6 N hydrochloric acid (aq.) (9:1) (10 mL) and stirred at room temperature for 18 hours. The reaction mixture was poured into water (100
10 mL). The solid was filtered off and rinsed with water (3 x 10 mL) and dichloromethane (3 x 10 mL) to yield the title compound (127 mg). No further purification was necessary.


¹H-NMR (DMSO-*d*₆): δ 8.89 (1H, d), 8.29 (1H, d), 8.12 (1H, dd), 7.90 (1H, d), 7.72 (1H, d), 7.53 (1H, t), 7.36-7.27 (4H, m), 7.08 (1H, bt), 5.78 (2H, s); HPLC-MS (Method B): *m/z*: 360 (M+1); *R*_t = 5.07 min.

15

The compounds in the following examples were prepared in a similar fashion. Optionally, the compounds can be further purified by recrystallization from e.g. aqueous sodium hydroxide (1 N) or by chromatography.

- 20 Example 721 (General Procedure (J)).


9-(4-Chlorobenzyl)-3-(2*H*-tetrazol-5-yl)-9*H*-carbazole



HPLC-MS (Method C): *m/z*: 360 (M+1); *R*_t = 4.31 min.

- 25 Example 722 (General Procedure (J)).


9-(4-Methylbenzyl)-3-(2*H*-tetrazol-5-yl)-9*H*-carbazole



HPLC-MS (Method C): m/z : 340 ($M+1$); R_t = 4.26 min.

Example 723 (General Procedure (J)).

3-(2*H*-Tetrazol-5-yl)-9-(4-trifluoromethylbenzyl)-9*H*-carbazole




5

HPLC-MS (Method C): m/z : 394 ($M+1$); R_t = 4.40 min.

Example 724 (General Procedure (J)).

9-(4-Benzyloxybenzyl)-3-(2*H*-tetrazol-5-yl)-9*H*-carbazole




10

HPLC-MS (Method C): m/z : 432 ($M+1$); R_t = 4.70 min.

Example 725 (General Procedure (J)).

9-(3-Methylbenzyl)-3-(2*H*-tetrazol-5-yl)-9*H*-carbazole




15

HPLC-MS (Method C): m/z : 340 ($M+1$); R_t = 4.25 min.

Example 726 (General Procedure (J)).


9-Benzyl-3-(2*H*-tetrazol-5-yl)-9*H*-carbazole



$^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 8.91 (1H, dd), 8.30 (1H, d), 8.13 (1H, dd), 7.90 (1H, d), 7.73 (1H, d), 7.53 (1H, t), 7.36-7.20 (6H, m), 5.77 (2H, s).

5 Example 727 (General Procedure (J)).

9-(4-Phenylbenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole




$^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 8.94 (1H, s), 8.33 (1H, d), 8.17 (1H, dd), 7.95 (1H, d), 7.77 (1H, d), 7.61-7.27 (11H, m), 5.82 (2H, s).

10

Example 728 (General Procedure (J)).

9-(3-Methoxybenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole




HPLC-MS (Method C): m/z : 356 ($M+1$); R_t = 3.99 min.

15

Example 729 (General Procedure (J)).


9-(Naphthalen-2-ylmethyl)-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method C): m/z : 376 ($M+1$); R_t = 4.48 min.

Example 730 (General Procedure (J)).


5 9-(3-Bromobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method C): m/z : 404 ($M+1$); R_t = 4.33 min.

Example 731 (General Procedure (J)).


10 9-(Biphenyl-2-ylmethyl)-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method C): m/z : 402 ($M+1$); R_t = 4.80 min.


Example 732 (General Procedure (J)).

15 3-(2H-Tetrazol-5-yl)-9-[4-(1,2,3-thiadiazol-4-yl)benzyl]-9H-carbazole



Example 733 (General Procedure (J)).

9-(2'-Cyanobiphenyl-4-ylmethyl)-3-(2H-tetrazol-5-yl)-9H-carbazole




5

¹H-NMR (DMSO-*d*₆): δ 8.91 (1H, d), 8.31 (1H, d), 8.13 (1H, dd), 7.95 (1H, d), 7.92 (1H, d), 7.78 (1H, d), 7.75 (1H, dt), 7.60-7.47 (5H, m), 7.38-7.28 (3H, m), 5.86 (2H, s); HPLC-MS (Method C): *m/z*: 427 (M+1); *R*_t = 4.38 min.

10 Example 734 (General Procedure (J)).


9-(4-Iodobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method C): *m/z*: 452 (M+1); *R*_t = 4.37 min.

15 Example 735 (General Procedure (J)).


9-(3,5-Bis(trifluoromethyl)benzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method C): m/z : 462 ($M+1$); R_t = 4.70 min.

Example 736 (General Procedure (J)).

5 9-(4-Bromobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole




$^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 8.89 (1H, d), 8.29 (1H, d), 8.11 (1H, dd), 7.88 (1H, d), 7.70 (1H, d), 7.52 (1H, t), 7.49 (2H, d), 7.31 (1H, t), 7.14 (2H, d), 5.74 (2H, s); HPLC-MS (Method C): m/z : 404 ($M+1$); R_t = 4.40 min.

10

Example 737 (General Procedure (J)).

9-(Anthracen-9-ylmethyl)-3-(2H-tetrazol-5-yl)-9H-carbazole




HPLC-MS (Method C): m/z : 426 ($M+1$); R_t = 4.78 min.

15

Example 738 (General Procedure (J)).

9-(4-Carboxybenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole




3.6 fold excess sodium hydride was used.

¹H-NMR (DMSO-*d*₆): δ 12.89 (1H, bs), 8.89 (1H, d), 8.30 (1H, d), 8.10 (1H, dd), 7.87 (1H, d),
 5 7.86 (2H, d), 7.68 (1H, d), 7.51 (1H, t), 7.32 (1H, t), 7.27 (2H, d), 5.84 (2H, s); HPLC-MS
 (Method C): m/z: 370 (M+1); Rt = 3.37 min.

Example 739 (General Procedure (J)).

9-(2-Chlorobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole




10

HPLC-MS (Method B): m/z: 360 (M+1); Rt = 5.30 min.

Example 740 (General Procedure (J)).

9-(4-Fluorobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole




15

¹H-NMR (DMSO-*d*₆): δ 8.88 (1H, d), 8.28 (1H, d), 8.10 (1H, dd), 7.89 (1H, d), 7.72 (1H, d),
 7.52 (1H, t), 7.31 (1H, t), 7.31-7.08 (4H, m), 5.74 (2H, s); HPLC-MS (Method C): m/z: 344
 (M+1); Rt = 4.10 min.

20 Example 741 (General Procedure (J)).

9-(3-Fluorobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole




$^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 8.89 (1H, d), 8.29 (1H, d), 8.12 (1H, dd), 7.90 (1H, d), 7.72 (1H, d), 7.53 (1H, t), 7.37-7.27 (2H, m), 7.12-7.02 (2H, m), 6.97 (1H, d), 5.78 (2H, s); HPLC-MS (Method C): m/z : 344 ($M+1$); R_t = 4.10 min.

5

Example 742 (General Procedure (J)).

9-(2-Iodobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole




HPLC-MS (Method C): m/z : 452 ($M+1$); R_t = 4.58 min.

10

Example 743 (General Procedure (J)).

9-(3-Carboxybenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole




3.6 fold excess sodium hydride was used.

15

$^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 12.97 (1H, bs), 8.90 (1H, bs), 8.30 (1H, d), 8.12 (1H, bd), 7.89 (1H, d), 7.82 (1H, m), 7.77 (1H, bs), 7.71 (1H, d), 7.53 (1H, t), 7.46-7.41 (2H, m), 7.32 (1H, t), 5.84 (2H, s); HPLC-MS (Method C): m/z : 370 ($M+1$); R_t = 3.35 min.

Example 744 (General Procedure (J)).


9-[4-(2-Propyl)benzyl]-3-(2*H*-tetrazol-5-yl)-9*H*-carbazole



5 ¹H-NMR (DMSO-*d*₆): δ 8.87 (1H, d), 8.27 (1H, d), 8.10 (1H, dd), 7.87 (1H, d), 7.71 (1H, d), 7.51 (1H, t), 7.31 (1H, t), 7.15 (2H, d), 7.12 (2H, d), 5.69 (2H, s), 2.80 (1H, sept), 1.12 (6H, d); HPLC-MS (Method C): *m/z*: 368 (M+1); *R*_t = 4.73 min.

Example 745 (General Procedure (J)).

9-(3,5-Dimethoxybenzyl)-3-(2*H*-tetrazol-5-yl)-9*H*-carbazole




10

HPLC-MS (Method C): *m/z*: 386 (M+1); *R*_t = 4.03 min.

Example 746 (General Procedure (J)).

3-(2*H*-Tetrazol-5-yl)-9-(2,4,5-trifluorobenzyl)-9*H*-carbazole




15

HPLC-MS (Method B): *m/z*: 380 (M+1); *R*_t = 5.00 min.

Example 747 (General Procedure (J)).


N-Methyl-*N*-phenyl-2-[3-(2*H*-tetrazol-5-yl)carbazol-9-yl]acetamide



HPLC-MS (Method B): m/z : 383 ($M+1$); R_t = 4.30 min.

Example 748 (General Procedure (J)).

5 9-(4-Methoxybenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole




$^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 8.86 (1H, d), 8.26 (1H, d), 8.10 (1H, dd), 7.90 (1H, d), 7.73 (1H, d), 7.51 (1H, t), 7.30 (1H, t), 7.18 (2H, d), 6.84 (2H, d), 5.66 (2H, s), 3.67 (3H, s); HPLC-MS (Method B): m/z : 356 ($M+1$); R_t = 4.73 min.

10

Example 749 (General Procedure (J)).


9-(2-Methoxybenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole



15 $^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 8.87 (1H, d), 8.27 (1H, d), 8.09 (1H, dd), 7.77 (1H, d), 7.60 (1H, d), 7.49 (1H, t), 7.29 (1H, t), 7.23 (1H, bt), 7.07 (1H, bd), 6.74 (1H, bt), 6.61 (1H, bd), 5.65 (2H, s), 3.88 (3H, s); HPLC-MS (Method B): m/z : 356 ($M+1$); R_t = 4.97 min.

Example 750 (General Procedure (J)).


9-(4-Cyanobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method C): m/z: 351 (M+1); Rt = 3.74 min.

Example 751 (General Procedure (J)).


5 9-(3-Cyanobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method C): m/z: 351 (M+1); Rt = 3.73 min.

Example 752 (General Procedure (J)).

10 9-(5-Chloro-2-methoxybenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole




$^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 8.87 (1H, d), 8.35 (1H, d), 8.10 (1H, dd), 7.73 (1H, d), 7.59 (1H, d), 7.49 (1H, t), 7.29 (1H, t), 7.27 (1H, dd), 7.11 (1H, d), 6.51 (1H, d), 5.63 (2H, s), 3.88 (3H, s);
HPLC-MS (Method C): m/z: 390 (M+1); Rt = 4.37 min.

15

Example 753 (General Procedure (J)).

N-Phenyl-2-[3-(2H-tetrazol-5-yl)carbazol-9-yl]acetamide




¹H-NMR (DMSO-*d*₆): δ 10.54 (1H, s), 8.87 (1H, bs), 8.27 (1H, d), 8.12 (1H, bd), 7.83 (1H, d), 7.66 (1H, d), 7.61 (2H, d), 7.53 (1H, t), 7.32 (1H, t), 7.32 (2H, t), 7.07 (1H, t), 5.36 (2H, s); HPLC-MS (Method C): *m/z*: 369 (M+1); *R*_t = 3.44 min.

5

Example 754 (General Procedure (J)).

N-Butyl-2-[3-(2*H*-tetrazol-5-yl)carbazol-9-yl]acetamide




¹H-NMR (DMSO-*d*₆): δ 8.85 (1H, d), 8.31 (1H, t), 8.25 (1H, d), 8.10 (1H, dd), 7.75 (1H, d), 7.58 (1H, d), 7.52 (1H, t), 7.30 (1H, t), 5.09 (2H, s), 3.11 (2H, q), 1.42 (2H, quint), 1.30 (2H, sext), 0.87 (3H, t); HPLC-MS (Method C): *m/z*: 349 (M+1); *R*_t = 3.20 min.

10

Example 755 (General Procedure (J)).

9-(2,4-Dichlorobenzyl)-3-(2*H*-tetrazol-5-yl)-9*H*-carbazole




15

¹H-NMR (DMSO-*d*₆): δ 8.92 (1H, d), 8.32 (1H, d), 8.09 (1H, dd), 7.76 (1H, d), 7.74 (1H, d), 7.58 (1H, d), 7.51 (1H, t), 7.33 (1H, t), 7.23 (1H, dd), 6.42 (1H, d), 5.80 (2H, s); HPLC-MS (Method B): *m/z*: 394 (M+1); *R*_t = 5.87 min.

Example 756 (General Procedure (J)).


9-(2-Methylbenzyl)-3-(2*H*-tetrazol-5-yl)-9*H*-carbazole



¹H-NMR (DMSO-*d*₆): δ 8.92 (1H, d), 8.32 (1H, d), 8.08 (1H, dd), 7.72 (1H, d), 7.55 (1H, d), 7.48 (1H, t), 7.32 (1H, t), 7.26 (1H, d), 7.12 (1H, t), 6.92 (1H, t), 6.17 (1H, d), 5.73 (2H, s), 2.46 (3H, s); HPLC-MS (Method B): *m/z*: 340 (*M*+1); *R*_t = 5.30 min.

Example 757 (General Procedure (J)).

9-(3-Nitrobenzyl)-3-(2*H*-tetrazol-5-yl)-9*H*-carbazole




10

HPLC-MS (Method C): *m/z*: 371 (*M*+1); *R*_t = 3.78 min.

Example 758 (General Procedure (J)).

9-(3,4-Dichlorobenzyl)-3-(2*H*-tetrazol-5-yl)-9*H*-carbazole




15

HPLC-MS (Method B): *m/z*: 394 (*M*+1); *R*_t = 5.62 min.

Example 759 (General Procedure (J)).

9-(2,4-Difluorobenzyl)-3-(2*H*-tetrazol-5-yl)-9*H*-carbazole




¹H-NMR (DMSO-*d*₆): δ 8.89 (1H, d), 8.29 (1H, d), 8.11 (1H, dd), 7.88 (1H, d), 7.69 (1H, d), 7.52 (1H, t), 7.36-7.24 (2H, m), 7.06-6.91 (2H, m), 5.78 (2H, s); HPLC-MS (Method B): *m/z*: 362 (M+1); Rt = 5.17 min.

5

Example 760 (General Procedure (J)).

9-(3,5-Difluorobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole




¹H-NMR (DMSO-*d*₆): δ 8.90 (1H, bs), 8.31 (1H, d), 8.13 (1H, bd), 7.90 (1H, d), 7.73 (1H, d), 7.54 (1H, t), 7.34 (1H, t), 7.14 (1H, t), 6.87 (2H, bd), 5.80 (2H, s); HPLC-MS (Method B): *m/z*: 362 (M+1); Rt = 5.17 min.

10

Example 761 (General Procedure (J)).

9-(3,4-Difluorobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole




¹H-NMR (DMSO-*d*₆): δ 8.89 (1H, bs), 8.29 (1H, d), 8.12 (1H, bd), 7.92 (1H, d), 7.74 (1H, d), 7.54 (1H, t), 7.42-7.25 (3H, m), 6.97 (1H, bm), 5.75 (2H, s); HPLC-MS (Method B): *m/z*: 362 (M+1); Rt = 5.17 min.

15

20 Example 762 (General Procedure (J)).


9-(3-Iodobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method B): m/z : 452 ($M+1$); R_t = 5.50 min.

Example 763 (General Procedure (J)).

5 3-(2H-Tetrazol-5-yl)-9-[3-(trifluoromethyl)benzyl]-9H-carbazole




$^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ 8.89 (1H, d), 8.30 (1H, d), 8.11 (1H, dd), 7.90 (1H, d), 7.72 (1H, d), 7.67 (1H, bs), 7.62 (1H, bd), 7.53 (1H, t), 7.50 (1H, bt), 7.33 (1H, bd), 7.32 (1H, t), 5.87 (2H, s); HPLC-MS (Method B): m/z : 394 ($M+1$); R_t = 5.40 min.

10

Example 764 (General Procedure (J)).

N-(4-Carboxyphenyl)-2-[3-(2H-tetrazol-5-yl)carbazol-9-yl]acetamide




15 3.6 fold excess sodium hydride was used.

HPLC-MS (Method B): m/z : 413 ($M+1$); R_t = 3.92 min.

Example 765 (General Procedure (J)).


20 *N*-(2-Propyl)-2-[3-(2H-tetrazol-5-yl)carbazol-9-yl]acetamide



HPLC-MS (Method B): m/z : 335 ($M+1$); R_t = 3.70 min.

Example 766 (General Procedure (J)).


5 *N*-Benzyl-*N*-phenyl-2-[3-(2*H*-tetrazol-5-yl)carbazol-9-yl]acetamide



HPLC-MS (Method B): m/z : 459 ($M+1$); R_t = 5.37 min.

Example 767 (General Procedure (J)).


10 *N*-[4-(2-Methyl-2-propyl)phenyl]-2-[3-(2*H*-tetrazol-5-yl)carbazol-9-yl]acetamide



HPLC-MS (Method B): m/z : 425 ($M+1$); R_t = 5.35 min.

Example 768 (General Procedure (J)).


15 *N*-Phenethyl-2-[3-(2*H*-tetrazol-5-yl)carbazol-9-yl]acetamide



HPLC-MS (Method C): m/z: 397 (M+1); Rt = 3.43 min.

Example 769 (General Procedure (J)).


5 3-(2H-Tetrazol-5-yl)-9-[2-(trifluoromethyl)benzyl]-9H-carbazole



HPLC-MS (Method C): m/z: 394 (M+1); Rt = 4.44 min.

Example 770 (General Procedure (J)).


10 9-[2-Fluoro-6-(trifluoromethyl)benzyl]-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method C): m/z: 412 (M+1); Rt = 4.21 min.

Example 771 (General Procedure (J)).


15 9-[2,4-Bis(trifluoromethyl)benzyl]-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method C): m/z : 462 ($M+1$); R_t = 4.82 min.

Example 772 (General Procedure (J)).


- 5 3-(2H-Tetrazol-5-yl)-9-(2,4,6-trimethylbenzyl)-9H-carbazole



HPLC-MS (Method C): m/z : 368 ($M+1$); R_t = 4.59 min.

Example 773 (General Procedure (J)).


- 10 9-(2,3,5,6-Tetramethylbenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method C): m/z : 382 ($M+1$); R_t = 4.47 min.

Example 774 (General Procedure (J)).


- 15 9-[(Naphthalen-1-yl)methyl]-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method C): m/z : 376 ($M+1$); R_t = 4.43 min.

Example 775 (General Procedure (J)).


5 9-[Bis(4-fluorophenyl)methyl]-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method C): m/z : 438 ($M+1$); R_t = 4.60 min.

Example 776 (General Procedure (J)).


10 9-(2-Bromobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method C): m/z : 404 ($M+1$); R_t = 4.50 min.

Example 777 (General Procedure (J)).


15 9-(2-Fluorobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method C): m/z: 344 (M+1); Rt = 4.09 min.

Example 778 (General Procedure (J)).

5 9-(4-Carboxy-2-methylbenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole




In this preparation, a 3.6-fold excess of sodium hydride was used.

HPLC-MS (Method C): m/z: 384 (M+1); Rt = 3.56 min.

10 Example 779 (General Procedure (J)).


9-(2-Phenylethyl)-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method C): m/z: 340 (M+1); Rt = 4.08 min.

15 Example 780 (General Procedure (J)).


9-[2-Fluoro-5-(trifluoromethyl)benzyl]-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method C): m/z : 412 ($M+1$); R_t = 4.34 min.

Example 781 (General Procedure (J)).

5 9-(4-Carboxy-2-fluorobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole




3-Fluoro-4-methylbenzoic acid (3.0 g, 19.5 mmol) and benzoyl peroxide (0.18 g, 0.74 mmol) were suspended in benzene. The mixture was purged with N_2 and heated to reflux. *N*-Bromosuccinimide (3.47 g, 19.5 mmol) was added portionwise, and reflux was maintained for 10 18 hours. The reaction mixture was concentrated, and the residue was washed with water (20 mL) at 70 °C for 1 hour. The crude product was isolated by filtration and washed with additional water (2 x 10 mL). The dry product was recrystallized from heptanes. Filtration furnished 4-bromomethyl-3-fluorobenzoic acid (1.92 g) which was used in the following step according to General Procedure (J).

15 In this preparation, a 3.6-fold excess of sodium hydride was used.

HPLC-MS (Method C): m/z : 388 ($M+1$); R_t = 3.49 min.

Example 782 (General Procedure (J)).

5-{4-[[3-(2H-Tetrazol-5-yl)carbazol-9-yl)methyl]naphthalen-1-yl]oxy}pentanoic Acid



20

5-[(4-Formylnaphthalen-1-yl)oxy]pentanoic acid intermediate obtained in example 470 (3.0 g, 11.0 mmol) was dissolved in a mixture of methanol and tetrahydrofuran (9:1) (100 mL), and


sodium borohydride (1.67 g, 44.1 mmol) was added portionwise at ambient temperature. After 30 minutes, the reaction mixture was concentrated to 50 mL and added to hydrochloric acid (0.1 N, 500 mL). Additional hydrochloric acid (1 N, 40 mL) was added, and 5-[(4-hydroxymethyl-naphthalen-1-yl)oxy]pentanoic acid (2.90 g) was collected by filtration. To the
5 crude product was added concentrated hydrochloric acid (100 mL), and the suspension was stirred vigorously for 48 hours at room temperature. The crude product was filtered off and washed with water, until the pH was essentially neutral. The material was washed with heptanes to furnish 5-[(4-chloromethylnaphthalen-1-yl)oxy]pentanoic acid (3.0 g) which was used in the following step according to General Procedure (J).

10

In this preparation, a 3.6-fold excess of sodium hydride was used.
HPLC-MS (Method C): m/z: 492 (M+1); Rt = 4.27 min.

Example 783 (General procedure (J))


15 9-(2,3-Difluorobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method C): m/z= 362 (M+1); Rt = 4.13 min.

Example 784 (General procedure (J))


20 9-(2,5-Difluorobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole



HPLC-MS (Method C): m/z = 362 (M+1); Rt = 4.08 min.

Example 785 (General procedure (J))


9-Pentafluorophenylmethyl-3-(2H-tetrazol-5-yl)-9H-carbazole

HPLC-MS (Method C): $m/z = 416$ ($M+1$); $R_t = 4.32$ min.

5

Example 786 (General procedure (J))

9-(2,6-Difluorobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole

HPLC-MS (Method C): $m/z = 362$ ($M+1$); $R_t = 3.77$ min.

10


Further compounds of the invention that may be prepared according to general procedure (J), and includes:

| | | |
|------------------------|------------------------|------------------------|
| Example 787 | Example 788 | Example 789 |
| Example 790 | Example 791 | Example 792 |
| Example 793 | Example 794 | Example 795 |
| Example 796 | Example 797 | Example 798 |
| Example 799 | | |

The following compounds of the invention may be prepared eg. from 9-(4-bromobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole (example 736) or from 9-(3-bromobenzyl)-3-(2H-tetrazol-5-yl)-9H-carbazole (example 730) and aryl boronic acids *via* the Suzuki coupling reaction eg as described in Littke, Dai & Fu *J. Am. Chem. Soc.*, 2000, 122, 4020-8 (or references cited therein), or using the methodology described in general procedure (E), optionally changing the palladium catalyst to bis(tri-*tert*-butylphosphine)palladium (0).

| | | |
|------------------------|------------------------|------------------------|
| Example 800 | Example 801 | Example 802 |
| Example 803 | Example 804 | Example 805 |

General procedure (K) for preparation of compounds of general formula I₁₀:




5

wherein T is as defined above.

The general procedure (K) is further illustrated by the following example:

Example 806 (General procedure (K)).

10 1-Benzyl-5-(2H-tetrazol-5-yl)-1H-indole



5-Cyanoindole (1.0 g; 7.0 mmol) was dissolved in *N,N*-dimethylformamide (14 mL) and cooled in an ice-water bath. Sodium hydride (0.31 g, 60 %, 7.8 mmol) was added, and the resulting suspension was stirred for 30 min. Benzyl chloride (0.85 mL, 0.94 g, 7.4 mmol) was

added, and the cooling was discontinued. The stirring was continued for 65 hours at room temperature. Water (150 mL) was added, and the mixture was extracted with ethyl acetate (3 x 25 mL). The combined organic phases were washed with brine (30 mL) and dried with sodium sulfate (1 hour). Filtration and concentration yielded the crude material. Purification
5 by flash chromatography on silica gel eluting with ethyl acetate/heptanes = 1:3 afforded 1.60 g 1-benzyl-1*H*-indole-5-carbonitrile.

HPLC-MS (Method C): m/z : 233 ($M+1$); R_t = 4.17 min.

10 1-Benzyl-1*H*-indole-5-carbonitrile was transformed into 1-benzyl-5-(2*H*-tetrazol-5-yl)-1*H*-indole by the method described in general procedure (J) and in example 594. Purification was done by flash chromatography on silica gel eluting with dichloromethane/methanol = 9:1.


HPLC-MS (Method C): m/z : 276 ($M+1$); R_t = 3.35 min.

15

The compounds in the following examples were prepared by the same procedure.

Example 807 (General procedure (K)).

1-(4-Bromobenzyl)-5-(2*H*-tetrazol-5-yl)-1*H*-indole




20

HPLC-MS (Method C): m/z : 354 ($M+1$); R_t = 3.80 min.

Example 808 (General procedure (K)).

1-(4-Phenylbenzyl)-5-(2*H*-tetrazol-5-yl)-1*H*-indole



25

$^1\text{H-NMR}$ (200 MHz, $\text{DMSO}-d_6$): δ = 5.52 (2H, s), 6.70 (1H, d), 7.3-7.45 (6H, m), 7.6 (4H, m), 7.7-7.8 (2H, m), 7.85 (1H, dd), 8.35 (1H, d).


Calculated for $C_{22}H_{17}N_5$, H_2O :

73.32% C; 5.03% H; 19.43% N. Found:

73.81% C; 4.90% H; 19.31% N.

5 Example 809

4'-[5-(2H-Tetrazol-5-yl)indol-1-ylmethyl]biphenyl-4-carboxylic acid



5-(2H-Tetrazol-5-yl)-1H-indole (Syncom BV, Groningen, NL) (1.66g; 8.9 mmol) was treated with trityl chloride (2.5 g, 8.9 mmol) and triethyl amine (2.5 mL, 17.9 mmol) in DMF (25 mL) by stirring at RT overnight. The resulting mixture was treated with water. The gel was isolated, dissolved in methanol, treated with activated carbon; filtered and evaporated to dryness *in vacuo*. This afforded 3.6 g (94%) of crude 5-(2-trityl-2H-tetrazol-5-yl)-1H-indole.

HPLC-MS (Method C): m/z = 450 ($M+23$); R_t = 5.32 min.

15

4-Methylphenylbenzoic acid (5 g, 23.5 mmol) was mixed with CCl_4 (100 mL) and under an atmosphere of nitrogen, the slurry was added *N*-Bromosuccinimide (4.19 g, 23.55 mmol) and dibenzoyl peroxide (0.228 g, 0.94 mmol). The mixture was subsequently heated to reflux for 0.5 hour. After cooling, DCM and water (each 30 mL) were added. The resulting precipitate was isolated, washed with water and a small amount of methanol. The solid was dried *in vacuo* to afford 5.27 g (77%) of 4'-bromomethylbiphenyl-4-carboxylic acid.

20

HPLC-MS (Method C): m/z = 291 ($M+1$); R_t = 3.96 min.

25


5-(2-Trityl-2H-tetrazol-5-yl)-1H-indole (3.6 g, 8.4 mmol) was dissolved in DMF (100 mL). Under nitrogen, NaH (60 % suspension in mineral oil, 34 mmol) was added slowly. 4'-Bromomethylbiphenyl-4-carboxylic acid (2.7 g, 9.2 mmol) was added over 5 minutes and the

resulting slurry was heated at 40 °C for 16 hours. The mixture was poured into water (100mL) and the precipitate was isolated by filtration and treated with THF/6N HCl (9/1) (70 mL) at room temperature for 16 hours. The mixture was subsequently evaporated to dryness *in vacuo*, the residue was treated with water and the solid was isolated by filtration and washed thoroughly 3 times with DCM. The solid was dissolved in hot THF (400 mL) treated with activated carbon and filtered. The filtrate was evaporated *in vacuo* to dryness. This afforded 1.6 g (50%) of the title compound.

HPLC-MS (Method C): m/z = 396 (M+1); R_t = 3.51 min.

Example 810 (General procedure (K)).

5-(2*H*-Tetrazol-5-yl)-1*H*-indole




5-(2*H*-Tetrazol-5-yl)-1*H*-indole was prepared from 5-cyanoindole according to the method described in example 594.

HPLC-MS (Method C): m/z : 186 (M+1); R_t = 1.68 min.

Example 811 (General procedure (K)).

1-Benzyl-4-(2*H*-tetrazol-5-yl)-1*H*-indole




1-Benzyl-1*H*-indole-4-carbonitrile was prepared from 4-cyanoindole according to the method described in example 806.

HPLC-MS (Method C): m/z : 233 (M+1); R_t = 4.24 min.

1-Benzyl-4-(2*H*-tetrazol-5-yl)-1*H*-indole was prepared from 1-benzyl-1*H*-indole-4-carbonitrile according to the method described in example 594.


HPLC-MS (Method C): m/z: 276 (M+1); Rt = 3.44 min.

General procedure (L) for preparation of compounds of general formula I₁₁:



wherein T is as defined above and


Pol- is a polystyrene resin loaded with a 2-chlorotrityl linker, graphically shown below:



This general procedure (L) is further illustrated by the following example:

Example 812 (General procedure (L)).

5-(2H-Tetrazol-5-yl)-1-[3-(trifluoromethyl)benzyl]-1H-indole



2-Chlorotritylchloride resin (100 mg, 0.114 mmol active chloride) was swelled in dichloromethane (2 mL) for 30 min. The solvent was drained, and a solution of 5-(2H-


tetrazol-5-yl)-1*H*-indole (example 810) (63 mg, 0.34 mmol) in a mixture of *N,N*-dimethylformamide, dichloromethane and *N,N*-di(2-propyl)ethylamine (DIPEA) (5:5:2) (1.1 mL) was added. The reaction mixture was shaken at room temperature for 20 hours. The solvent was removed by filtration, and the resin was washed consecutively with *N,N*-dimethylformamide (2 x 4 mL), dichloromethane (6 x 4 mL) and methyl sulfoxide (2 x 4 mL). Methyl sulfoxide (1 mL) was added, followed by the addition of a solution of lithium bis(trimethylsilyl)amide in tetrahydrofuran (1.0 M, 0.57 mL, 0.57 mmol). The mixture was shaken for 30 min at room temperature, before 3-(trifluoromethyl)benzyl bromide (273 mg, 1.14 mmol) was added as a solution in methyl sulfoxide (0.2 mL). The reaction mixture was shaken for 20 hours at room temperature. The drained resin was washed consecutively with methyl sulfoxide (2 x 4 mL), dichloromethane (2 x 4 mL), methanol (2 x 4 mL), dichloromethane (2 x 4 mL) and tetrahydrofuran (4 mL). The resin was treated with a solution of hydrogen chloride in tetrahydrofuran, ethyl ether and ethanol = 8:1:1 (0.1 M, 3 mL) for 6 hours at room temperature. The resin was drained and the filtrate was concentrated *in vacuo*. The crude product was re-suspended in dichloromethane (1.5 mL) and concentrated three times to afford the title compound (35 mg). No further purification was necessary.

HPLC-MS (Method B): *m/z*: 344 (*M*+1); *R*_t = 4.35 min.

¹H-NMR (DMSO-*d*₆): δ 8.29 (1H, s), 7.80 (1H, dd), 7.72 (2H, m), 7.64 (2H, bs), 7.56 (1H, t), 7.48 (1H, d), 6.70 (1H, d), 5.62 (2H, s).


The compounds in the following examples were prepared in a similar fashion. Optionally, the compounds can be further purified by recrystallization or by chromatography.

Example 813 (General procedure (L)).
1-(4-Chlorobenzyl)-5-(2*H*-tetrazol-5-yl)-1*H*-indole



HPLC-MS (Method B): *m/z*: 310 (*M*+1); *R*_t = 4.11 min.


Example 814 (General procedure (L)).
1-(2-Chlorobenzyl)-5-(2*H*-tetrazol-5-yl)-1*H*-indole



HPLC-MS (Method B): m/z : 310 ($M+1$); R_t = 4.05 min.

Example 815 (General procedure (L)).


5 1-(4-Methoxybenzyl)-5-(2H-tetrazol-5-yl)-1H-indole



HPLC-MS (Method B): m/z : 306 ($M+1$); R_t = 3.68 min.

Example 816 (General procedure (L)).


10 1-(4-Methylbenzyl)-5-(2H-tetrazol-5-yl)-1H-indole



HPLC-MS (Method B): m/z : 290 ($M+1$); R_t = 3.98 min.

Example 817 (General procedure (L)).


15 5-(2H-Tetrazol-5-yl)-1-[4-(trifluoromethyl)benzyl]-1H-indole



HPLC-MS (Method B): m/z : 344 ($M+1$); R_t = 4.18 min.

Example 818 (General procedure (L)).

1-(3-Chlorobenzyl)-5-(2*H*-tetrazol-5-yl)-1*H*-indole




HPLC-MS (Method B): m/z : 310 ($M+1$); R_t = 4.01 min.

5

Example 819 (General procedure (L)).

1-(3-Methylbenzyl)-5-(2*H*-tetrazol-5-yl)-1*H*-indole




HPLC-MS (Method B): m/z : 290 ($M+1$); R_t = 3.98 min.

10

Example 820 (General procedure (L)).

1-(2,4-Dichlorobenzyl)-5-(2*H*-tetrazol-5-yl)-1*H*-indole




HPLC-MS (Method B): m/z : 344 ($M+1$); R_t = 4.41 min.

15

Example 821 (General procedure (L)).


1-(3-Methoxybenzyl)-5-(2*H*-tetrazol-5-yl)-1*H*-indole



HPLC-MS (Method B): m/z: 306 (M+1); Rt = 3.64 min.

Example 822 (General procedure (L)).


5 1-(4-Fluorobenzyl)-5-(2H-tetrazol-5-yl)-1H-indole



HPLC-MS (Method B): m/z: 294 (M+1); Rt = 3.71 min.

Example 823 (General procedure (L)).


10 1-(3-Fluorobenzyl)-5-(2H-tetrazol-5-yl)-1H-indole



HPLC-MS (Method B): m/z: 294 (M+1); Rt = 3.68 min.

Example 824 (General procedure (L)).


15 1-(2-Iodobenzyl)-5-(2H-tetrazol-5-yl)-1H-indole



HPLC-MS (Method B): m/z: 402 (M+1); Rt = 4.11 min.

Example 825 (General procedure (L)).


1-[(Naphthalen-2-yl)methyl]-5-(2*H*-tetrazol-5-yl)-1*H*-indole



5 HPLC-MS (Method B): *m/z*: 326 (*M*+1); *R*_t = 4.18 min.

Example 826 (General procedure (L)).


1-(3-Bromobenzyl)-5-(2*H*-tetrazol-5-yl)-1*H*-indole



10 HPLC-MS (Method B): *m/z*: 354 (*M*+1); *R*_t = 4.08 min.

Example 827 (General procedure (L)).

1-(4-Carboxybenzyl)-5-(2*H*-tetrazol-5-yl)-1*H*-indole




15 In this preparation, a larger excess of lithium bis(trimethylsilyl)amide in tetrahydrofuran (1.0 M, 1.7 mL, 1.7 mmol) was used.

HPLC-MS (Method B): *m/z*: 320 (*M*+1); *R*_t = 2.84 min.

Example 828 (General procedure (L)).

20 1-(3-Carboxybenzyl)-5-(2*H*-tetrazol-5-yl)-1*H*-indole




In this preparation, a larger excess of lithium bis(trimethylsilyl)amide in tetrahydrofuran (1.0 M, 1.7 mL, 1.7 mmol) was used.

HPLC-MS (Method B): m/z: 320 (M+1); Rt = 2.91 min.

5

Example 829 (General procedure (L)).


1-(2,4-Difluorobenzyl)-5-(2H-tetrazol-5-yl)-1H-indole



10 HPLC-MS (Method B): m/z: 312 (M+1); Rt = 3.78 min.

Example 830 (General procedure (L)).

1-(3,5-Difluorobenzyl)-5-(2H-tetrazol-5-yl)-1H-indole




15

HPLC-MS (Method B): m/z: 312 (M+1); Rt = 3.78 min.

Example 831 (General procedure (L)).


1-(3,4-Difluorobenzyl)-5-(2H-tetrazol-5-yl)-1H-indole



HPLC-MS (Method B): m/z : 312 ($M+1$); R_t = 3.81 min.

Example 832 (General procedure (L)).


5 1-[4-(2-Propyl)benzyl]-5-(2*H*-tetrazol-5-yl)-1*H*-indole



HPLC-MS (Method B): m/z : 318 ($M+1$); R_t = 4.61 min.

Example 833 (General procedure (L)).


10 1-(3,5-Dimethoxybenzyl)-5-(2*H*-tetrazol-5-yl)-1*H*-indole



HPLC-MS (Method B): m/z : 336 ($M+1$); R_t = 3.68 min.

Example 834 (General procedure (L)).


15 1-(2'-Cyanobiphenyl-4-ylmethyl)-5-(2*H*-tetrazol-5-yl)-1*H*-indole



HPLC-MS (Method B): m/z: 377 (M+1); Rt = 4.11 min.






















Example 835 (General procedure (L)).


5 1-(2-Methylbenzyl)-5-(2H-tetrazol-5-yl)-1H-indole




HPLC-MS (Method B): m/z: 290 (M+1); Rt = 3.98 min.

- 10 Further compounds of the invention that may be prepared according to general procedure (K) and/or (L) includes:


| | | |
|---|---|--|
| Example 836  | Example 837  | Example 838  |
| Example 839  | Example 840  | Example 841  |
| Example 842  | Example 843  | Example 844  |
| Example 845  | Example 846  | Example 847  |
| Example 848  | Example 849  | Example 850  |
| Example 851  | Example 852  | Example 853  |
| Example 854  | Example 855  | Example 856  |



The following compounds of the invention may be prepared eg. from 1-(4-bromobenzyl)-5-(2H-tetrazol-5-yl)-1H-indole (example 807) or from the analogue 1-(3-bromobenzyl)-5-(2H-tetrazol-5-yl)-1H-indole and aryl boronic acids *via* the Suzuki coupling reaction eg as described in Littke, Dai & Fu *J. Am. Chem. Soc.*, 2000, 122, 4020-8 (or references cited therein), or using the methodology described in general procedure (E), optionally changing the palladium catalyst to bis(tri-*tert*-butylphosphine)palladium (0).



10 **General procedure (M) for preparation of compounds of general formula I₁₂:**




wherein T is as defined above.

The general procedure (M) is further illustrated by the following example:

Example 865 (General procedure (M)).

5 1-Benzoyl-5-(2*H*-tetrazol-5-yl)-1*H*-indole



To a solution of 5-cyanoindole (1.0 g, 7.0 mmol) in dichloromethane (8 mL) was added 4-(dimethylamino)pyridine (0.171 g, 1.4 mmol), triethylamine (1.96 mL, 1.42 g, 14 mmol) and benzoyl chloride (0.89 mL, 1.08 g, 7.7 mmol). The resulting mixture was stirred for 18 hours at room temperature. The mixture was diluted with dichloromethane (80 mL) and washed consecutively with a saturated solution of sodium hydrogencarbonate (40 mL) and brine (40 mL). The organic phase was dried with magnesium sulfate (1 hour). Filtration and concentration furnished the crude material which was purified by flash chromatography on silica gel, eluting with ethyl acetate/heptanes = 2:3. 1-Benzoyl-1*H*-indole-5-carbonitrile was obtained as a solid.

HPLC-MS (Method C): *m/z*: 247 (*M*+1); *R*_t = 4.07 min.


1-Benzoyl-1*H*-indole-5-carbonitrile was transformed into 1-benzoyl-5-(2*H*-tetrazol-5-yl)-1*H*-indole by the method described in example 594.

HPLC (Method C): *R*_t = 1.68 min.

25 The compound in the following example was prepared by the same procedure.

Example 866 (General procedure (M)).

1-Benzoyl-4-(2*H*-tetrazol-5-yl)-1*H*-indole



1-Benzoyl-1*H*-indole-4-carbonitrile was prepared from 4-cyanoindole according to the method described in example 865.

HPLC-MS (Method C): m/z : 247 ($M+1$); R_t = 4.24 min.

5


1-Benzoyl-4-(2*H*-tetrazol-5-yl)-1*H*-indole was prepared from 1-benzoyl-1*H*-indole-4-carbonitrile according to the method described in example 594.

HPLC (Method C): R_t = 1.56 min.

10

Example 867 (General procedure (M))

(2-Fluoro-3-trifluoromethylphenyl)-[5-(2*H*-tetrazol-5-yl)-indol-1-yl]-methanone




HPLC-MS (Method B): m/z = 376 ($M+1$); R_t = 4.32 min.

15

Example 868 (General procedure (M))

(4-Methoxyphenyl)-[5-(2*H*-tetrazol-5-yl)-indol-1-yl]-methanone




HPLC-MS (Method B): m/z = 320 ($M+1$); R_t = 3.70 min.

20

Example 869 (General procedure (M))


(3-Nitrophenyl)-[5-(2H-tetrazol-5-yl)-indol-1-yl]-methanone

HPLC-MS (Method B): $m/z = 335$ (M+1); $R_t = 3.72$ min.

5

Example 870 (General procedure (M))


(4-Nitrophenyl)-[5-(2H-tetrazol-5-yl)-indol-1-yl]-methanone

HPLC-MS (Method B): $m/z = 335$ (M+1); $R_t = 3.71$ min.

10

Example 871 (General procedure (M))


Naphthalen-2-yl-[5-(2H-tetrazol-5-yl)-indol-1-yl]-methanone

HPLC-MS (Method C): $m/z = 340$ (M+1); $R_t = 4.25$ min.

15

Example 872 (General procedure (M))

(2,3-Difluorophenyl)-[5-(2H-tetrazol-5-yl)-indol-1-yl]-methanone




HPLC-MS (Method B: $m/z = 326$ ($M+1$); $R_t = 3.85$ min.

The following known and commercially available compounds do all bind to the His B10 Zn^{2+} site of the insulin hexamer:

5

Example 873


1-(4-Fluorophenyl)-5-(2H-tetrazol-5-yl)-1H-indole



10

Example 874


1-Amino-3-(2H-tetrazol-5-yl)benzene



15

Example 875

1-Amino-4-(2H-tetrazol-5-yl)benzene




20

A mixture of 4-aminobenzonitrile (10 g, 84.6 mmol), sodium azide (16.5 g, 254 mmol) and ammonium chloride (13.6 g, 254 mmol) in DMF was heated at 125 °C for 16 hours. The cooled mixture was filtered and the filtrate was concentrated *in vacuo*. The residue was added water (200 mL) and diethyl ether (200 mL) which resulted in crystallisation. The mixture was filtered and the solid was dried *in vacuo* at 40 °C for 16 hours to afford 5-(4-aminophenyl)-2H-tetrazole.


¹H NMR DMSO-*d*₆: δ = 5.7 (3H, bs), 6.69 (2H, d), 7.69 (2H, d).

HPLC-MS (Method C): m/z: 162 (M+1); Rt = 0,55 min.


Example 8761-Nitro-4-(2H-tetrazol-5-yl)benzene



Example 8771-Bromo-4-(2H-tetrazol-5-yl)benzene



General procedure (N) for solution phase preparation of amides of general formula I₁₃:




wherein Frag is any fragment carrying a carboxylic acid group, R is hydrogen, optionally substituted aryl or C₁₋₈-alkyl and R' is hydrogen or C₁₋₄-alkyl.

Frag-CO₂H may be prepared eg by general procedure (D) or by other similar procedures described herein, or may be commercially available.


The procedure is further illustrated in the following example 878:

Example 878 (General procedure (N))

N-(4-Chlorobenzyl)-2-[3-(2,4-dioxothiazolidin-5-ylidenemethyl)-1*H*-indol-1-yl]acetamide


- 5 [3-(2,4-Dioxothiazolidin-5-ylidenemethyl)indol-1-yl]acetic acid (example 478, 90.7 mg, 0.3 mmol) was dissolved in NMP (1 mL) and added to a mixture of 1-ethyl-3-(3-dimethylamino-propyl)carbodiimide, hydrochloride (86.4 mg, 0.45 mmol) and 1-hydroxybenzotriazol (68.8 mg, 0.45 mmol) in NMP (1 mL). The resulting mixture was shaken at RT for 2 h. 4-Chlorobenzylamine (51 mg, 0.36 mmol) and DIPEA (46.4 mg, 0.36 mmol) in NMP (1 mL)
- 10 were added to the mixture and the resulting mixture shaken at RT for 2 days. Subsequently ethyl acetate (10 mL) was added and the resulting mixture washed with 2x10 mL water followed by saturated ammonium chloride (5 mL). The organic phase was evaporated to dryness giving 75 mg (57%) of the title compound.
- 15 HPLC-MS (Method C): m/z: 426 (M+1); Rt. = 3.79 min.

Example 879 (General procedure (N))

N-(4-Chlorobenzyl)-4-[2-chloro-4-(2,4-dioxothiazolidin-5-ylidenemethyl)phenoxy]butyramide


- 20 HPLC-MS (Method A): m/z: 465 (M+1); Rt = 4.35 min.

Example 880 (General procedure (N))

N-(4-Chlorobenzyl)-4-[4-(2,4-dioxothiazolidin-5-ylidenemethyl)phenoxy]butyramide


- 25 HPLC-MS (Method A): m/z: 431 (M+1); Rt = 3.68 min.

Example 881 (General procedure (N))

2-[2-Bromo-4-(2,4-dioxothiazolidin-5-ylidenemethyl)phenoxy]-*N*-(4-chlorobenzyl)acetamideHPLC-MS (Method A): *m/z*: 483 (*M*+1); *R*_t = 4.06 min.


5

Example 882 (General procedure (N))

N-(4-Chlorobenzyl)-2-[3-(2,4-dioxothiazolidin-5-ylidenemethyl)phenoxy]acetamideHPLC-MS (Method A): *m/z*: 403 (*M*+1); *R*_t = 4.03 min.


10

Example 883 (General procedure (N))

N-(4-Chlorobenzyl)-3-[4-(2,4-dioxothiazolidin-5-ylidenemethyl)phenyl]acrylamideHPLC-MS (Method A): *m/z*: 399 (*M*+1); *R*_t = 3.82.


15

Example 884 (General procedure (N))

N-(4-Chlorobenzyl)-4-[3-(2,4-dioxothiazolidin-5-ylidenemethyl)phenoxy]butyramideHPLC-MS (Method A): *m/z*: 431 (*M*+1); *R*_t = 3.84 min.


20

Example 885 (General procedure (N))

4-[2-Bromo-4-(2,4-dioxothiazolidin-5-ylidenemethyl)phenoxy]-*N*-(4-chlorobenzyl)butyramideHPLC-MS (Method A): *m/z*: 511 (*M*+1); *R*_t = 4.05 min.

Example 886 (General procedure (N))


4-[2-Bromo-4-(4-oxo-2-thioxothiazolidin-5-ylidenemethyl)-phenoxy]-*N*-(4-chlorobenzyl)-butyramide



HPLC-MS (Method A): m/z : 527 ($M+1$); R_t = 4.77 min.

Example 887 (General procedure (N))


N-(4-Chlorobenzyl)-2-[4-(2,4-dioxothiazolidin-5-ylidenemethyl)naphthalen-1-yloxy]acetamide



HPLC-MS (Method C): m/z : 431 ($M+1$); R_t = 4.03 min.

Example 888 (General procedure (N))


N-(4-Chlorobenzyl)-3-[3-(2,4-dioxothiazolidin-5-ylidenemethyl)-1*H*-indol-1-yl]propionamide



HPLC-MS (Method C): m/z : 440 ($M+1$); R_t = 3.57 min.

Example 889 (General procedure (N))


N-(4-Chlorobenzyl)-4-[4-(2,4-dioxothiazolidin-5-ylidenemethyl)naphthalen-1-yloxy]butyramide



HPLC-MS (Method C): m/z : 481 ($M+1$); R_t = 4.08 min.

Example 890 (General procedure (N))


4-[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)-naphthalen-1-yloxy]-N-hexylbutyramide



5 HPLC-MS (Method C): m/z: 441 (M+1); Rt = 4.31 min.

Example 891 (General Procedure (N))

4-([3-(2,4-Dioxothiazolidin-5-ylidenemethyl)indole-7-carbonyl]amino)methyl)benzoic acid methyl ester




10

HPLC-MS (Method C): m/z: 436 (M+1); Rt.= 3.55 min.

Example 892 (General procedure (N))


N-(4-Chlorobenzyl)-4-[3-(2H-tetrazol-5-yl)carbazol-9-ylmethyl]benzamide



15 HPLC-MS (Method C): m/z:493 (M+1); Rt = 4.19 min.

Example 893 (General procedure (N))


N-(4-Chlorobenzyl)-3-[3-(2H-tetrazol-5-yl)carbazol-9-ylmethyl]benzamide



HPLC-MS (Method C): m/z : 493 ($M+1$); R_t = 4.20 min.

Example 894 (General Procedure (N))


- 5 *N*-(4-Chlorobenzyl)-3-methyl-4-[3-(2*H*-tetrazol-5-yl)-carbazol-9-ylmethyl]benzamide



HPLC-MS (Method C): m/z : 507 ($M+1$); R_t = 4.37 min.

Example 895 (General procedure (N))


- 10 5-{2-[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)-naphthalen-1-yloxy]-acetylamino}-isophthalic acid dimethyl ester



HPLC-MS (Method C): m/z = 521 ($M+1$); R_t = 4.57 min.

- 15 Example 896 (General procedure (N))

5-{2-[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)-naphthalen-1-yloxy]-acetylamino}-isophthalic acid




HPLC-MS (Method C): m/z = 515 ($M+23$); R_t = 3.09 min.

5

Example 897 (General procedure (N))


5-(3-{2-[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)-naphthalen-1-yloxy]-ethyl}-ureido)-isophthalic acid monomethyl ester



10 HPLC-MS (Method C): m/z = 536 ($M+1$); R_t = 3,58 min.

Example 898 (General Procedure (N)).

2-[4-[3-(2H-Tetrazol-5-yl)carbazol-9-ylmethyl]benzoylamino]succinic acid dimethyl ester



15 4-[3-(1H-Tetrazol-5-yl)carbazol-9-ylmethyl]benzoic acid (2.00 g, 5.41 mmol), 1-hydroxybenzotriazole (1.46 g, 10.8 mmol) and N,N-di(2-propyl)ethylamine (4.72 mL, 3.50 g, 27.1 mmol) were dissolved in dry N,N-dimethylformamide (60 mL). The mixture was cooled in an ice-water bath, and 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (1.45 g, 7.56 mmol) and (S)-aminosuccinic acid dimethyl ester hydrochloride (1.28 g, 6.48


mmol) were added. The cooling was discontinued, and the reaction mixture was stirred at room temperature for 18 hours before it was poured into hydrochloric acid (0.1 N, 600 mL). The solid was collected by filtration and washed with water (2 X 25 mL) to furnish the title compound.

5 HPLC-MS (Method C): m/z: 513 (M+1); Rt = 3.65 min.

¹H-NMR (DMSO-d₆): δ 8.90 (1H, d), 8.86 (1H, d), 8.29 (1H, d), 8.11 (1H, dd), 7.87 (1H, d), 7.75 (2H, d), 7.69 (1H, d), 7.51 (1H, t), 7.32 (1H, t), 7.28 (2H, d), 5.82 (2H, s), 4.79 (1H, m), 3.61 (3H, s), 3.58 (3H, s), 2.92 (1H, dd), 2.78 (1H, dd).

10 Example 899 (General Procedure (N)).

2-{4-[3-(2H-Tetrazol-5-yl)carbazol-9-ylmethyl]benzoylamino}succinic acid



2-{4-[3-(2H-Tetrazol-5-yl)carbazol-9-ylmethyl]benzoylamino}succinic acid dimethyl ester (1.20 g, 2.34 mmol) was dissolved in tetrahydrofuran (30 mL). Aqueous sodium hydroxide (1 N, 14 mL) was added, and the resulting mixture was stirred at room temperature for 18 hours. The reaction mixture was poured into hydrochloric acid (0.1 N, 500 mL). The solid was collected by filtration and washed with water (2 X 25 mL) and diethyl ether (2 X 25 mL) to furnish the title compound.

HPLC-MS (Method C): m/z: 485 (M+1); Rt = 2.94 min.


20 ¹H-NMR (DMSO-d₆): δ 12.44 (2H, s (br)), 8.90 (1H, d), 8.68 (1H, d), 8.29 (1H, d), 8.11 (1H, dd), 7.87 (1H, d), 7.75 (2H, d), 7.68 (1H, d), 7.52 (1H, t), 7.32 (1H, t), 7.27 (2H, d), 5.82 (2H, s), 4.70 (1H, m), 2.81 (1H, dd), 2.65 (1H, dd).

The compounds in the following examples were prepared in a similar fashion.

25

Example 900 (General procedure (N))


2-{4-[3-(2H-Tetrazol-5-yl)-carbazol-9-ylmethyl]-benzoylamino}-succinic acid dimethyl ester



HPLC-MS (Method C): m/z = 513 ($M+1$); R_t = 3.65min.

Example 901 (General procedure (N))


- 5 2-{4-[3-(2H-Tetrazol-5-yl)carbazol-9-ylmethyl]benzoylamino}pentanedioic acid dimethyl ester



HPLC-MS (Method C): m/z = 527 ($M+1$); R_t = 3.57min.

Example 902 (General procedure (N))


- 10 (Methoxycarbonylmethyl-{4-[3-(2H-tetrazol-5-yl)-carbazol-9-ylmethyl]-benzoyl}-amino)-acetic acid methyl ester



HPLC-MS (Method C): m/z = 513 ($M+1$); R_t = 3,55min.

- 15 Example 903 (General procedure (N))


2-{4-[3-(2H-Tetrazol-5-yl)carbazol-9-ylmethyl]benzoylamino}pentanedioic acid



HPLC-MS (Method C): m/z = 499 ($M+1$); R_t = 2.87 min.

Example 904 (General procedure (N))


- 5 (Ethoxycarbonylmethyl-4-[3-(2H-tetrazol-5-yl)-carbazol-9-ylmethyl]-benzoyl)-amino)-acetic acid ethyl ester



HPLC-MS (Method C): m/z = 541 ($M+1$); R_t = 3.91 min.

- 10 Example 905 (General procedure (N))

3-(3-{4-[4-(2,4-Dioxo-thiazolidin-5-ylidenemethyl)-naphthalen-1-yloxy]-butyrylamino}-propylamino)-hexanedioic acid dimethyl ester




HPLC-MS (Method C: m/z = 585 ($M+1$); R_t = 2,81 min.

15

Example 906 (General procedure (N))


3-(3-{4-[4-(2,4-Dioxo-thiazolidin-5-ylidenemethyl)-naphthalen-1-yloxy]-butyrylamino}-propylamino)-hexanedioic acid



HPLC-MS (Method C): m/z = 554 (M-3); R_t = 3.19 min.

Example 907 (General procedure (N))


5 (Carboxymethyl-{4-[3-(2H-tetrazol-5-yl)-carbazol-9-ylmethyl]-benzoyl}-amino)-acetic acid



HPLC-MS (Method C): m/z = 485 (M+1); R_t = 3.04 min.

Example 908 (General procedure (N))


10 4-(3-{4-[4-(2,4-Dioxothiazolidin-5-ylidenemethyl)-naphthalen-1-yloxy]-butyrylamino}-propylamino)-cyclohexane-1,3-dicarboxylic acid dimethyl ester



HPLC-MS (Method C): m/z = 612 (M+1); Rt = 3,24 min.

15 Example 909 (General procedure (N))


2-{3-[3-(2H-Tetrazol-5-yl)carbazol-9-ylmethyl]benzoylamino}pentanedioic acid dimethyl ester



HPLC-MS (Method C): m/z = 527 (M+1); R_t = 3.65min.

Example 910 (General procedure (N))


2-{3-[3-(2H-Tetrazol-5-yl)carbazol-9-ylmethyl]benzoylamino}pentanedioic acid dimethyl ester

HPLC-MS (Method C): m/z = 527 ($M+1$); R_t = 3.65min.

5

Example 911 (General procedure (N))


2-{3-[3-(2H-Tetrazol-5-yl)carbazol-9-ylmethyl]benzoylamino}pentanedioic acid dimethyl ester

HPLC-MS (Method C): m/z = 527 ($M+1$); R_t = 3.65min.

10

Example 912 (General procedure (N))


2-{3-[3-(2H-Tetrazol-5-yl)carbazol-9-ylmethyl]benzoylamino}pentanedioic acid

HPLC-MS (Method C): m/z = 499 ($M+1$); R_t = 3.00 min.

15

Example 913 (General procedure (N))

(Methoxycarbonylmethyl-{3-[3-(2H-tetrazol-5-yl)carbazol-9-ylmethyl]benzoyl}amino)acetic acid methyl ester




$^1\text{H-NMR}$ (DMSO-d_6): δ 8.88 (1H, d), 8.29 (1H, d), 8.10 (1H, dd), 7.85 (1H, d), 7.67 (1H, d), 7.52 (1H, t), 7.39 (1H, t), 7.30 (2H, m), 7.17 (2H, m), 5.79 (2H, s), 4.17 (2H, s), 4.02 (2H, s), 3.62 (3H, s), 3.49 (3H, s).

5

Example 914 (General procedure (N))

2-{3-[3-(2H-Tetrazol-5-yl)carbazol-9-ylmethyl]benzoylamino}succinic acid dimethyl ester




HPLC-MS (Method C): m/z = 513 ($M+1$); R_t = 3.70 min.

10

Example 915 (General procedure (N))

2-{3-[3-(2H-Tetrazol-5-yl)-carbazol-9-ylmethyl]-benzoylamino}-succinic acid




HPLC-MS (Method C): m/z = 485 ($M+1$); R_t = 2.96 min.

15

Example 916 (General procedure (N))


(Carboxymethyl-{3-[3-(2H-tetrazol-5-yl)carbazol-9-ylmethyl]benzoyl}amino)acetic acid



HPLC-MS (Method C): m/z = 485 ($M+1$); R_t = 2.87 min.

Example 917 (General procedure (N))

- 5 4-(4-(3-Carboxy-propylcarbamoyl)-4-[3-(2H-tetrazol-5-yl)carbazol-9-ylmethyl]-benzoylamino)-butyrylamino)-butyric acid




- 10 The title compound was prepared by coupling of (S)-2-[4-[3-(2H-tetrazol-5-yl)carbazol-9-ylmethyl]benzoylamino]pentanedioic acid bis-(2,5-dioxopyrrolidin-1-yl) ester (prepared from (S)-2-[4-[3-(2H-tetrazol-5-yl)carbazol-9-ylmethyl]benzoylamino]pentanedioic acid by essentially the same procedure as described for the synthesis of 4-[3-(2H-tetrazol-5-yl)carbazol-9-ylmethyl]benzoic acid 2,5-dioxopyrrolidin-1-yl ester) with 4-aminobutyric acid according to the procedure described for the preparation of 4-[4-[3-(2H-tetrazol-5-yl)carbazol-9-ylmethyl]benzoylamino]butyric acid .

15

HPLC-MS (Method C): m/z : 669 ($M+1$); R_t = 2.84 min.

Example 918 (General procedure (N))


[2-(2-[4-[3-(2H-Tetrazol-5-yl)-carbazol-9-ylmethyl]benzoylamino]ethoxy)ethoxy]acetic acid



HPLC-MS (Method C): m/z : 515 (M+1); R_t = 3.10 min.

Example 919 (General procedure (N))


- 5 2-{4-[3-(2H-Tetrazol-5-yl)-carbazol-9-ylmethyl]-benzoylamino}-pentanedioic acid di-tert-butyl ester



HPLC-MS (Method C): m/z = 611 (M+1); R_t = 4.64 min.

- 10 Example 920 (General Procedure (N)).

4-{4-[3-(2H-Tetrazol-5-yl)carbazol-9-ylmethyl]benzoylamino}butyric Acid




HPLC-MS (Method C): m/z : 455 (M+1); R_t = 3.13 min.

- 15 Example 921 (General Procedure (N)).

[2-(2-{4-[3-(2H-Tetrazol-5-yl)carbazol-9-ylmethyl]benzoylamino}ethoxy)ethoxy]acetic acid

340




The title compound was prepared by coupling of 4-[3-(2H-tetrazol-5-yl)carbazol-9-ylmethyl]benzoic acid 2,5-dioxopyrrolidin-1-yl ester with [2-(2-aminoethoxy)ethoxy]acetic acid (prepared from [2-[2-(Fmoc-amino)ethoxy]ethoxy]acetic acid by treatment with PS-Trisamine resin in DMF).

HPLC-MS (Method C): m/z: 515 (M+1); Rt = 3.10 min.

The commercially available compounds in the following examples do all bind to the HisB10 Zn²⁺ site:


Example 922

1-(4-Bromo-3-methylphenyl)-1,4-dihydrotetrazole-5-thione




Example 923

1-(4-Iodophenyl)-1,4-dihydrotetrazole-5-thione




Example 924

1-(2,4,5-Trichlorophenyl)-1H-tetrazole-5-thiol




Example 925

1-(2,6-Dimethylphenyl)-1,4-dihydrotetrazole-5-thione




5 Example 926

1-(2,4,6-Trimethylphenyl)-1,4-dihydrotetrazole-5-thione




Example 927

10 1-(4-Dimethylaminophenyl)-1H-tetrazole-5-thiol



Example 928


1-(3,4-Dichlorophenyl)-1,4-dihydro-1H-tetrazole-5-thione



15

Example 929


1-(4-Propylphenyl)-1,4-dihydro-1H-tetrazole-5-thione



20


Example 930

1-(3-Chlorophenyl)-1,4-dihydro-1H-tetrazole-5-thione




5 Example 931

1-(2-Fluorophenyl)-1,4-dihydro-1H-tetrazole-5-thione




Example 932

10 1-(2,4-Dichlorophenyl)-1,4-dihydro-1H-tetrazole-5-thione



Example 933


1-(4-Trifluoromethoxyphenyl)-1,4-dihydro-1H-tetrazole-5-thione



15

Example 934


N-[4-(5-Mercaptotetrazol-1-yl)-phenyl]-acetamide



20


Example 935

1-(4-Chlorophenyl)-1,4-dihydrotetrazole-5-thione




5 Example 936

1-(4-Methoxyphenyl)-1,4-dihydrotetrazole-5-thione




Example 937

10 1-(3-Fluoro-4-pyrrolidin-1-ylphenyl)-1,4-dihydrotetrazole-5-thione



Example 938


N-[3-(5-Mercaptotetrazol-1-yl)phenyl]acetamide




15

Example 939

1-(4-Hydroxyphenyl)-5-mercaptotetrazole



Example 940




5

Preparation of 1-aryl-1,4-dihydro-1H-tetrazole-5-thiones (or the tautomeric 1-aryltetrazole-5-thiols) is described in the literature (eg. by Kauer & Sheppard, *J. Org. Chem.*, **32**, 3580-92 (1967)) and is generally performed eg. by reaction of aryl-isothiocyanates with sodium azide followed by acidification

1-Aryl-1,4-dihydro-1H-tetrazole-5-thiones with a carboxylic acid tethered to the aryl group may be prepared as shown in the following scheme:

15










Step 1 is a phenol alkylation and is very similar to steps 1 and 2 of general procedure (D) and may also be prepared similarly as described in example 481.

Step 2 is a reduction of the nitro group. SnCl_2 , H_2 over Pd/C and many other procedures known to those skilled in the art may be utilised.

Step 3 is formation of an arylisothiocyanate from the corresponding aniline. As reagents CS_2 , CSCl_2 , or other reagents known to those skilled in the art, may be utilised.

Step 4 is a conversion to mercaptotetrazole as described above.


Compounds of the invention include:

| | |
|--|---|
| <p>Example 941</p>  | <p>Example 942</p>  |
| <p>Example 943</p>  | <p>Example 944</p>  |
| <p>Example 945</p>  | <p>Example 946</p>  |
| <p>Example 947</p>  | |

10

Example 948

4-(4-Hydroxyphenyl)-1H-[1,2,3]triazole-5-carbonitrile



Phenylsulphonyl acetonitrile (2.0 g, 11.04 mmol) was mixed with 4-hydroxybenzaldehyde (1.35 g, 11.04 mmol) in DMF (10 mL) and toluene (20 mL). The mixture was refluxed for 3 hours and subsequently evaporated to dryness *in vacuo*. The residue was treated with diethyl ether and toluene. The solid formed was filtered to afford 2.08 g (66%) of 2-benzenesulfonyl-3-(4-hydroxyphenyl)acrylonitrile.


HPLC-MS (Method C): m/z: 286 (M+1); Rt. = 3.56 min.

A mixture of 2-benzenesulfonyl-3-(4-hydroxyphenyl)acrylonitrile (2.08 g, 7.3 mmol) and sodium azide (0.47g, 7.3 mmol) in DMF (50 mL) was heated at reflux temperature 2 hours. After cooling, the mixture was poured on ice. The mixture was evaporated *in vacuo* to almost dryness and toluene was added. After filtration, the organic phase was evaporated *in vacuo*. The residue was purified by silicagel chromatography eluting with a mixture of ethyl acetate and heptane (1:2). This afforded 1.2 g (76%) of the title compound.

¹H NMR (DMSO-*d*₆): 10.2 (broad, 1H); 7.74 (d, 2H); 6.99 (d, 2H); 3.6-3.2 (broad, 1H).

HPLC-MS (Method C) m/z: = 187 (M+1); Rt. = 1.93 min

General procedure (O) for preparation of compounds of general formula I₁₄:



wherein

AA is as defined above,

Steps 1 and 2 are described in the literature (eg Beck & Günther, *Chem. Ber.*, **106**, 2758-66 (1973))


Step 1 is a Knoevenagel condensation of the aldehyde AA-CHO with phenylsulfonyl-acetonitrile and step 2 is a reaction of the vinylsulfonyl compound obtained in step 1 with sodium azide. This reaction is usually performed in DMF at 90 – 110 °C.

5

This general procedure is further illustrated in the following example 949:

Example 949 (General Procedure (O))

[4-(5-Cyano-1H-[1,2,3]triazol-4-yl)phenoxy]acetic acid



10

Phenylsulfonylacetonitrile (0.1 g, 0.55 mmol) was mixed with 4-formylphenoxyacetic acid (0.099 g, 0.55 mmol) in DMF (3 mL) and heated to 110 °C for 3 h and subsequently cooled to RT. Sodium azide (0.036 g, 0.55 mmol) was added and the resulting mixture was heated to 110 °C for 3 h and cooled to RT. The mixture was poured into water (20 mL) and centrifuged. The supernatant was discarded, ethanol (5 mL) was added and the mixture was centrifuged again. After discarding the supernatant, the residue was dried *in vacuo* to afford 50 mg (37%) of [4-(5-Cyano-1H-[1,2,3]triazol-4-yl)phenoxy]acetic acid.


15

HPLC-MS (Method C): m/z: 245 (M+1) Rt. 2.19 min.

20

Example 950 (General Procedure (O))

5-(Naphthalen-1-yl)-3H-[1,2,3]triazole-4-carbonitrile




HPLC-MS (Method C): m/z: 221 (M+1); Rt. 3.43 min.

25

Example 951 (General Procedure (O))

5-(Naphthalen-2-yl)-3H-[1,2,3]triazole-4-carbonitrile




HPLC-MS (Method C): m/z: 221 (M+1); Rt = 3.66 min.

5

Example 952 (General procedure (O))

4-[3-(5-Cyano-[1,2,3]triazol-4-yl)-1,4-dimethylcarbazol-9-ylmethyl]-benzoic acid



HPLC-MS (Method C): m/z = 422 (M+1); Rt = 3.85 min.

10


Preparation of intermediary aldehyde:

1,4 Dimethylcarbazol-3-carbaldehyde (0.68 g, 3.08 mmol) was dissolved in dry DMF (15 mL), NaH (diethyl ether washed) (0.162 g, 6.7 mol) was slowly added under nitrogen and the mixture was stirred for 1 hour at room temperature. 4-Bromomethylbenzoic acid (0.73 g, 3.4 mmol) was slowly added and the resulting slurry was heated to 40 °C for 16 hours. Water (5 mL) and hydrochloric acid (6N, 3 mL) were added. After stirring for 20 min at room temperature, the precipitate was filtered off and washed twice with acetone to afford after drying 0.38 g (34%) of 4-(3-formyl-1,4-dimethylcarbazol-9-ylmethyl)benzoic acid.

20 HPLC-MS (Method C) : m/z = 358 (M+1), RT. = 4.15 min.

Example 953 (General Procedure (O))


5-(Anthracen-9-yl)-3H-[1,2,3]triazole-4-carbonitrile



HPLC-MS (Method C): m/z : 271 ($M+1$); R_t = 3.87 min.

Example 954 (General Procedure (O))


- 5 5-(4-Methoxynaphthalen-1-yl)-3H-[1,2,3]triazole-4-carbonitrile



HPLC-MS (Method C): m/z : 251 ($M+1$); R_t = 3.57 min.

Example 955 (General Procedure (O))


- 10 5-(1,4-Dimethyl-9H-carbazol-3-yl)-3H-[1,2,3]triazole-4-carbonitrile



HPLC-MS (Method C): m/z : 288 ($M+1$); R_t = 3.67 min.

Example 956 (General procedure (O))


- 15 5-(4'-Methoxybiphenyl-4-yl)-3H-[1,2,3]triazole-4-carbonitrile



HPLC-MS (Method C): $m/z = 277$ ($M+1$); $R_t = 3.60$ min.

Example 957 (General procedure (O))


5 5-(4-Styrylphenyl)-3H-[1,2,3]triazole-4-carbonitrile



HPLC-MS (Method C): $m/z = 273$ ($M+1$); $R_t = 4.12$ min.

Example 958 (General procedure (O))


10 5-(2,6-Dichloro-4-dibenzylaminophenyl)-3H-[1,2,3]triazole-4-carbonitrile



HPLC-MS (Method C): $m/z = 434$ ($M+1$); $R_t = 4.64$ min.

Example 959 (General procedure (O))


5-(1-Bromonaphthalen-2-yl)-3H-[1,2,3]triazole-4-carbonitrile



- 5 HPLC-MS (Method C: $m/z = 300$ (M+1); $R_t = 3.79$ min.

Example 960


4-(4-Bromophenyl)-1H-[1,2,3]triazole-5-carbonitrile



- 10 This compound is commercially available (MENAI).

Example 961


N-[4-(5-Cyano-1H-[1,2,3]triazol-4-yl)-phenyl]-acetamide



- 15 This compound is commercially available (MENAI).

Example 962 (General procedure (O))

5-(4'-Chlorobiphenyl-4-yl)-3H-[1,2,3]triazole-4-carbonitrile




HPLC-MS (Method C): m/z = 281 (M+1); R_t = 4.22 min.

- 5 The compounds in the following examples are commercially available and may be prepared using a similar methodology:

Example 963


4-(4-Trifluoromethoxyphenyl)-1H-[1,2,3]triazole-5-carbonitrile



10

Example 964


4-Benzo[1,3]dioxol-5-yl-1H-[1,2,3]triazole-5-carbonitrile



15


Example 965

4-(3-Trifluoromethylphenyl)-1H-[1,2,3]triazole-5-carbonitrile



Example 966


4-Pyridin-3-yl-1H-[1,2,3]triazole-5-carbonitrile



5

Example 967


4-(2,6-Dichlorophenyl)-1H-[1,2,3]triazole-5-carbonitrile




10

Example 968


4-Thiophen-2-yl-1H-[1,2,3]triazole-5-carbonitrile




15 Example 969

3,5-Dimethylisoxazole-4-carboxylic acid 4-(5-cyano-1*H*-[1,2,3]triazol-4-yl)phenyl ester

Example 970


5 3,3-Dimethyl-butylric acid 4-(5-cyano-1*H*-[1,2,3]triazol-4-yl)phenyl ester

Example 971


4-Methyl-[1,2,3]thiadiazole-5-carboxylic acid 4-(5-cyano-1*H*-[1,2,3]triazol-4-yl)phenyl ester

10

Example 972


4-Chlorobenzoic acid 4-(5-cyano-1*H*-[1,2,3]triazol-4-yl)phenyl ester

Example 973


4-(3-Phenoxyphenyl)-1*H*-[1,2,3]triazole-5-carbonitrile

5

Example 974

4-(5-Bromo-2-methoxyphenyl)-1*H*-[1,2,3]triazole-5-carbonitrile

10 Example 975

4-(2-Chloro-6-fluorophenyl)-1*H*-[1,2,3]triazole-5-carbonitrile

The following cyanotriazoles are also compounds of the invention:

15

4-(2-Chloro-6-fluorophenyl)-1*H*-[1,2,3]triazole-5-carbonitrile.Terephthalic acid mono[4-(5-cyano-1*H*-[1,2,3]triazol-4-yl)phenyl] ester.*N*- [4-(5-cyano-1*H*-[1,2,3]triazol-4-yl)-phenyl]terephthalamide4-(4-Octyloxyphenyl)-1*H*-[1,2,3]triazole-5-carbonitrile20 4-(4-Styrylphenyl)-1*H*-[1,2,3]triazole-5-carbonitrile.4-(4'-Trifluoromethylbiphenyl-4-yl)-1*H*-[1,2,3]triazole-5-carbonitrile.4-(4'-Chlorobiphenyl-4-yl)-1*H*-[1,2,3]triazole-5-carbonitrile.4-(4'-Methoxybiphenyl-4-yl)-1*H*-[1,2,3]triazole-5-carbonitrile.4-(1-Naphthyl)-1*H*-[1,2,3]triazole-5-carbonitrile.

4-(9-Anthranyl)-1*H*-[1,2,3]triazole-5-carbonitrile.


4-(4-Methoxy-1-naphthyl)-1*H*-[1,2,3]triazole-5-carbonitrile.

4-(4-Aminophenyl)-1*H*-[1,2,3]triazole-5-carbonitrile.

4-(2-Naphthyl)-1*H*-[1,2,3]triazole-5-carbonitrile.

5

General procedure (P) for preparation of compounds of general formula I₁₅:



10 wherein

n is 1 or 3-20,

AA is as defined above,

R'' is a standard carboxylic acid protecting group, such as C_1 - C_6 -alkyl or benzyl and Lea is a leaving group, such as chloro, bromo, iodo, methanesulfonyloxy, toluenesulfonyloxy or the like.

15

This procedure is very similar to general procedure (D), steps 1 and 2 are identical.

Steps 3 and 4 are described in the literature (eg Beck & Günther, *Chem. Ber.*, **106**, 2758-66 (1973))


20

Step 3 is a Knoevenagel condensation of the aldehyde obtained in step 2 with phenylsulfonylacetonitrile and step 4 is a reaction of the vinylsulfonyl compound obtained in step 3 with sodium azide. This reaction is usually performed in DMF at 90 – 110 °C.

This General procedure (P) is further illustrated in the following two examples

Example 976 (General procedure (P))

5 5-[6-(5-Cyano-1H-[1,2,3]triazol-4-yl)-naphthalen-2-yloxy]-pentanoic acid ethyl ester



6-Hydroxynaphthalene-2-carbaldehyde (Syncom BV. NL, 15.5 g, 90 mmol) and K_2CO_3 (62.2 g, 450 mmol) were mixed in DMF (300mL) and stirred at room temperature for 1hour. Ethyl 5-bromovalerate (21.65 g, 103.5 mmol) was added and the mixture was stirred at room
10 temperature for 16 hours. Activated carbon was added and the mixture was filtered. The filtrate was evaporated to dryness *in vacuo* to afford 28.4 g of crude 5-(6-formylnaphthalen-2-yloxy)pentanoic acid ethyl ester, which was used without further purification.

HPLC-MS (Method C): m/z = 301 ($M+1$); R_t = 4.39 min.

15

5-(6-Formylnaphthalen-2-yloxy)pentanoic acid ethyl ester (28.4 g, 94.5 mmol), phenylsulfonylacetonitrile (20.6 g, 113.5 mmol), and piperidine (0.94 mL) were dissolved in DMF (200 mL) and the mixture was heated at 50 °C for 16 hours. The resulting mixture was evaporated to dryness *in vacuo* and the residue was dried for 16 hours at 40 °C *in vacuo*.
20 The solid was recrystallised from 2-propanol (800 mL) and dried again as described above. This afforded 35 g (80%) of 5-[6-(2-benzenesulfonyl-2-cyanovinyl)naphthalen-2-yloxy]pentanoic acid ethyl ester.

HPLC-MS (Method C): m/z = 486 ($M+23$); R_t = 5.09 min.

25

5-[6-(2-Benzenesulfonyl-2-cyanovinyl)naphthalen-2-yloxy]pentanoic acid ethyl ester (35 g, 74.6 mmol) and sodium azide (4.9 g, 75.6 mmol) were dissolved in DMF (100 mL) and stirred


for 16 hours at 50 °C. The mixture was evaporated to dryness *in vacuo*, redissolved in THF / ethanol and a small amount of precipitate was filtered off. The resulting filtrate was poured into water (2.5 L). Filtration afforded after drying 24.5 g (88%) of 5-[6-(5-cyano-1H-[1,2,3]triazol-4-yl)naphthalen-2-yloxy]pentanoic acid ethyl ester (24.5 g, 88%).

5

HPLC-MS (Method C): m/z = 365 (M+1); R_t = 4.36 min.

Example 977 (General procedure (B))

5-[6-(5-Cyano-1H-[1,2,3]triazol-4-yl)-naphthalen-2-yloxy]-pentanoic acid



5-[6-(5-Cyano-1H-[1,2,3]triazol-4-yl)naphthalen-2-yloxy]pentanoic acid ethyl ester (24.5 g, 67.4 mmol) was dissolved in THF (150 mL) and mixed with sodium hydroxide (8.1 g, 202 mmol) dissolved in water (50 mL). The mixture was stirred for 2 days and the volatiles were evaporated *in vacuo*. The resulting aqueous solution was poured into a mixture of water (1 L) and hydrochloric acid (1N, 250 mL). The solid was isolated by filtration, dissolved in sodium hydroxide (1N, 200 mL), and the solution was washed with DCM and then ethyl acetate, the aqueous layer was acidified with hydrochloric acid (12N). The precipitate was isolated by filtration, dissolved in THF / diethyl ether, the solution was treated with $MgSO_4$ and activated carbon, filtrated and evaporated *in vacuo* to almost dryness followed by precipitation by addition of pentane (1L). This afforded after drying *in vacuo* 17.2 g (76%) of the title compound.


15

20

HPLC-MS (Method C): m/z = 337 (M+1); R_t = 3.49 min.

25 Example 978 (General procedure (P))


6-[6-(5-Cyano-1H-[1,2,3]triazol-4-yl)naphthalen-2-yloxy]hexanoic acid



HPLC-MS (Method C): $m/z = 351$ ($M+1$); $R_t = 3.68$ min.

Example 979 (General procedure (P))


- 5 11-[6-(5-Cyano-1H-[1,2,3]triazol-4-yl)-naphthalen-2-yloxy]-undecanoic acid



HPLC-MS (Method C): $m/z = 443$ ($M+23$); $R_t = 4.92$ min.

Example 980 (General procedure (P))


- 10 2-{3-[6-(5-Cyano-1H-[1,2,3]triazol-4-yl)-naphthalen-2-yloxy]-propyl}-malonic acid diethyl ester



HPLC-MS (Method C): $m/z = 465$ ($M+1$); $R_t = 4.95$ min.

Example 981 (General procedure (P))


- 15 2-{5-[6-(5-Cyano-1H-[1,2,3]triazol-4-yl)-naphthalen-2-yloxy]-pentyl}-malonic acid diethyl ester



HPLC-MS (Method C): m/z = 465 ($M+1$); R_t = 4.95 min.

Example 982 (General procedure (P))


- 5 2-{3-[6-(5-Cyano-1H-[1,2,3]triazol-4-yl)-naphthalen-2-yloxy]-propyl}-malonic acid



HPLC-MS (Method C): m/z = 381 ($M+1$); R_t = 3.12 min.

Example 983 (General procedure (P))


- 10 2-{5-[6-(5-Cyano-1H-[1,2,3]triazol-4-yl)-naphthalen-2-yloxy]-pentyl}-malonic acid



HPLC-MS (Method C): m/z 0 409 ($M+1$); R_t = 3.51 min.

Example 984 (General procedure (P))

- 15 4-[4-(5-Cyano-1H-[1,2,3]triazol-4-yl)-phenoxy]butyric acid





HPLC-MS (Method C): $m/z = 273$ (M+1); $R_t = 2.44$ min.


The following compounds may be prepared according to this general procedure (P):

5

4-(4-(5-Cyano-1*H*-[1,2,3]triazol-4-yl)phenoxy)butyric acid:



2-(4-(5-Cyano-1*H*-[1,2,3]triazol-4-yl)phenoxy)acetic acid:



10


4-(4-(5-Cyano-1*H*-[1,2,3]triazol-4-yl)phenoxy)butyric acid ethyl ester

5-(4-(5-Cyano-1*H*-[1,2,3]triazol-4-yl)phenoxy)pentanoic acid

8-(4-(5-Cyano-1*H*-[1,2,3]triazol-4-yl)phenoxy)octanoic acid

10-(4-(5-Cyano-1*H*-[1,2,3]triazol-4-yl)phenoxy)decanoic acid


15 12-(4-(5-Cyano-1*H*-[1,2,3]triazol-4-yl)phenoxy)dodecanoic acid

General procedure (R) for preparation of compounds of general formula I₁₂:

- 5 wherein T is as defined above and R² and R³ are hydrogen, aryl or lower alkyl, both optionally substituted.

The general procedure (R) is further illustrated by the following example:

- 10 Example 985 (General procedure (R))
Phenyl-[3-(2H-tetrazol-5-yl)-carbazol-9-yl]-methanone



- 15 2-Chlorotriptylchloride resin (100 mg, 0.114 mmol active chloride) was swelled in dichloromethane (4 mL) for 30 minutes. The solvent was drained, and a solution of 3-(2H-tetrazol-5-yl)-9H-carbazole (80 mg, 0.34 mmol) in a mixture of N,N-dimethylformamide / dichloromethane / N,N-di(2-propyl)ethylamine (5:5:1) (3 mL) was added. The reaction mixture was shaken at room temperature for 20 hours. The solvent was removed by filtration, and the resin was washed thoroughly with N,N-dimethylformamide (2 x 4 mL) and

dichloromethane (6 x 4 mL). A solution of 4-(dimethylamino)pyridine (14 mg, 0.11 mmol) and N,N-di(2-propyl)ethylamine (0.23 mL, 171 mg, 1.32 mmol) in N,N-dimethylformamide (2 mL) was added followed by benzoyl chloride (0.13 mL, 157 mg, 1.12 mmol). The mixture was shaken for 48 hours at room temperature. The drained resin was washed consecutively with
5 dichloromethane (2 x 4 mL), methanol (2 x 4 mL) and tetrahydrofuran (4 mL). The resin was treated for 2 hours at room temperature with a solution of dry hydrogen chloride in tetrahydrofuran / ethyl ether / ethanol = 8:1:1 (0.1 M, 3 mL). The reaction mixture was drained and concentrated. The crude product was stripped with dichloromethane (1.5 mL) three times to yield the title compound.

10


HPLC-MS (Method C): m/z: 340 (M+1); Rt = 3.68 min.

¹H-NMR (DMSO-d₆): δ 8.91 (1H, s), 8.34 (1H, d), 8.05 (1H, d), 7.78 (3H, m), 7.63 (3H, m), 7.46 (2H, m), 7.33 (1H, dd).

15 The compounds in the following examples were prepared in a similar fashion.

Example 986 (General procedure (R))

Phenyl-[5-(2H-tetrazol-5-yl)-indol-1-yl]-methanone




20 HPLC-MS (Method C): m/z: 290 (M+1); Rt = 3.04 min.

¹H-NMR (DMSO-d₆): δ 8.46 (1H, d), 8.42 (1H, d), 8.08 (1H, dd), 7.82 (2H, d), 7.74 (1H, t), 7.64 (2H, t), 7.55 (1H, d), 6.93 (1H, d).

Example 987 (General procedure (R))


25 (2,3-Difluorophenyl)-[5-(2H-tetrazol-5-yl)-indol-1-yl]-methanone



HPLC-MS (Method B): m/z = 326 ($M+1$); R_t = 3.85 min.

Example 988 (General procedure (R))


5 (2-Fluoro-3-(trifluoromethyl)phenyl)-[5-(2H-tetrazol-5-yl)-indol-1-yl]-methanone



HPLC-MS (Method B): m/z = 376 ($M+1$); R_t = 4.32 min.

Example 989 (General procedure (R))


10 (3-Nitrophenyl)-[5-(2H-tetrazol-5-yl)-indol-1-yl]-methanone



HPLC-MS (Method B): m/z = 335 ($M+1$); R_t = 3.72 min.

Example 990 (General procedure (R))


15 (4-Nitrophenyl)-[5-(2H-tetrazol-5-yl)-indol-1-yl]-methanone




HPLC-MS (Method B): m/z = 335 ($M+1$); R_t = 3.71 min.

Example 991 (General procedure (R))

Naphthalen-2-yl-[5-(2H-tetrazol-5-yl)-indol-1-yl]-methanone


5 HPLC-MS (Method C): $m/z = 340$ ($M+1$); $R_t = 4.25$ min.

Example 992 (General procedure (R))


HPLC-MS (Method C): $m/z: 354$ ($M+1$); $R_t = 3.91$ min.

10


Example 993 (General procedure (R))

HPLC-MS (Method C): $m/z: 418$ ($M+1$); $R_t = 4.39$ min.

15 Example 994 (General procedure (R))

HPLC-MS (Method C): $m/z: 370$ ($M+1$); $R_t = 4.01$ min.


Example 995 (General procedure (R))



HPLC-MS (Method C): m/z: 374 (M+1); Rt = 4.28 min.


5

Example 996 (General procedure (R))




HPLC-MS (Method C): m/z: 416 (M+1); Rt = 4.55 min.

10 Example 997 (General procedure (R))




HPLC-MS (Method C): m/z: 354 (M+1); Rt = 4.22 min.

Example 998 (General procedure (R))




HPLC-MS (Method C): m/z: 358 (M+1); Rt = 3.91 min.

5 Example 999 (General procedure (R))



HPLC-MS (Method C): m/z: 390 (M+1); Rt = 4.38 min.


Example 1000 (General procedure (R))



10

HPLC-MS (Method C): m/z: 418 (M+1); Rt = 4.36 min.


Example 1001 (General procedure (R))



15 HPLC-MS (Method C): m/z: 304 (M+1); Rt = 3.32 min.


368

Example 1002 (General procedure (R))




HPLC-MS (Method C): m/z: 368 (M+1); Rt = 3.84 min.

5 Example 1003 (General procedure (R))



HPLC-MS (Method C): m/z: 320 (M+1); Rt = 3.44 min.


Example 1004 (General procedure (R))



10


HPLC-MS (Method C): m/z: 324 (M+1); Rt = 3.73 min.

Example 1005 (General procedure (R))




HPLC-MS (Method C): m/z: 304 (M+1); Rt = 3.64 min.

5 Example 1006 (General procedure (R))



HPLC-MS (Method A): m/z: 308 (M+1); Rt = 3.61 min.


Example 1007 (General procedure (R))



10

HPLC-MS (Method C): m/z: 368 (M+1); Rt = 3.77 min.


Example 1008 (General procedure (R))



15 HPLC-MS (Method A): (sciex) m/z: 326 (M+1); Rt = 3.73 min.

HPLC-MS (Method C): m/z: 326 (M+1); Rt = 3.37 min.

Example 1009 (General procedure (R))



5 HPLC-MS (Method C): m/z: 374 (M+1); Rt = 4.03 min.

Example 1010 Characterization of ligand effects on physical stability of formulations by the Thioflavine T fluorescence assay.


- 10 Low physical stability of insulin formulations may lead to amyloid fibril formation, which is observed as well-ordered, thread-like macromolecular structures in the sample eventually resulting in gel formation. This has traditionally been measured by visual inspection of the sample. However, the application of a small molecule indicator probe is much more preferable. Thioflavin T is such a probe and has a distinct fluorescence signature when
- 15 binding to fibrils (or rather β -sheet rich proteins) [Naiki et al. (1989) Anal. Biochem. **177**, 244-249; LeVine (1999) Methods. Enzymol. **309**, 274-284]. Its application to insulin fibrillation has recently been validated [Nielsen et al. (2001) Biochemistry **40**, 6036-6046].

The time course for fibril formation can be described by a sigmoidal curve with the following expression:

20

$$F = f_i + m_i t + \frac{f_f + m_f t}{1 + e^{-[(t-t_0)/\tau]}} \quad \text{Eq.(2)}$$

- Here, F is the ThT fluorescence at the time t . The constant t_0 is the time needed to reach 50% of maximum fluorescence. The minimum and maximum fluorescence is denoted f_i and f_f , respectively, and the expressions $m_i t$ and $m_f t$ describe the linear development of the
- 25 bottom and top base lines. The two important parameters describing fibril formation are the lag-time calculated by $t_0 - 2\tau$ and the apparent rate constant $k_{app} = 1/\tau$.



Formation of a partially folded intermediate of the protein is suggested as a general initiating mechanism for fibrillation. Few of those intermediates nucleate to form a template onto which further intermediates may assemble and the fibrillation is initiated. The lag-time corresponds to the interval in which the critical mass of nucleus is built up and the apparent rate constant is the rate with which the fibril itself is formed.

In accordance with this mechanism, insulin needs to dissociate to its monomeric form before a partially folded intermediate may be formed. Keeping insulin on a multimeric form may therefore result in increased physical stability. Ligands binding to the insulin hexamer zinc site should stabilize the hexameric form and draw the equilibrium even further away from the monomeric form. Hence, an increased physical stability could be achieved.

Sample preparation

Insulin formulations were prepared freshly before each assay from appropriate stock solutions. Typical final concentrations were 0.6 mM human insulin or insulin aspart analogue, 0.2 mM ZnAc, 30 mM phenol, 10 mM Tris pH 8. ThT was added from a 1 mM stock solution in water to a final concentration of 1 μ M. The formulations were typically prepared in double concentration and mixed with an equal volume of test compound in appropriate concentration in 4% DMSO, 10 mM Tris pH 8.

Alternatively, insulin aspart formulations (100 U/ml) from the production line were used directly. ThT was added to 1 μ M and DMSO containing test compound in appropriate concentration to 2%.

Sample aliquots of 200 μ l were placed in a 96 well microtiter plate (Packard OptiPlate™-96, white polystyrene). Usually, eight replica of each sample (corresponding to one test

compound concentration) was placed in one column of wells. The plate was sealed with Scotch Pad (Qiagen).

Control experiments for possible test compound quenching of the ThT emission were carried out using human insulin without Zn^{2+} and phenol i.e. in a non-hexameric configuration.

- 5 Hence, the fibrillation process as well as the ThT emission should be unaffected by the presence of test compound, unless it quenched the ThT signal.

Incubation and fluorescence measurement

- 10 Temperature incubation, shaking and measurement of the ThT fluorescence were done in a Fluoroskan Ascent FL fluorescence platereader (Thermo Labsystems). Temperature setting is possible up till 45 °C, but usually sat at 30 °C. Heating was initiated at first measurement. The orbital shaking is selectable up till 1200 rpm, but adjusted to 960 rpm in all the presented data with an amplitude of 1 mm.

- 15 Fluorescence measurement was done using excitation trough a 444 nm filter and measurement of emission through a 485 nm filter. Each run was initiated by a measurement and intervals between measurements were usually 20 min. The plate was shaken and heated as adjusted between each measurement. The assay time was regulated by the number of measurements and the interval in between. Usually the plate was measured 46 times with 20 min between, i.e. over 15 hours.

20

Data handling









- The measurement points were saved in Microsoft Excel format for further processing and curve drawing and fitting was performed using GraphPad Prism. The background emission from ThT in the absence of fibrils was negligible. Some test compounds had background
25 fluorescence under the applied experimental conditions. This was eliminated by subtracting the mean value of the first measurement from the data set for this test compound. The data points are shown with standard deviation.

- The data set may be fitted to Eq. (2). However, since the stabilizing effect of the test compounds/ligands were so significant that a full sigmoid curve was not obtained during the
30 usual assay time, curve fitting to such a data set would be imprecise and hence meaningless.

Only data obtained in the same experiment (i.e. samples on the same plate) are presented in the same graph.

- 35 Examples & Results

The various ligands are shown below with structure and affinity towards the zinc site as measured by the TZD-assay described in "Analytical Methods".

| Reference | Example | Compound | $K_d(\text{app})$ (nM) |
|-----------|---------|---|---------------------------|
| A | 533 |  | 383 |
| C | 462 |  | 58 |
| D | 738 |  | 171 |
| E | 68 |  | 23 |
| F | 756 |  | 3 |
| G | 1 |  | 3879 |
| H | 76 |  | 82 |
| I | 759 |  | 23 |

The ThT assays of various combinations of insulin formulations and ligands are shown in Fig. 1 – 8.

Addition of ligands improves the physical stability of insulin formulations. This holds for human insulin formulations (see Fig 1) as well as insulin aspart formulations (rest of data set).

The improved stability can be obtained by using various compound classes as zinc site anchor, e.g. benzothiazoles (G, Fig 1), naphthosalicylic acids (A, Fig 2), thiazolidine-diones (E, Figs 4, 7; C, Fig 2; H, Figs 6, 8) and tetrazoles (D, Fig 3; F, Fig 8; I, Fig 5).

- 5 Increased affinity of the ligand results in higher stability of the formulation. Compare the effect of the weakest binding ligand in 2 mM (G, Fig. 1) with the effect on an insulin aspart formulation of 0,5 mM E (Fig. 4). Also compare the effects of similar concentrations of A and C (Fig 2); and of D (fig. 3) and E (Fig 4) on insulin aspart formulations.
- 10 Increasing the concentration of ligand tends to improve the stabilization (see Figs. 1, 3, 4, 5, 6). In some instances, more pronounced effects are seen with the ligand in slight molar excess to the zinc sites, see Figs. 4, 6, whereas it seems to plateau around the stoichiometric concentration in other instances (Figs. 3, 5).
- 15 Of the presented ligands, A, C, D, E, F were tested in a disappearance assay for the effect on release of insulin aspart from a subcutaneous inject site. Surprisingly, the ligands had no effect on the insulin aspart disappearance. In a very limited way, this can be mimicked in the ThT assay by increasing the assay temperature to 37 °C (see Figs. 7, 8). The stabilizing effect is somewhat attenuated, e.g. compare E at 30 °C (Fig 4) and 37 °C (Fig 7), and H (Fig 6 and 8). The ligand with highest affinity (F) has the most stabilizing effect at 37 °C (Fig. 8).
- 20





Fig 3

Aspart Formulation, 1 μ M ThT, 30 °C, 960 rpm.




Fig 4

Aspart formulation, 1 μ M ThT, 30 °C, 960 rpm.




Fig 5

Aspart formulation, 1 μ M ThT, pH 8, 30 °C, 960 rpm.







Fig 6

Aspart formulation, 1 μ M ThT, pH 8, 30 °C, 960 rpm.











Example 1011 Retention of fast absorption characteristics of formulations stabilized by addition of His^{B10} Zn²⁺-site ligands.

5

Formulations of the present invention were characterized by the disappearance rate from the subcutaneous depot following injection in pigs. Formulations of B28 Asp human insulin containing A14Tyr(¹²⁵I) B28 Asp human insulin were followed with an external γ -counter (Ribel et al., The pig as a model for subcutaneous absorption in man. In: Serrano-Ritos & Lefebvre (Eds.): Diabetes (1985) proceedings of the 12th congress of the international diabetes federation, Madrid, Spain, 1985 (Excerpta Medica, Amsterdam (1986) 891-896. Formulations of Insulin Aspart (0.6 mM, U100) containing 0.3 mM Zn²⁺, 30 mM phenol, 2 mM phosphate buffer, and 1.6% glycerol, pH 7.4, were compared with the corresponding formulations containing 0.3 mM of the ligands shown below: where T_{50%} is the time when 50% of the A14Tyr(¹²⁵I) B28 Asp insulin has disappeared from the site of injection and K_d is the affinity of the ligand as measured by the TZD-assay described in "Analytical methods" below. It is evident that the stabilizing ligands do not affect the fast absorption properties of the formulations

20

| | Example # | Compound | T50% | K _d (app) (nM) |
|-----------|-----------|----------|----------|------------------------------|
| reference | | | 1.3± 0.3 | |

| | | | | |
|---|-----|---|-----------|-----|
| A | 533 |  | 1.0± 0.23 | 383 |
| B | 476 |  | 1.1± 0.37 | 68 |
| C | 462 |  | 0.9± 0.19 | 58 |
| D | 738 |  | 1.3± 0.36 | 171 |
| E | 68 |  | 1.2± 0.41 | 23 |
| F | 756 |  | 1.2± 0.27 | 3 |






Figure 9

Example 1012

Reference experiment

- 5 The chemical stability of insulin formulations of the invention was characterised by HPLC (RPC, reverse phase chromatography and SEC, size exclusion chromatography). As reference, insulin formulated without ligands of the invention but with 0.3 % DMSO was also investigated and shown below:



10

Figure 10

Fig 10. Reverse phase chromatography of formulated human insulin with 3 Zn^{2+} per hexamer, 30 mM phenol, 150 mM mannitol, 3 mM phosphoric acid, sodium hydroxide to pH 7.4 and 0.3 % DMSO corresponding to 3 ligands per hexamer at start (upper panel) and after storage for 2 weeks at 37 °C (lower panel): Preservatives before 20 min., "hydrophilic

15

derivatives" (desamido-insulins) 20 min to main top insulin, "hydrophobic derivatives 1" main top to 64 min., and "hydrophobic derivatives 2" (insulin dimers) after 64 min.


Storage in HPLC 1 ml vials at 45 °C (5 d), 37 °C (2 w), 30 °C (6 w), 25 °C (10 w), 15 °C (30 w) gave about the same increase of transformation products correlating to an increase in reactions constants of a factor 3-4 per 10 degree..

RPC (reverse phase chromatography) on Waters SymmetryShield RP₈ column, 150×4.6 mm and 3.5 μm, eluted by **A**: 0.2 M sodium sulfate + 0.04 M sodium phosphate pH 7.2 + 10 % acetonitrile and isocratically (i) or a gradient (g) of **B**: 70 % acetonitrile [minutes/%B(i/g): 0/19, 21/24(i)(sudden change), 51/24(i), 81/39(g), 81.1/0(i), 82.3/19(i)] at flow of 0.9 mL/min and 30 °C.

SEC (size exclusion chromatography) on Waters insulin HMWP column, 300×7.8 mm, eluted by 15:20:65 of acetic acid: acetonitrile: arginine 1 g/L at flow of 1 mL/min and ambient temperature.

| % formed/year | RPC | | | SEC |
|---------------|---------|---------|---------|---------|
| | hphil | hphob1 | hphob2 | dimer |
| 45°C | 153.3 | 47.5 | 23.4 | 21.9 |
| 37°C | 57.2 | 15.6 | 5.20 | 6.24 |
| 30°C | 24.5 | 4.16 | 1.30 | 2.08 |
| 25°C | 14.2 | 2.13 | 0.68 | 1.09 |
| 15°C | 4.12 | 0.43 | 0.22 | 0.31 |
| 5°predicted | (1.001) | (0.066) | (0.027) | (0.052) |

Arrhenius Plot




The chemical stability of insulin formulations of the invention was likewise characterised by HPLC (RPC, reverse phase chromatography and SEC, size exclusion chromatography). Compared to the reference the formulations of the invention were shown to be more chemically stable.

5

Example 1013

Chemical stability of insulin formulated with the compound of example 533, 7-bromo-3-hydroxy-2-naphthoic acid:



10

Figure 11

Fig 11. Reverse phase chromatography of formulated human insulin as described for the reference example and added 3 ligands of Example 533 and 3 Zn^{2+} per hexamer at start (upper panel) and after storage for 2 weeks at 37 °C (lower panel): Preservatives before 20 min., "hydrophilic derivatives" (desamido-insulins) 20 min to main top insulin, "hydrophobic derivatives 1" main top to 64 min., and "hydrophobic derivatives 2" (insulin dimers) after 64 min.

15

Storage in HPLC 1 ml vials at 45 °C (5 d), 37 °C (2 w), 30 °C (6 w), 25 °C (10 w), 15 °C (30 w) will give about the same increase in transformation products correlating to an increase in reaction constants of a factor 3-4 per 10 degrees...

5

RPC (reverse phase chromatography) on Waters SymmetryShield RP₈ column, 150×4.6 mm and 3.5 µm, eluted by **A**: 0.2 M sodium sulfate + 0.04 M sodium phosphate pH 7.2 + 10 % acetonitrile and isocratically (i) or a gradient (g) of **B**: 70 % acetonitrile [minutes/%B(i/g): 0/19, 21/24(i)(sudden change), 51/24(i), 81/39(g), 81.1/0(i), 82.3/19(i)] at flow of 0.9 mL/min and 30 °C.


10

SEC (size exclusion chromatography) on Waters insulin HMWP column, 300×7.8 mm, eluted by 15:20:65 of acetic acid: acetonitrile:arginine 1 g/L at flow of 1 mL/min and ambient temperature.

15


| % formed/year | RPC | | | SEC |
|---------------|--------|--------|---------|---------|
| | hphil | hphob1 | hphob2 | dimer |
| 45°C | 30.7 | 62.1 | 23.4 | 26.3 |
| 37°C | 12.5 | 22.4 | 7.02 | 7.02 |
| 30°C | 6.76 | 6.42 | 2.25 | 2.86 |
| 25°C | 4.06 | 5.67 | 1.46 | 1.61 |
| 15°C | 1.80 | 0.85 | 0.42 | 0.45 |
| 5°predicted | (0.55) | (0.19) | (0.079) | (0.087) |

Arrhenius Plot



Example 1014

Chemical stability of insulin formulated with the compound of Example 462, 3-[4-(2,4-dioxothiazolidin-5-ylidenemethyl)phenyl]acrylic acid:



5

Figure 12

Fig 12. Reverse phase chromatography of formulated human insulin as described for the reference example and added 3 ligands of example 462 and 3 Zn^{2+} per hexamer at start (upper panel) and after storage for 2 weeks at 37 °C (lower panel): Preservatives before 20 min., "hydrophilic derivatives" (desamido-insulins) 20 min to main top insulin, "hydrophobic derivatives 1" main top to 64 min., and "hydrophobic derivatives 2" (insulin dimers) after 64 min.

Storage in HPLC 1 ml vials at 45 °C (5 d), 37 °C (2 w), 30 °C (6 w), 25 °C (10 w), 15 °C (30 w) will give about the same increase in transformation products correlating to an increase in reaction constants of a factor 3-4 per 10 degrees.


- 5 **RPC** (reverse phase chromatography) on Waters SymmetryShield RP₈ column, 150×4.6 mm and 3.5 µm, eluted by **A**: 0.2 M sodium sulfate + 0.04 M sodium phosphate pH 7.2 + 10 % acetonitrile and isocratically (i) or a gradient (g) of **B**: 70 % acetonitrile [minutes/%B(i/g): 0/19, 21/24(i)(sudden change), 51/24(i), 81/39(g), 81.1/0(i), 82.3/19(i)] at flow of 0.9 mL/min and 30 °C.

10

SEC (size exclusion chromatography) on Waters insulin HMWP column, 300×7.8 mm, eluted by 15:20:65 of acetic acid: acetonitrile:arginine 1 g/L at flow of 1 mL/min and ambient temperature.

| % formed/year | RPC | | | SEC |
|---------------|---------|---------|---------|---------|
| | hphil | hphob1 | hphob2 | Dimer |
| 45°C | 20.4 | 43.8 | 17.5 | 21.2 |
| 37°C | 9.88 | 19.5 | 4.68 | 6.24 |
| 30°C | 4.07 | 4.42 | 1.65 | 2.17 |
| 25°C | 2.60 | 6.14 | 0.78 | 1.09 |
| 15°C | 1.02 | 1.28 | 0.38 | 0.28 |
| 5°predicted | (0.293) | (0.336) | (0.062) | (0.048) |

15

**Example 1015**

Chemical stability of insulin formulated with the compound of example 461, [3-(2,4-Dioxothiazolidin-5-ylidenemethyl)phenoxy]acetic acid:




Fig 13. Reverse phase chromatography of formulated human insulin added 3 ligands (#) and 3 Zn per hexamer at start and storage of 2 w 37 °C: Preservatives before 20 min., “hydrophilic derivatives” (desamido-insulins) 20 min to main top insulin, “hydrophobic derivatives 1” main top to 64 min., and “hydrophobic derivatives 2” (insulin dimers) after 64 min.

Storage in HPLC 1 ml vials at 45 °C (5 d), 37 °C (2 w), 30 °C (6 w), 25 °C (10 w), 15 °C (30 w) will give about the same increase of 0.7 % hfil, 0.6 % hfob1, 0.3 % hfob2 and 0.3 % dimer solution 1, correlating to Q_{10} of 3 below 30 °C and 4 at higher temperature for ref..


RPC (reverse phase chromatography) on Waters SymmetryShield RP₈ column, 150×4.6 mm and 3.5 μm, eluted by **A**: 0.2 M sodium sulfate + 0.04 M sodium phosphate pH 7.2 + 10 % acetonitrile and isocratically (i) or a gradient (g) of **B**: 70 % acetonitrile [minutes/%B(i/g): 0/19, 21/24(i)(sudden change), 51/24(i), 81/39(g), 81.1/0(i), 82.3/19(i)] at flow of 0.9 mL/min and 30 °C.

SEC (size exclusion chromatography) on Waters insulin HMWP column, 300×7.8 mm, eluted by 15:20:65 of acetic acid: acetonitrile:arginine 1 g/L at flow of 1 mL/min and ambient temperature.

| % formed/year | RPC | | | SEC |
|---------------|---------|---------|---------|---------|
| | hphil | hphob1 | hphob2 | dimer |
| 45°C | 29.9 | 52.6 | 23.4 | 23.4 |
| 37°C | 11.2 | 20.0 | 7.02 | 6.50 |
| 30°C | 5.12 | 7.11 | 1.99 | 2.34 |
| 25°C | 2.86 | 4.78 | 0.99 | 1.20 |
| 15°C | 1.14 | 0.67 | 0.29 | 0.33 |
| 5°predicted | (0.287) | (0.153) | (0.044) | (0.057) |

5

Arrhenius Plot



Example 1016

Chemical stability of insulin formulated with the compound of example 70 , 5-(4-Diethylaminobenzylidene)thiazolidine-2,4-dione

10




Fig 14. Reverse phase chromatography of formulated human insulin added 3 ligands (#) and 3 Zn per hexamer at start and storage of 2 w 37 °C: Preservatives before 20 min.,

- 5 "hydrophilic derivatives" (desamido-insulins) 20 min to main top insulin, "hydrophobic derivatives 1" main top to 64 min., and "hydrophobic derivatives 2" (insulin dimers) after 64 min.


- 10 **Storage** in HPLC 1 ml vials at 45 °C (5 d), 37 °C (2 w), 30 °C (6 w), 25 °C (10 w), 15 °C (30 w) will give about the same increase of 0.7 % hfil, 0.6 % hfob1, 0.3 % hfob2 and 0.3 % dimer solution 1, correlating to Q_{10} of 3 below 30 °C and 4 at higher temperature for ref..

- 15 **RPC** (reverse phase chromatography) on Waters SymmetryShield RP₈ column, 150×4.6 mm and 3.5 μ m, eluted by **A**: 0.2 M sodium sulfate + 0.04 M sodium phosphate pH 7.2 + 10 % acetonitrile and isocratically (i) or a gradient (g) of **B**: 70 % acetonitrile [minutes/%B(i/g): 0/19, 21/24(i)(sudden change), 51/24(i), 81/39(g), 81.1/0(i), 82.3/19(i)] at flow of 0.9 mL/min and 30 °C.

SEC (size exclusion chromatography) on Waters insulin HMWP column, 300×7.8 mm, eluted by 15:20:65 of acetic acid: acetonitrile:arginine 1 g/L at flow of 1 mL/min and ambient temperature.

| % formed/year | RPC | | | SEC |
|---------------|--------|--------|---------|--------|
| | hphil | hphob1 | hphob2 | dimer |
| 45°C | 56.2 | 78.8 | 27.0 | 29.2 |
| 37°C | 23.7 | 14.8 | 6.50 | 8.84 |
| 30°C | 12.1 | 6.94 | 3.03 | 3.55 |
| 25°C | 6.08 | 4.99 | 0.94 | 2.24 |
| 15°C | 1.95 | 1.09 | 0.42 | 0.69 |
| 5°predicted | (0.54) | (0.22) | (0.064) | (0.15) |

5



ANALYTICAL METHODS

Assays to quantify the binding affinity of ligands to the metal site of the insulin R₆ hexamers:

5 4H3N-assay:

The binding affinity of ligands to the metal site of insulin R₆ hexamers are measured in a UV/vis based displacement assay. The UV/vis spectrum of 3-hydroxy-4-nitro benzoic acid (4H3N) which is a known ligand for the metal site of insulin R₆ shows a shift in absorption maximum upon displacement from the metal site to the solution (Huang et al., 1997,

10 Biochemistry 36, 9878-9888). Titration of a ligand to a solution of insulin R₆ hexamers with 4H3N mounted in the metal site allows the binding affinity of these ligands to be determined following the reduction of absorption at 444 nm.

A stock solution with the following composition 0.2 mM human insulin, 0.067 mM Zn-acetate, 40 mM phenol, 0.101 mM 4H3N is prepared in a 10mL quantum as described below. Buffer

15 is always 50mM tris buffer adjusted to pH=8.0 with NaOH/CIO₄⁻.

1000 µL of 2.0mM human insulin in buffer

66.7 µL of 10mM Zn-acetate in buffer

20 800 µL of 500mM phenol in H₂O

201 µL of 4H3N in H₂O

7.93 ml buffer

The ligand is dissolved in DMSO to a concentration of 20 mM.

25 The ligand solution is titrated to a cuvette containing 2 mL stock solution and after each addition the UV/vis spectrum is measured. The titration points are listed in Table 3 below.

Table 3

| ligand addition (μ l) | ligand conc. (mM) | dilution factor |
|-------------------------------|----------------------|-----------------|
| 1 | 0.010 | 1.0005 |
| 1 | 0.020 | 1.0010 |
| 1 | 0.030 | 1.0015 |
| 2 | 0.050 | 1.0025 |
| 5 | 0.100 | 1.0050 |
| 10 | 0.198 | 1.0100 |
| 20 | 0.392 | 1.0200 |
| 20 | 0.583 | 1.0300 |
| 20 | 0.769 | 1.0400 |
| 20 | 0.952 | 1.0500 |

5 The UV/vis spectra resulting from a titration of the compound 3-hydroxy-2-naphthoic acid is shown in Figure 5. Inserted in the upper right corner is the absorbance at 444nm vs. the concentration of ligand.

The following equation is fitted to these datapoints to determine the two parameters $K_D(\text{obs})$, the observed dissociation constant, and abs_{max} the absorbance at maximal ligand concentration.

$$10 \quad \text{abs}([\text{ligand}]_{\text{free}}) = (\text{abs}_{\text{max}} * [\text{ligand}]_{\text{free}}) / (K_D(\text{obs}) + [\text{ligand}]_{\text{free}})$$

The observed dissociation constant is recalculated to obtain the apparent dissociation constant

$$15 \quad K_D(\text{app}) = K_D(\text{obs}) / (1 + [4\text{H3N}]/K_{4\text{H3N}})$$

The value of $K_{4\text{H3N}}=50 \mu\text{M}$ is taken from Huang et al., 1997, Biochemistry 36, 9878-9888.

TZD-assay:

20 The binding affinity of ligands to the metal site of insulin R_6 hexamers are measured in a fluorescence based displacement assay. The fluorescence of 5-(4-dimethylaminobenzylidene)thiazolidine-2,4-dione (TZD) which is a ligand for the metal site of insulin R_6 is quenched upon displacement from the metal site to the solution. Titration of a ligand to a stock solution of insulin R_6 hexamers with this compound mounted in the metal

site allows the binding affinity of these ligands to be determined measuring the fluorescence at 455nm upon excitation at 410nm.

Preparation

- 5 Stock solution: 0.02 mM human insulin, 0.007 mM Zn-acetate, 40 mM phenol, 0.01 mM TZD in 50mM tris buffer adjusted to pH=8.0 with NaOH/CIO₄⁻.

The ligand is dissolved in DMSO to a concentration of 5 mM and added in aliquots to the stock solution to final concentrations of 0-250 μ M.

10 Measurements

Fluorescence measurements were carried out on a Perkin Elmer Spectrofluorometer LS50B. The main absorption band was excited at 410 nm and emission was detected at 455 nm. The resolution was 10 nm and 2.5 nm for excitation and emission, respectively.

15 Data analysis

This equation is fitted to the datapoints

$$\Delta F(455\text{nm}) = \Delta F_{\text{max}} * [\text{ligand}]_{\text{free}} / (K_D(\text{app}) * (1 + [\text{TZD}]/K_{\text{TZD}}) + [\text{ligand}]_{\text{free}})$$

K_D(app) is the apparent dissociation constant and F_{max} is the fluorescence at maximal ligand concentration. The value of K_{TZD} is measured separately to 230 nM


20

Two different fitting-procedures can be used. One in which both parameters, K_D(app) and F_{max}, are adjusted to best fit the data and a second in which the value of F_{max} is fixed (F_{max}=1) and only K_D(app) is adjusted. The given data are from the second fitting procedure. The Solver module of Microsoft Excel can be used to generate the fits from the data points.

25

CLAIMS

1. A pharmaceutical composition comprising insulin and a zinc-binding ligand which reversibly binds to a His^{B10} Zn²⁺ site of an insulin hexamer, wherein the ligand is selected from the group consisting of
- 5 benzotriazoles, 3-hydroxy 2-napthoic acids, salicylic acids, tetrazoles, thiazolidinediones, 5-mercaptotetrazoles, or 4-cyano-1,2,3-triazoles, or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.
- 10 2. A pharmaceutical composition according to claim 1 wherein the zinc-binding ligand is



wherein

X is =O, =S or =NH

Y is -S-, -O- or -NH-

15

R¹, R^{1A} and R⁴ are independently selected from hydrogen or C₁-C₆-alkyl,

R² and R^{2A} are hydrogen or C₁-C₆-alkyl or aryl, R¹ and R² may optionally be combined to form a double bond, R^{1A} and R^{2A} may optionally be combined to form a double bond,

- 15 R³, R^{3A} and R⁵ are independently selected from hydrogen, halogen, aryl optionally substituted with one or more substituents independently selected from R¹⁶, C₁-C₆-alkyl, or -C(O)NR¹¹R¹²,
- 20

A, A¹ and B are independently selected from C₁-C₆-alkyl, aryl, aryl-C₁-C₆-alkyl, -NR¹¹-aryl, aryl-C₂-C₆-alkenyl or heteroaryl, wherein the alkyl or alkenyl is optionally substituted with one or more substituents independently selected from R⁶ and the aryl or heteroaryl is optionally substituted with up to four substituents R⁷, R⁸, R⁹, and R¹⁰,

25

A and R³ may be connected through one or two valence bonds, B and R⁵ may be connected through one or two valence bonds,

R⁶ is independently selected from halogen, -CN, -CF₃, -OCF₃, aryl, -COOH and -NH₂,

- 30 R⁷, R⁸, R⁹ and R¹⁰ are independently selected from

• hydrogen, halogen, -CN, -CH₂CN, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃,
 -OCF₂CHF₂, -S(O)₂CF₃, -OS(O)₂CF₃, -SCF₃, -NO₂, -OR¹¹, -NR¹¹R¹², -SR¹¹,
 -NR¹¹S(O)₂R¹², -S(O)₂NR¹¹R¹², -S(O)NR¹¹R¹², -S(O)R¹¹, -S(O)₂R¹¹, -OS(O)₂R¹¹,
 -C(O)NR¹¹R¹², -OC(O)NR¹¹R¹², -NR¹¹C(O)R¹², -CH₂C(O)NR¹¹R¹²,
 -OC₁-C₆-alkyl-C(O)NR¹¹R¹², -CH₂OR¹¹, -CH₂OC(O)R¹¹, -CH₂NR¹¹R¹², -OC(O)R¹¹,
 -OC₁-C₁₅-alkyl-C(O)OR¹¹, -OC₁-C₆-alkyl-OR¹¹, -SC₁-C₆-alkyl-C(O)OR¹¹,
 -C₂-C₆-alkenyl-C(=O)OR¹¹, -NR¹¹-C(=O)-C₁-C₆-alkyl-C(=O)OR¹¹,
 -NR¹¹-C(=O)-C₁-C₆-alkenyl-C(=O)OR¹¹, -C(O)OR¹¹, C(O)R¹¹, or -C₂-C₆-alkenyl-
 C(=O)R¹¹, =O, or -C₂-C₆-alkenyl-C(=O)-NR¹¹R¹²,

• C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl, each of which may optionally be substituted with one or more substituents independently selected from R¹³,

• aryl, aryloxy, aryloxycarbonyl, aroyl, arylsulfanyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aroyl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl, heteroaryl-C₂-C₆-alkynyl, or C₃-C₆ cycloalkyl,

of which each cyclic moiety may optionally be substituted with one or more substituents independently selected from R¹⁴,

R¹¹ and R¹² are independently selected from hydrogen, OH, C₁-C₂₀-alkyl, aryl-C₁-C₆-alkyl or aryl, wherein the alkyl groups may optionally be substituted with one or more substituents independently selected from R¹⁵, and the aryl groups may optionally be substituted one or more substituents independently selected from R¹⁶; R¹¹ and R¹² when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom, the heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds,

R¹³ is independently selected from halogen, -CN, -CF₃, -OCF₃, -OR¹¹, -C(O)OR¹¹, -NR¹¹R¹², and -C(O)NR¹¹R¹²,

R¹⁴ is independently selected from halogen, -C(O)OR¹¹, -CH₂C(O)OR¹¹, -CH₂OR¹¹, -CN, -CF₃, -OCF₃, -NO₂, -OR¹¹, -NR¹¹R¹², -NR¹¹C(O)R¹¹, -S(O)₂R¹¹, aryl and C₁-C₆-alkyl,

R¹⁵ is independently selected from halogen, -CN, -CF₃, =O, -OCF₃, -OC₁-C₆-alkyl, -C(O)OC₁-C₆-alkyl, -COOH and -NH₂.

5 R¹⁶ is independently selected from halogen, -C(O)OC₁-C₆-alkyl, -COOH, -CN, -CF₃, -OCF₃, -NO₂, -OH, -OC₁-C₆-alkyl, -NH₂, C(=O) or C₁-C₆-alkyl, or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.

3. A pharmaceutical composition according to claim 2 wherein X is =O or =S.

10 4. A pharmaceutical composition according to claim 3 wherein X is =O.

5. A pharmaceutical composition according to claim 3 wherein X is =S.

6. A pharmaceutical composition according to any one of the claims 2 to 5 wherein Y is -O- or -S-.

7. A pharmaceutical composition according to claim 6 wherein Y is -O-.

15 8. A pharmaceutical composition according to claim 6 wherein Y is -NH-.


9. A pharmaceutical composition according to claim 6 wherein Y is -S-.

10. A pharmaceutical composition according to any one of the claims 2 to 9 wherein A is aryl optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.

20 11. A pharmaceutical composition according to claim 10 wherein A is selected from ArG1 optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.

12. A pharmaceutical composition according to claim 11 wherein A is phenyl or naphthyl optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.

25 13. A pharmaceutical composition according to claim 12 wherein A is



14. A pharmaceutical composition according to claim 12 wherein A is phenyl.

15. A pharmaceutical composition according to any one of the claims 2 to 9 wherein A is heteroaryl optionally substituted with up to four substituents, R⁷, R⁸, R⁹, and R¹⁰ which may be the same or different.

30

16. A pharmaceutical composition according to claim 15 wherein A is selected from Het1 optionally substituted with up to four substituents, R^7 , R^8 , R^9 , and R^{10} which may be the same or different.


17. A pharmaceutical composition according to claim 16 wherein A is selected from Het2
5 optionally substituted with up to four substituents, R^7 , R^8 , R^9 , and R^{10} which may be the same or different.

18. A pharmaceutical composition according to claim 17 wherein A is selected from Het3 optionally substituted with up to four substituents, R^7 , R^8 , R^9 , and R^{10} which may be the same or different.

10 19. A pharmaceutical composition according to claim 18 wherein A is selected from the group consisting of indolyl, benzofuranyl, quinolyl, furyl, thienyl, or pyrrolyl, wherein each heteroaryl may optionally substituted with up to four substituents, R^7 , R^8 , R^9 , and R^{10} which may be the same or different.


20. A pharmaceutical composition according to claim 18 wherein A is benzofuranyl optionally
15 substituted with up to four substituents R^7 , R^8 , R^9 , and R^{10} which may be the same or different.

21. A pharmaceutical composition according to claim 20 wherein A is




22. A pharmaceutical composition according to claim 18 wherein A is carbazolyl optionally
20 substituted with up to four substituents R^7 , R^8 , R^9 , and R^{10} which may be the same or different.

23. A pharmaceutical composition according to claim 22 wherein A is




24. A pharmaceutical composition according to claim 18 wherein A is quinolyl optionally
25 substituted with up to four substituents R^7 , R^8 , R^9 , and R^{10} which may be the same or different.

25. A pharmaceutical composition according to claim 24 wherein A is



26. A pharmaceutical composition according to claim 18 wherein A is indolyl optionally substituted with up to four substituents R^7 , R^8 , R^9 , and R^{10} which may be the same or different.

5 27. A pharmaceutical composition according to claim 26 wherein A is



28. A pharmaceutical composition according to any one of the claims 2 to 27 wherein R^1 is hydrogen.

29. A pharmaceutical composition according to any one of the claims 2 to 28 wherein R^2 is hydrogen.

30. A pharmaceutical composition according to any one of the claims 2 to 27 wherein R^1 and R^2 are combined to form a double bond.

31. A pharmaceutical composition according to any one of the claims 2 to 30 wherein R^3 is C_1 - C_6 -alkyl, halogen, or $C(O)NR^{16}R^{17}$.

15 32. A pharmaceutical composition according to claim 31 wherein R^3 is C_1 - C_6 -alkyl or $C(O)NR^{16}R^{17}$.

33. A pharmaceutical composition according to claim 32 wherein R^3 is methyl.

34. A pharmaceutical composition according to any one of the claims 2 to 9 wherein B is phenyl optionally substituted with up to four substituents, R^7 , R^8 , R^9 , and R^{10} which may be the same or different.

35. A pharmaceutical composition according to any one of the claims 2 to 9 or 34 wherein R^4 is hydrogen.

36. A pharmaceutical composition according to any one of the claims 2 to 9 or 34 to 35 wherein R^5 is hydrogen.

25 37. A pharmaceutical composition according to any one of the claims 2 to 36 wherein R^6 is aryl.

38. A pharmaceutical composition according to claim 37 wherein R^6 is phenyl.

39. A pharmaceutical composition according to any one of the claims 2 to 38 wherein R^7 , R^8 , R^9 and R^{10} are independently selected from

• hydrogen, halogen, $-\text{NO}_2$, $-\text{OR}^{11}$, $-\text{NR}^{11}\text{R}^{12}$, $-\text{SR}^{11}$, $-\text{NR}^{11}\text{S}(\text{O})_2\text{R}^{12}$, $-\text{S}(\text{O})_2\text{NR}^{11}\text{R}^{12}$,
5 $-\text{S}(\text{O})\text{NR}^{11}\text{R}^{12}$, $-\text{S}(\text{O})\text{R}^{11}$, $-\text{S}(\text{O})_2\text{R}^{11}$, $-\text{OS}(\text{O})_2\text{R}^{11}$, $-\text{NR}^{11}\text{C}(\text{O})\text{R}^{12}$, $-\text{CH}_2\text{OR}^{11}$, $-\text{CH}_2\text{OC}(\text{O})\text{R}^{11}$, $-\text{CH}_2\text{NR}^{11}\text{R}^{12}$, $-\text{OC}(\text{O})\text{R}^{11}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{11}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{NR}^{11}\text{R}^{12}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-OR}^{11}$, $-\text{SC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{11}$, $-\text{C}_2\text{-C}_6\text{-alkenyl-C}(\text{=O})\text{OR}^{11}$, $-\text{C}(\text{O})\text{OR}^{11}$, or $-\text{C}_2\text{-C}_6\text{-alkenyl-C}(\text{=O})\text{R}^{11}$,

10 • $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_2\text{-C}_6\text{-alkenyl}$ or $\text{C}_2\text{-C}_6\text{-alkynyl}$, which may each optionally be substituted with one or more substituents independently selected from R^{13}

• aryl, aryloxy, aroyl, arylsulfanyl, aryl- $\text{C}_1\text{-C}_6\text{-alkoxy}$, aryl- $\text{C}_1\text{-C}_6\text{-alkyl}$, aryl- $\text{C}_2\text{-C}_6\text{-alkenyl}$, aroyl- $\text{C}_2\text{-C}_6\text{-alkenyl}$, aryl- $\text{C}_2\text{-C}_6\text{-alkynyl}$, heteroaryl, heteroaryl- $\text{C}_1\text{-C}_6\text{-alkyl}$,
15 wherein each of the cyclic moieties optionally may be substituted with one or more substituents independently selected from R^{14}

40. A pharmaceutical composition according to claim 39 wherein R^7 , R^8 , R^9 and R^{10} are independently selected from

20 • hydrogen, halogen, $-\text{NO}_2$, $-\text{OR}^{11}$, $-\text{NR}^{11}\text{R}^{12}$, $-\text{SR}^{11}$, $-\text{S}(\text{O})_2\text{R}^{11}$, $-\text{OS}(\text{O})_2\text{R}^{11}$, $-\text{CH}_2\text{OC}(\text{O})\text{R}^{11}$, $-\text{OC}(\text{O})\text{R}^{11}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{11}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-OR}^{11}$, $-\text{SC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{11}$, $-\text{C}(\text{O})\text{OR}^{11}$, or $-\text{C}_2\text{-C}_6\text{-alkenyl-C}(\text{=O})\text{R}^{11}$,

25 • $\text{C}_1\text{-C}_6\text{-alkyl}$ or $\text{C}_1\text{-C}_6\text{-alkenyl}$ which may each optionally be substituted with one or more substituents independently selected from R^{13}

• aryl, aryloxy, aroyl, aryl- $\text{C}_1\text{-C}_6\text{-alkoxy}$, aryl- $\text{C}_1\text{-C}_6\text{-alkyl}$, heteroaryl,

30 of which each of the cyclic moieties optionally may be substituted with one or more substituents independently selected from R^{14}

41. A pharmaceutical composition according to claim 40 wherein R^7 , R^8 , R^9 and R^{10} are independently selected from

• hydrogen, halogen, -NO₂, -OR¹¹, -NR¹¹R¹², -SR¹¹, -S(O)₂R¹¹, -OS(O)₂ R¹¹, -CH₂OC(O)R¹¹, -OC(O)R¹¹, -OC₁-C₆-alkyl-C(O)OR¹¹, -OC₁-C₆-alkyl-OR¹¹, -SC₁-C₆-alkyl-C(O)OR¹¹, -C(O)OR¹¹, or -C₂-C₆-alkenyl-C(=O)R¹¹,

5 • C₁-C₆-alkyl or C₁-C₆- which may each optionally be substituted with one or more substituents independently selected from R¹³

• aryl, aryloxy, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, heteroaryl,

10 of which each of the cyclic moieties optionally may be substituted with one or more substituents independently selected from R¹⁴.

42. A pharmaceutical composition according to claim 41 wherein R⁷, R⁸, R⁹ and R¹⁰ are independently selected from

15 • hydrogen, halogen, -OR¹¹, -OC₁-C₆-alkyl-C(O)OR¹¹, or -C(O)OR¹¹,

• C₁-C₆-alkyl which may each optionally be substituted with one or more substituents independently selected from R¹³

20 • aryl, aryloxy, aryl-C₁-C₆-alkoxy,

of which each of the cyclic moieties optionally may be substituted with one or more substituents independently selected from R¹⁴.

25 43. A pharmaceutical composition according to claim 42 wherein R⁷, R⁸, R⁹ and R¹⁰ are independently selected from

• hydrogen, halogen, -OR¹¹, -OC₁-C₆-alkyl-C(O)OR¹¹, or -C(O)OR¹¹,

30 • C₁-C₆-alkyl which may each optionally be substituted with one or more substituents independently selected from R¹³

• ArG1, ArG1oxy, ArG1-C₁-C₆-alkoxy,

35 of which each of the cyclic moieties optionally may be substituted with one or more substituents independently selected from R¹⁴.

44. A pharmaceutical composition according to claim 43 wherein R⁷, R⁸, R⁹ and R¹⁰ are independently selected from

- hydrogen, halogen, -OR¹¹, -OC₁-C₆-alkyl-C(O)OR¹¹, or -C(O)OR¹¹,

5 • C₁-C₆-alkyl which may optionally be substituted with one or more substituents independently selected from R¹³

10 • phenyl, phenoxy, phenyl-C₁-C₆-alkoxy, wherein each of the cyclic moieties optionally may be substituted with one or more substituents independently selected from R¹⁴.

45. A pharmaceutical composition according to any one of the claims 2 to 44 wherein R¹¹ and R¹² are independently selected from hydrogen, C₁-C₂₀-alkyl, aryl or aryl-C₁-C₆-alkyl, wherein the alkyl groups may optionally be substituted with one or more substituents independently
15 selected from R¹⁵, and the aryl groups may optionally be substituted one or more substituents independently selected from R¹⁶; R¹¹ and R¹² when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom, the heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds.

20 46. A pharmaceutical composition according to claim 45 wherein R¹¹ and R¹² are independently selected from hydrogen, C₁-C₂₀-alkyl, aryl or aryl-C₁-C₆-alkyl, wherein the alkyl groups may optionally be substituted with one or more substituents independently selected from R¹⁵, and the aryl groups may optionally be substituted one or more substituents independently selected from R¹⁶.

25 47. A pharmaceutical composition according to claim 46 wherein R¹¹ and R¹² are independently selected from phenyl or phenyl-C₁-C₆-alkyl.

48. A pharmaceutical composition according to claim 46 wherein one or both of R¹¹ and R¹² are methyl.

30 49. A pharmaceutical composition according to any one of the claims 2 to 48 wherein R¹³ is independently selected from halogen, CF₃, OR¹¹ or NR¹¹R¹².

50. A pharmaceutical composition according to claim 49 wherein R¹³ is independently selected from halogen or OR¹¹.

51. A pharmaceutical composition according to claim 50 wherein R¹³ is OR¹¹.

52. A pharmaceutical composition according to any one of the claims 2 to 51 wherein R^{14} is independently selected from halogen, $-C(O)OR^{11}$, $-CN$, $-CF_3$, $-OR^{11}$, $S(O)_2R^{11}$, and C_1 - C_6 -alkyl.

53. A pharmaceutical composition according to claim 52 wherein R^{14} is independently selected from halogen, $-C(O)OR^{11}$, or $-OR^{11}$.


54. A pharmaceutical composition according to any one of the claims 2 to 53 wherein R^{15} is independently selected from halogen, $-CN$, $-CF_3$, $-C(O)OC_1-C_6$ -alkyl, and $-COOH$.

55. A pharmaceutical composition according to claim 54 wherein R^{15} is independently selected from halogen or $-C(O)OC_1-C_6$ -alkyl.

56. A pharmaceutical composition according to any one of the claims 2 to 55 wherein R^{16} is independently selected from halogen, $-C(O)OC_1-C_6$ -alkyl, $-COOH$, $-NO_2$, $-OC_1-C_6$ -alkyl, $-NH_2$, $C(=O)$ or C_1-C_6 -alkyl.

57. A pharmaceutical composition according to claim 56 wherein R^{16} is independently selected from halogen, $-C(O)OC_1-C_6$ -alkyl, $-COOH$, $-NO_2$, or C_1-C_6 -alkyl.

58. A pharmaceutical composition according to claim 1 wherein the zinc-binding ligand is



wherein

R^{19} is hydrogen or C_1 - C_6 -alkyl,

R^{20} is hydrogen or C_1 - C_6 -alkyl,

D , D^1 and F are a valence bond, C_1 - C_6 -alkylene or C_1 - C_6 -alkenylene optionally substituted with one or more substituents independently selected from R^{72} ,

R^{72} is independently selected from hydroxy, C_1 - C_6 -alkyl, or aryl,

E is C_1 - C_6 -alkyl, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with up to three substituents R^{21} , R^{22} and R^{23} ,

G and G¹ are C₁-C₆-alkyl, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with up to three substituents R²⁴, R²⁵ and R²⁶,

R¹⁷, R¹⁸, R²¹, R²², R²³, R²⁴, R²⁵ and R²⁶ are independently selected from

5

• hydrogen, halogen, -CN, -CH₂CN, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -S(O)₂CF₃, -SCF₃, -NO₂, =O, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷S(O)₂R²⁸, -S(O)₂NR²⁷R²⁸, -S(O)NR²⁷R²⁸, -S(O)R²⁷, -S(O)₂R²⁷, -C(O)NR²⁷R²⁸, -OC(O)NR²⁷R²⁸, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -CH₂C(O)NR²⁷R²⁸, -OCH₂C(O)NR²⁷R²⁸, -CH₂OR²⁷, -CH₂NR²⁷R²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -NR²⁷-C(=O)-C₁-C₆-alkyl-C(=O)OR²⁷, -NR²⁷-C(=O)-C₁-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

10

• C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl,

which may optionally be substituted with one or more substituents independently selected from R²⁹,

15

• aryl, aryloxy, aryloxycarbonyl, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl or heteroaryl-C₂-C₆-alkynyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰,

20

R²⁷ and R²⁸ are independently selected from hydrogen, C₁-C₆-alkyl, aryl-C₁-C₆-alkyl or aryl, or R²⁷ and R²⁸ when attached to the same nitrogen atom together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds,

25

R²⁹ is independently selected from halogen, -CN, -CF₃, -OCF₃, -OR²⁷, and -NR²⁷R²⁸,

30

R^{30} is independently selected from halogen, $-C(O)OR^{27}$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OR^{27}$, $-NR^{27}R^{28}$ and C_1 - C_6 -alkyl, or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.

59. A pharmaceutical composition according to claim 58 wherein D is a valence bond.

5 60. A pharmaceutical composition according to claim 58 wherein D is C_1 - C_6 -alkylene optionally substituted with one or more hydroxy, C_1 - C_6 -alkyl, or aryl.


61. A pharmaceutical composition according to any one of the claims 58 to 60 wherein E is aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with up to three substituents independently selected from R^{21} , R^{22} and R^{23} .

10 62. A pharmaceutical composition according to claim 61 wherein E is aryl optionally substituted with up to three substituents independently selected from R^{21} , R^{22} and R^{23} .

63. A pharmaceutical composition according to claim 62 wherein E is selected from ArG1 and optionally substituted with up to three substituents independently selected from R^{21} , R^{22} and R^{23} .

15 64. A pharmaceutical composition according to claim 63 wherein E is phenyl optionally substituted with up to three substituents independently selected from R^{21} , R^{22} and R^{23} .

65. A pharmaceutical composition according to claim 64 wherein the zinc-binding ligand is



20 66. A pharmaceutical composition according to any one of the claims 58 to 65 wherein R^{21} , R^{22} and R^{23} are independently selected from

25 •hydrogen, halogen, $-CHF_2$, $-CF_3$, $-OCF_3$, $-OCHF_2$, $-OCH_2CF_3$, $-OCF_2CHF_2$, $-SCF_3$, $-NO_2$, $-OR^{27}$, $-NR^{27}R^{28}$, $-SR^{27}$, $-C(O)NR^{27}R^{28}$, $-OC(O)NR^{27}R^{28}$, $-NR^{27}C(O)R^{28}$, $-NR^{27}C(O)OR^{28}$, $-CH_2C(O)NR^{27}R^{28}$, $-OCH_2C(O)NR^{27}R^{28}$, $-CH_2OR^{27}$, $-CH_2NR^{27}R^{28}$, $-OC(O)R^{27}$, $-OC_1-C_6$ -alkyl- $C(O)OR^{27}$, $-SC_1-C_6$ -alkyl- $C(O)OR^{27}$, $-C_2-C_6$ -alkenyl- $C(=O)OR^{27}$, $-NR^{27}-C(=O)-C_1-C_6$ -alkyl- $C(=O)OR^{27}$, $-NR^{27}-C(=O)-C_1-C_6$ -alkenyl- $C(=O)OR^{27}$, $-C(=O)NR^{27}-C_1-C_6$ -alkyl- $C(=O)OR^{27}$, $-C_1-C_6$ -alkyl- $C(=O)OR^{27}$, or $-C(O)OR^{27}$,

30

• C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl or C_2 - C_6 -alkynyl,

which may optionally be substituted with one or more substituents independently selected from R²⁹

• aryl, aryloxy, aryloxy carbonyl, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl or heteroaryl-C₂-C₆-alkynyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

67. A pharmaceutical composition according to claim 66 wherein R²¹, R²² and R²³ are independently selected from

• hydrogen, halogen, -OCF₃, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

• C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R²⁹

• aryl, aryloxy, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, heteroaryl, heteroaryl-C₁-C₆-alkyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

68. A pharmaceutical composition according to claim 67 wherein R²¹, R²² and R²³ are independently selected from

• hydrogen, halogen, -OCF₃, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

•methyl, ethyl propyl optionally substituted with one or more substituents independently selected from R²⁹

•aryl, aryloxy, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, heteroaryl, heteroaryl-C₁-C₆-alkyl

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

69. A pharmaceutical composition according to claim 68 wherein R²¹, R²² and R²³ are independently selected from

•hydrogen, halogen, -OCF₃, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

•methyl, ethyl propyl optionally substituted with one or more substituents independently selected from R²⁹

•ArG1, ArG1-O-, ArG1-C(O)-, ArG1-C₁-C₆-alkoxy, ArG1-C₁-C₆-alkyl, Het3, Het3-C₁-C₆-alkyl

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

70. A pharmaceutical composition according to claim 69 wherein R²¹, R²² and R²³ are independently selected from

•hydrogen, halogen, -OCF₃, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

•C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R²⁹

•phenyl, phenyloxy, phenyl-C₁-C₆-alkoxy, phenyl-C₁-C₆-alkyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

71. A pharmaceutical composition according to any one of the claims 58 to 70 wherein R¹⁹ is hydrogen or methyl.
72. A pharmaceutical composition according to claim 71 wherein R¹⁹ is hydrogen.
73. A pharmaceutical composition according to any one of the claims 58 to 72 wherein R²⁷ is Hydrogen, C₁-C₆-alkyl or aryl.
74. A pharmaceutical composition according to claim 73 wherein R²⁷ is hydrogen or C₁-C₆-alkyl.
75. A pharmaceutical composition according to any one of the claims 58 to 74 wherein R²⁸ is hydrogen or C₁-C₆-alkyl.
76. A pharmaceutical composition according to claim 58 wherein F is a valence bond.
77. A pharmaceutical composition according to claim 58 wherein F is C₁-C₆-alkylene optionally substituted with one or more hydroxy, C₁-C₆-alkyl, or aryl.
78. A pharmaceutical composition according to any one of the claims 58 or 76 to 77 wherein G is C₁-C₆-alkyl or aryl, wherein the aryl is optionally substituted with up to three substituents R²⁴, R²⁵ and R²⁶.
79. A pharmaceutical composition according to any one of the claims 58 or 76 to 77 wherein G is C₁-C₆-alkyl or ArG1, wherein the aryl is optionally substituted with up to three substituents R²⁴, R²⁵ and R²⁶.
80. A pharmaceutical composition according to claim 78 wherein G is C₁-C₆-alkyl.
81. A pharmaceutical composition according to claim 80 wherein G is phenyl optionally substituted with up to three substituents R²⁴, R²⁵ and R²⁶.
82. A pharmaceutical composition according to any one of the claims 58 to 81 wherein R²⁴, R²⁵ and R²⁶ are independently selected from

• hydrogen, halogen, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -SCF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -C(O)NR²⁷R²⁸, -OC(O)NR²⁷R²⁸, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -CH₂C(O)NR²⁷R²⁸, -OCH₂C(O)NR²⁷R²⁸, -CH₂OR²⁷, -CH₂NR²⁷R²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -NR²⁷-C(=O)-C₁-C₆-alkyl-C(=O)OR²⁷, -NR²⁷-C(=O)-C₁-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

- C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl,

which may optionally be substituted with one or more substituents independently selected from R²⁹

5

- aryl, aryloxy, aryloxycarbonyl, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl or heteroaryl-C₂-C₆-alkynyl,

10

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

83. A pharmaceutical composition according to claim 82 wherein R²⁴, R²⁵ and R²⁶ are independently selected from

15

- hydrogen, halogen, -OCF₃, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -NR²⁷C(O)R²⁸, -NR²⁷C(O)OR²⁸, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-alkyl-C(O)OR²⁷, -C₂-C₆-alkenyl-C(=O)OR²⁷, -C(=O)NR²⁷-C₁-C₆-alkyl-C(=O)OR²⁷, -C₁-C₆-alkyl-C(=O)OR²⁷, or -C(O)OR²⁷,

20

- C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl,

which may optionally be substituted with one or more substituents independently selected from R²⁹

25

- aryl, aryloxy, aryloxycarbonyl, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl or heteroaryl-C₂-C₆-alkynyl,

30

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

84. A pharmaceutical composition according to claim 83 wherein R²⁴, R²⁵ and R²⁶ are independently selected from

35

• hydrogen, halogen, $-\text{OCF}_3$, $-\text{OR}^{27}$, $-\text{NR}^{27}\text{R}^{28}$, $-\text{SR}^{27}$, $-\text{NR}^{27}\text{C}(\text{O})\text{R}^{28}$, $-\text{NR}^{27}\text{C}(\text{O})\text{OR}^{28}$, $-\text{OC}(\text{O})\text{R}^{27}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{SC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}_2\text{-C}_6\text{-alkenyl-C}(\text{=O})\text{OR}^{27}$, $-\text{C}(\text{=O})\text{NR}^{27}\text{-C}_1\text{-C}_6\text{-alkyl-C}(\text{=O})\text{OR}^{27}$, $-\text{C}_1\text{-C}_6\text{-alkyl-C}(\text{=O})\text{OR}^{27}$, or $-\text{C}(\text{O})\text{OR}^{27}$,

5

• $\text{C}_1\text{-C}_6\text{-alkyl}$ optionally substituted with one or more substituents independently selected from R^{29}

• aryl, aryloxy, aroyl, aryl- $\text{C}_1\text{-C}_6\text{-alkoxy}$, aryl- $\text{C}_1\text{-C}_6\text{-alkyl}$, heteroaryl, heteroaryl- $\text{C}_1\text{-C}_6\text{-alkyl}$,

10

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R^{30} .

15 85. A pharmaceutical composition according to claim 84 wherein R^{21} , R^{22} and R^{23} are independently selected from

• hydrogen, halogen, $-\text{OCF}_3$, $-\text{OR}^{27}$, $-\text{NR}^{27}\text{R}^{28}$, $-\text{SR}^{27}$, $-\text{NR}^{27}\text{C}(\text{O})\text{R}^{28}$, $-\text{NR}^{27}\text{C}(\text{O})\text{OR}^{28}$, $-\text{OC}(\text{O})\text{R}^{27}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{SC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}_2\text{-C}_6\text{-alkenyl-C}(\text{=O})\text{OR}^{27}$, $-\text{C}(\text{=O})\text{NR}^{27}\text{-C}_1\text{-C}_6\text{-alkyl-C}(\text{=O})\text{OR}^{27}$, $-\text{C}_1\text{-C}_6\text{-alkyl-C}(\text{=O})\text{OR}^{27}$, or $-\text{C}(\text{O})\text{OR}^{27}$,

20

• methyl, ethyl propyl optionally substituted with one or more substituents independently selected from R^{29}

25

• ArG1 , ArG1-O- , $\text{ArG1-C}(\text{O})\text{-}$, $\text{ArG1-C}_1\text{-C}_6\text{-alkoxy}$, $\text{ArG1-C}_1\text{-C}_6\text{-alkyl}$, Het3 , $\text{Het3-C}_1\text{-C}_6\text{-alkyl}$

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R^{30} .

30 86. A pharmaceutical composition according to claim 85 wherein R^{21} , R^{22} and R^{23} are independently selected from

• hydrogen, halogen, $-\text{OCF}_3$, $-\text{OR}^{27}$, $-\text{NR}^{27}\text{R}^{28}$, $-\text{SR}^{27}$, $-\text{NR}^{27}\text{C}(\text{O})\text{R}^{28}$, $-\text{NR}^{27}\text{C}(\text{O})\text{OR}^{28}$, $-\text{OC}(\text{O})\text{R}^{27}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{SC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{27}$, $-\text{C}_2\text{-C}_6\text{-alkenyl-}$

$C(=O)OR^{27}$, $-C(=O)NR^{27}-C_1-C_6\text{-alkyl}-C(=O)OR^{27}$, $-C_1-C_6\text{-alkyl}-C(=O)OR^{27}$, or $-C(O)OR^{27}$,

• methyl, ethyl propyl optionally substituted with one or more substituents independently selected from R^{29}

• ArG1, ArG1-O-, ArG1-C(O)-, ArG1-C₁-C₆-alkoxy, ArG1-C₁-C₆-alkyl, Het3, Het3-C₁-C₆-alkyl

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R^{30} .

87. A pharmaceutical composition according to claim 86 wherein R^{21} , R^{22} and R^{23} are independently selected from

• hydrogen, halogen, $-OCF_3$, $-OR^{27}$, $-NR^{27}R^{28}$, $-SR^{27}$, $-NR^{27}C(O)R^{28}$, $-NR^{27}C(O)OR^{28}$, $-OC(O)R^{27}$, $-OC_1-C_6\text{-alkyl}-C(O)OR^{27}$, $-SC_1-C_6\text{-alkyl}-C(O)OR^{27}$, $-C_2-C_6\text{-alkenyl}-C(=O)OR^{27}$, $-C(=O)NR^{27}-C_1-C_6\text{-alkyl}-C(=O)OR^{27}$, $-C_1-C_6\text{-alkyl}-C(=O)OR^{27}$, or $-C(O)OR^{27}$,

• methyl, ethyl propyl optionally substituted with one or more substituents independently selected from R^{29}

• ArG1, ArG1-O-, ArG1-C₁-C₆-alkoxy, ArG1-C₁-C₆-alkyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R^{30} .

88. A pharmaceutical composition according to any one of the claims 58 or 76 to 87 wherein R^{20} is hydrogen or methyl.

89. A pharmaceutical composition according to claim 88 wherein R^{20} is hydrogen.

90. A pharmaceutical composition according to any one of the claims 58 or 76 to 89 wherein R^{27} is hydrogen, C₁-C₆-alkyl or aryl.

91. A pharmaceutical composition according to claim 90 wherein R^{27} is hydrogen or C₁-C₆-alkyl or ArG1.

92. A pharmaceutical composition according to claim 91 wherein R^{27} is hydrogen or C₁-C₆-alkyl.

93. A pharmaceutical composition according to any one of the claims 58 or 76 to 91 wherein R^{28} is hydrogen or C₁-C₆-alkyl.

94. A pharmaceutical composition according to claim 58 wherein R¹⁷ and R¹⁸ are independently selected from

• hydrogen, halogen, -CN, -CF₃, -OCF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, -SR²⁷, -S(O)R²⁷,
-S(O)₂R²⁷, -C(O)NR²⁷R²⁸, -CH₂OR²⁷, -OC(O)R²⁷, -OC₁-C₆-alkyl-C(O)OR²⁷, -SC₁-C₆-
alkyl-C(O)OR²⁷, or -C(O)OR²⁷,

• C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl, optionally substituted with one or more
substituents independently selected from R²⁹

• aryl, aryloxy, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, heteroaryl, heteroaryl-C₁-C₆-
alkyl,

of which the cyclic moieties optionally may be substituted with one or more
substituents selected from R³⁰.

95. A pharmaceutical composition according to claim 94 wherein R¹⁷ and R¹⁸ are independently selected from

• hydrogen, halogen, -CN, -CF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, or -C(O)OR²⁷,

• C₁-C₆-alkyl optionally substituted with one or more substituents independently
selected from R²⁹

• aryl, aryloxy, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, heteroaryl, heteroaryl-C₁-C₆-
alkyl,

of which the cyclic moieties optionally may be substituted with one or more
substituents selected from R³⁰.

96. A pharmaceutical composition according to claim 95 wherein R¹⁷ and R¹⁸ are independently selected from

• hydrogen, halogen, -CN, -CF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, or -C(O)OR²⁷

• methyl, ethyl propyl optionally substituted with one or more substituents
independently selected from R²⁹

- aryl, aryloxy, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, heteroaryl, heteroaryl-C₁-C₆-alkyl

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

- 5 97. A pharmaceutical composition according to claim 96 wherein R¹⁷ and R¹⁸ are independently selected from

- hydrogen, halogen, -CN, -CF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, or -C(O)OR²⁷
- methyl, ethyl propyl optionally substituted with one or more substituents independently selected from R²⁹

- 10 • ArG1, ArG1-O-, ArG1-C(O)-, ArG1-C₁-C₆-alkoxy, ArG1-C₁-C₆-alkyl, Het3, Het3-C₁-C₆-alkyl

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

- 15 98. A pharmaceutical composition according to claim 97 wherein R¹⁷ and R¹⁸ are independently selected from

- hydrogen, halogen, -CN, -CF₃, -NO₂, -OR²⁷, -NR²⁷R²⁸, or -C(O)OR²⁷
- C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R²⁹
- phenyl, phenyloxy, phenyl-C₁-C₆-alkoxy, phenyl-C₁-C₆-alkyl,

- 20 of which the cyclic moieties optionally may be substituted with one or more substituents selected from R³⁰.

99. A pharmaceutical composition according to any one of the claims 58 to 98 wherein R²⁷ is hydrogen or C₁-C₆-alkyl.


- 25 100. A pharmaceutical composition according to claim 99 wherein R²⁷ is hydrogen, methyl or ethyl.

101. A pharmaceutical composition according to any one of the claims 58 to 100 wherein R²⁸ is hydrogen or C₁-C₆-alkyl.

102. A pharmaceutical composition according to claim 101 wherein R²⁸ is hydrogen, methyl or ethyl.

- 30 103. A pharmaceutical composition according to any one of the claims 58 to 102 wherein R⁷² is -OH or phenyl.


104. A pharmaceutical composition according to claim 58 wherein the zinc-binding ligand is



105. A pharmaceutical composition according to claim 1 wherein the zinc-binding ligand is of the form H-I-J

5

wherein H is




wherein the phenyl, naphthalene or benzocarbazole rings are optionally substituted with one or more substituents independently selected from R^{31}

15

I is selected from

- a valence bond,
- $-\text{CH}_2\text{N}(\text{R}^{32})-$ or $-\text{SO}_2\text{N}(\text{R}^{33})-$,



- $-\text{Z}^1-\text{N}-\text{CH}_2-\text{CH}_2-\text{N}-\text{Z}^2-$ wherein Z^1 is $\text{S}(\text{O})_2$ or CH_2 , Z^2 is $-\text{NH}-$, $-\text{O}-$ or $-\text{S}-$, and n is 1 or 2,

20

J is

- C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl or C_2 - C_6 -alkynyl, which may each optionally be substituted with one or more substituents selected from R^{34} ,
- Aryl, aryloxy, aryl-oxycarbonyl-, aroyl, aryl- C_1 - C_6 -alkoxy-, aryl- C_1 - C_6 -alkyl-, aryl- C_2 - C_6 -alkenyl-, aryl- C_2 - C_6 -alkynyl-, heteroaryl, heteroaryl- C_1 - C_6 -alkyl-, heteroaryl- C_2 - C_6 -alkenyl- or heteroaryl- C_2 - C_6 -alkynyl-, wherein the cyclic moieties are optionally substituted with one or more substituents selected from R^{37} ,
- Hydrogen,

25

30

R^{31} is independently selected from hydrogen, halogen, $-\text{CN}$, $-\text{CH}_2\text{CN}$, $-\text{CHF}_2$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OCHF}_2$, $-\text{OCH}_2\text{CF}_3$, $-\text{OCF}_2\text{CHF}_2$, $-\text{S}(\text{O})_2\text{CF}_3$, $-\text{SCF}_3$, $-\text{NO}_2$, $-\text{OR}^{35}$, $-\text{C}(\text{O})\text{R}^{35}$, $-\text{NR}^{35}\text{R}^{36}$, $-\text{SR}^{35}$,

$-\text{NR}^{35}\text{S}(\text{O})_2\text{R}^{36}$, $-\text{S}(\text{O})_2\text{NR}^{35}\text{R}^{36}$, $-\text{S}(\text{O})\text{NR}^{35}\text{R}^{36}$, $-\text{S}(\text{O})\text{R}^{35}$, $-\text{S}(\text{O})_2\text{R}^{35}$, $-\text{C}(\text{O})\text{NR}^{35}\text{R}^{36}$,
 $-\text{OC}(\text{O})\text{NR}^{35}\text{R}^{36}$, $-\text{NR}^{35}\text{C}(\text{O})\text{R}^{36}$, $-\text{CH}_2\text{C}(\text{O})\text{NR}^{35}\text{R}^{36}$, $-\text{OCH}_2\text{C}(\text{O})\text{NR}^{35}\text{R}^{36}$, $-\text{CH}_2\text{OR}^{35}$,
 $-\text{CH}_2\text{NR}^{35}\text{R}^{36}$, $-\text{OC}(\text{O})\text{R}^{35}$, $-\text{OC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{35}$, $-\text{SC}_1\text{-C}_6\text{-alkyl-C}(\text{O})\text{OR}^{35}$, $-\text{C}_2\text{-C}_6\text{-alkenyl-}$
 $\text{C}(\text{=O})\text{OR}^{35}$, $-\text{NR}^{35}\text{-C}(\text{=O})\text{-C}_1\text{-C}_6\text{-alkyl-C}(\text{=O})\text{OR}^{35}$, $-\text{NR}^{35}\text{-C}(\text{=O})\text{-C}_1\text{-C}_6\text{-alkenyl-C}(\text{=O})\text{OR}^{35}$,
 5 $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_1\text{-C}_6\text{-alkanoyl}$ or $-\text{C}(\text{O})\text{OR}^{35}$,

R^{32} and R^{33} are independently selected from hydrogen, $\text{C}_1\text{-C}_6\text{-alkyl}$ or $\text{C}_1\text{-C}_6\text{-alkanoyl}$,

R^{34} is independently selected from halogen, $-\text{CN}$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OR}^{35}$, and $-\text{NR}^{35}\text{R}^{36}$,


10

R^{35} and R^{36} are independently selected from hydrogen, $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{aryl-C}_1\text{-C}_6\text{-alkyl}$ or aryl , or
 R^{35} and R^{36} when attached to the same nitrogen atom together with the said nitrogen atom
 may form a 3 to 8 membered heterocyclic ring optionally containing one or two further
 heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or
 15 two double bonds,

R^{37} is independently selected from halogen, $-\text{C}(\text{O})\text{OR}^{35}$, $-\text{C}(\text{O})\text{H}$, $-\text{CN}$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{NO}_2$, $-\text{OR}^{35}$, $-\text{NR}^{35}\text{R}^{36}$, $\text{C}_1\text{-C}_6\text{-alkyl}$ or $\text{C}_1\text{-C}_6\text{-alkanoyl}$,

20 or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.


106. A pharmaceutical composition according to claim 105 wherein the zinc-binding ligand is of the form H-I-J, wherein H is



25 wherein the phenyl, naphthalene or benzocarbazole rings are optionally substituted with one or more substituents independently selected from R^{31} ,

I is selected from

- a valence bond,
- 30 • $-\text{CH}_2\text{N}(\text{R}^{32})-$ or $-\text{SO}_2\text{N}(\text{R}^{33})-$,



wherein Z¹ is S(O)₂ or CH₂, Z² is N, -O- or -S-, and n is 1 or 2,

J is

- C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl, which may each optionally be substituted with one or more substituents selected from R³⁴,
- Aryl, aryloxy, aryl-oxycarbonyl-, aroyl, aryl-C₁-C₆-alkoxy-, aryl-C₁-C₆-alkyl-, aryl-C₂-C₆-alkenyl-, aryl-C₂-C₆-alkynyl-, heteroaryl, heteroaryl-C₁-C₆-alkyl-, heteroaryl-C₂-C₆-alkenyl- or heteroaryl-C₂-C₆-alkynyl-, wherein the cyclic moieties are optionally substituted with one or more substituents selected from R³⁷,
- hydrogen,

R³¹ is independently selected from hydrogen, halogen, -CN, -CH₂CN, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -S(O)₂CF₃, -SCF₃, -NO₂, -OR³⁵, -C(O)R³⁵, -NR³⁵R³⁶, -SR³⁵, -NR³⁵S(O)₂R³⁶, -S(O)₂NR³⁵R³⁶, -S(O)NR³⁵R³⁶, -S(O)R³⁵, -S(O)₂R³⁵, -C(O)NR³⁵R³⁶, -OC(O)NR³⁵R³⁶, -NR³⁵C(O)R³⁶, -CH₂C(O)NR³⁵R³⁶, -OCH₂C(O)NR³⁵R³⁶, -CH₂OR³⁵, -CH₂NR³⁵R³⁶, -OC(O)R³⁵, -OC₁-C₆-alkyl-C(O)OR³⁵, -SC₁-C₆-alkyl-C(O)OR³⁵, -C₂-C₆-alkenyl-C(=O)OR³⁵, -NR³⁵-C(=O)-C₁-C₆-alkyl-C(=O)OR³⁵, -NR³⁵-C(=O)-C₁-C₆-alkenyl-C(=O)OR³⁵, C₁-C₆-alkyl, C₁-C₆-alkanoyl or -C(O)OR³⁵,

R³² and R³³ are independently selected from hydrogen, C₁-C₆-alkyl or C₁-C₆-alkanoyl,

R³⁴ is independently selected from halogen, -CN, -CF₃, -OCF₃, -OR³⁵, and -NR³⁵R³⁶,

R³⁵ and R³⁶ are independently selected from hydrogen, C₁-C₆-alkyl, aryl-C₁-C₆-alkyl or aryl, or


R³⁵ and R³⁶ when attached to the same nitrogen atom together with the said nitrogen atom may form a 3 to 8 membered heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds,

R³⁷ is independently selected from halogen, -C(O)OR³⁵, -C(O)H, -CN, -CF₃, -OCF₃, -NO₂, -OR³⁵, -NR³⁵R³⁶, C₁-C₆-alkyl or C₁-C₆-alkanoyl,


or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base,

With the proviso that R^{31} and J cannot both be hydrogen.


- 5 107. A pharmaceutical composition according to any one of the claims 105 or 106 wherein H is



108. A pharmaceutical composition according to claim 107 wherein H is



109. A pharmaceutical composition according to claim 107 wherein H is



10

110. A pharmaceutical composition according to any one of the claims 105 to 109 wherein I is a valence bond, $-\text{CH}_2\text{N}(\text{R}^{32})-$, or $-\text{SO}_2\text{N}(\text{R}^{33})-$.

111. A pharmaceutical composition according to claim 110 wherein I is a valence bond.

112. A pharmaceutical composition according to any one of the claims 105 to 111 wherein J is

- hydrogen,
- C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl or C_2 - C_6 -alkynyl,

which may optionally be substituted with one or more substituents selected from halogen, $-\text{CN}$, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OR}^{35}$, and $-\text{NR}^{35}\text{R}^{36}$,

- 20 • aryl, or heteroaryl, wherein the cyclic moieties are optionally substituted with one or more substituents independently selected from R^{37} .

113. A pharmaceutical composition according to claim 112 wherein J is

- hydrogen,

- aryl or heteroaryl, wherein the cyclic moieties are optionally substituted with one or more substituents independently selected from R³⁷.

114. A pharmaceutical composition according to claim 112 wherein J is

- hydrogen,
- ArG1 or Het3, wherein the cyclic moieties are optionally substituted with one or more substituents independently selected from R³⁷.

115. A pharmaceutical composition according to claim 114 wherein J is

- hydrogen,
- phenyl or naphthyl optionally substituted with one or more substituents independently selected from R³⁷.

116. A pharmaceutical composition according to claim 115 wherein J is hydrogen.

117. A pharmaceutical composition according to any one of the claims 105 to 116 wherein R³² and R³³ are independently selected from hydrogen or C₁-C₆-alkyl.

118. A pharmaceutical composition according to any one of the claims 105 to 117 wherein R³⁴ is hydrogen, halogen, -CN, -CF₃, -OCF₃, -SCF₃, -NO₂, -OR³⁵, -C(O)R³⁵, -NR³⁵R³⁶, -SR³⁵, -C(O)NR³⁵R³⁶, -OC(O)NR³⁵R³⁶, -NR³⁵C(O)R³⁶, -OC(O)R³⁵, -OC₁-C₆-alkyl-C(O)OR³⁵, -SC₁-C₆-alkyl-C(O)OR³⁵ or -C(O)OR³⁵.

119. A pharmaceutical composition according to claim 118 wherein R³⁴ is hydrogen, halogen, -CF₃, -NO₂, -OR³⁵, -NR³⁵R³⁶, -SR³⁵, -NR³⁵C(O)R³⁶, or -C(O)OR³⁵.

120. A pharmaceutical composition according to claim 119 wherein R³⁴ is hydrogen, halogen, -CF₃, -NO₂, -OR³⁵, -NR³⁵R³⁶, or -NR³⁵C(O)R³⁶.

121. A pharmaceutical composition according to claim 120 wherein R³⁴ is hydrogen, halogen, or -OR³⁵.

122. A pharmaceutical composition according to any one of the claims 105 to 121 wherein R³⁵ and R³⁶ are independently selected from hydrogen, C₁-C₆-alkyl, or aryl.


123. A pharmaceutical composition according to claim 122 wherein R³⁵ and R³⁶ are independently selected from hydrogen or C₁-C₆-alkyl.

124. A pharmaceutical composition according to any one of the claims 105 to 123 wherein R³⁷ is halogen, -C(O)OR³⁵, -CN, -CF₃, -OR³⁵, -NR³⁵R³⁶, C₁-C₆-alkyl or C₁-C₆-alkanoyl.

125. A pharmaceutical composition according to claim 124 wherein R³⁷ is halogen, -C(O)OR³⁵, -OR³⁵, -NR³⁵R³⁶, C₁-C₆-alkyl or C₁-C₆-alkanoyl.

126. A pharmaceutical composition according to claim 125 wherein R³⁷ is halogen, -C(O)OR³⁵ or -OR³⁵.

127. A pharmaceutical composition according to claim 1 wherein the zinc-binding ligand is



wherein K is a valence bond, C₁-C₆-alkylene, -NH-C(=O)-U-, -C₁-C₆-alkyl-S-, -C₁-C₆-alkyl-O-, -C(=O)-, or -C(=O)-NH-, wherein any C₁-C₆-alkyl moiety is optionally substituted with R³⁸,

5

U is a valence bond, C₁-C₆-alkenylene, -C₁-C₆-alkyl-O- or C₁-C₆-alkylene wherein any C₁-C₆-alkyl moiety is optionally substituted with C₁-C₆-alkyl,

10 R³⁸ is C₁-C₆-alkyl, aryl, wherein the alkyl or aryl moieties are optionally substituted with one or more substituents independently selected from R³⁹,

R³⁹ is independently selected from halogen, cyano, nitro, amino,

15 M is a valence bond, arylene or heteroarylene, wherein the aryl or heteroaryl moieties are optionally substituted with one or more substituents independently selected from R⁴⁰,

R⁴⁰ is selected from

20 •hydrogen, halogen, -CN, -CH₂CN, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -S(O)₂CF₃, -OS(O)₂CF₃, -SCF₃, -NO₂, -OR⁴¹, -NR⁴¹R⁴², -SR⁴¹, -NR⁴¹S(O)₂R⁴², -S(O)₂NR⁴¹R⁴², -S(O)NR⁴¹R⁴², -S(O)R⁴¹, -S(O)₂R⁴¹, -OS(O)₂R⁴¹, -C(O)NR⁴¹R⁴², -OC(O)NR⁴¹R⁴², -NR⁴¹C(O)R⁴², -CH₂C(O)NR⁴¹R⁴², -OC₁-C₆-alkyl-C(O)NR⁴¹R⁴², -CH₂OR⁴¹, -CH₂OC(O)R⁴¹, -CH₂NR⁴¹R⁴², -OC(O)R⁴¹, -OC₁-C₆-alkyl-C(O)OR⁴¹, -OC₁-C₆-alkyl-OR⁴¹, -S-C₁-C₆-alkyl-C(O)OR⁴¹, -C₂-C₆-alkenyl-C(=O)OR⁴¹, -NR⁴¹-C(=O)-C₁-C₆-alkyl-C(=O)OR⁴¹, -NR⁴¹-C(=O)-C₁-C₆-alkenyl-C(=O)OR⁴¹, -C(O)OR⁴¹, -C₂-C₆-alkenyl-C(=O)R⁴¹, =O, -NH-C(=O)-O-C₁-C₆-alkyl, or -NH-C(=O)-C(=O)-O-C₁-C₆-alkyl,

25

•C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl, which may each optionally be substituted with one or more substituents selected from R⁴³,

30

•aryl, aryloxy, aryloxycarbonyl, aroyl, arylsulfanyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aroyl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl or heteroaryl-C₂-C₆-alkynyl, wherein the cyclic

moieties optionally may be substituted with one or more substituents selected from R^{44} ,

- 5 R^{41} and R^{42} are independently selected from hydrogen, -OH, C_1 - C_6 -alkyl, C_1 - C_6 -alkenyl, aryl-
 C_1 - C_6 -alkyl or aryl, wherein the alkyl moieties may optionally be substituted with one or more
substituents independently selected from R^{45} , and the aryl moieties may optionally be
substituted with one or more substituents independently selected from R^{46} ; R^{41} and R^{42} when
attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the
said nitrogen atom, the heterocyclic ring optionally containing one or two further heteroatoms
10 selected from nitrogen, oxygen and sulphur, and optionally containing one or two double
bonds,

R^{43} is independently selected from halogen, -CN, -CF₃, -OCF₃, -OR⁴¹, and -NR⁴¹R⁴²

- 15 R^{44} is independently selected from halogen, -C(O)OR⁴¹, -CH₂C(O)OR⁴¹, -CH₂OR⁴¹, -CN, -
CF₃, -OCF₃, -NO₂, -OR⁴¹, -NR⁴¹R⁴² and C_1 - C_6 -alkyl,
 R^{45} is independently selected from halogen, -CN, -CF₃, -OCF₃, -O- C_1 - C_6 -alkyl, -C(O)-O- C_1 -
 C_6 -alkyl, -COOH and -NH₂,
 R^{46} is independently selected from halogen, -C(O)OC₁- C_6 -alkyl, -COOH, -CN, -CF₃, -OCF₃, -
20 NO₂, -OH, -OC₁- C_6 -alkyl, -NH₂, C(=O) or C_1 - C_6 -alkyl,

- Q is a valence bond, C_1 - C_6 -alkylene, - C_1 - C_6 -alkyl-O-, - C_1 - C_6 -alkyl-NH-, -NH- C_1 - C_6 -alkyl,
-NH-C(=O)-, -C(=O)-NH-, -O- C_1 - C_6 -alkyl, -C(=O)-, or - C_1 - C_6 -alkyl-C(=O)-N(R^{47})- wherein the
alkyl moieties are optionally substituted with one or more substituents independently selected
25 from R^{48} ,

R^{47} and R^{48} are independently selected from hydrogen, C_1 - C_6 -alkyl, aryl optionally substituted
with one or more R^{49} ,

- 30 R^{49} is independently selected from halogen and -COOH,

T is

- hydrogen,

• C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₆-alkyloxy-carbonyl, wherein the alkyl, alkenyl and alkynyl moieties are optionally substituted with one or more substituents independently selected from R⁵⁰,

5 • aryl, aryloxy, aryloxy-carbonyl, aryl-C₁-C₆-alkyl, aroyl, aryl-C₁-C₆-alkoxy, aryl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl, heteroaryl-C₂-C₆-alkynyl,

wherein any alkyl, alkenyl, alkynyl, aryl and heteroaryl moiety is optionally substituted with one or more substituents independently selected from R⁵⁰,

10

R⁵⁰ is C₁-C₆-alkyl, C₁-C₆-alkoxy, aryl, aryloxy, aryl-C₁-C₆-alkoxy, -C(=O)-NH-C₁-C₆-alkyl-aryl, -C(=O)-NR^{50A}-C₁-C₆-alkyl, -C(=O)-NH-(CH₂CH₂O)_m-C₁-C₆-alkyl-COOH, heteroaryl, heteroaryl-C₁-C₆-alkoxy, -C₁-C₆-alkyl-COOH, -O-C₁-C₆-alkyl-COOH, -S(O)₂R⁵¹, -C₂-C₆-alkenyl-COOH, -OR⁵¹, -NO₂, halogen, -COOH, -CF₃, -CN, =O, -N(R⁵¹R⁵²), wherein m is 1, 2, 3 or 4, and wherein the aryl or heteroaryl moieties are optionally substituted with one or more R⁵³, and the alkyl moieties are optionally substituted with one or more R^{50B}.

15

R^{50A} and R^{50B} are independently selected from -C(O)OC₁-C₆-alkyl, -COOH, -C₁-C₆-alkyl-C(O)OC₁-C₆-alkyl, -C₁-C₆-alkyl-COOH, or C₁-C₆-alkyl,

R⁵¹ and R⁵² are independently selected from hydrogen and C₁-C₆-alkyl,

20

R⁵³ is independently selected from C₁-C₆-alkyl, C₁-C₆-alkoxy, -C₁-C₆-alkyl-COOH, -C₂-C₆-alkenyl-COOH, -OR⁵¹, -NO₂, halogen, -COOH, -CF₃, -CN, or -N(R⁵¹R⁵²),

or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.

25

128. A pharmaceutical composition according to claim 127 wherein K is a valence bond, C₁-C₆-alkylene, -NH-C(=O)-U-, -C₁-C₆-alkyl-S-, -C₁-C₆-alkyl-O-, or -C(=O)-, wherein any C₁-C₆-alkyl moiety is optionally substituted with R³⁸.

129. A pharmaceutical composition according to claim 128 wherein K is a valence bond, C₁-C₆-alkylene, -NH-C(=O)-U-, -C₁-C₆-alkyl-S-, or -C₁-C₆-alkyl-O-, wherein any C₁-C₆-alkyl moiety is optionally substituted with R³⁸.

30

130. A pharmaceutical composition according to claim 129 wherein K is a valence bond, C₁-C₆-alkylene, or -NH-C(=O)-U-, wherein any C₁-C₆-alkyl moiety is optionally substituted with R³⁸.

131. A pharmaceutical composition according to claim 130 wherein K is a valence bond or C₁-C₆-alkylene, wherein any C₁-C₆-alkyl moiety is optionally substituted with R³⁸.

35

132. A pharmaceutical composition according to claim 130 wherein K is a valence bond or -NH-C(=O)-U.

133. A pharmaceutical composition according to claim 131 wherein K is a valence bond.

134. A pharmaceutical composition according to any one of the claims 127 to 133 wherein U

5 is a valence bond or -C₁-C₆-alkyl-O-.

135. A pharmaceutical composition according to claim 134 wherein U is a valence bond

136. A pharmaceutical composition according to any one of the claims 127 to 135 wherein M is arylene or heteroarylene, wherein the arylene or heteroarylene moieties are optionally substituted with one or more substituents independently selected from R⁴⁰.

10 137. A pharmaceutical composition according to claim 136 wherein M is ArG1 or Het1, wherein the arylene or heteroarylene moieties are optionally substituted with one or more substituents independently selected from R⁴⁰.


138. A pharmaceutical composition according to claim 137 wherein M is ArG1 or Het2, wherein the arylene or heteroarylene moieties are optionally substituted with one or more
15 substituents independently selected from R⁴⁰.

139. A pharmaceutical composition according to claim 138 wherein M is ArG1 or Het3, wherein the arylene or heteroarylene moieties are optionally substituted with one or more substituents independently selected from R⁴⁰.

140. A pharmaceutical composition according to claim 139 wherein M is phenylene optionally
20 substituted with one or more substituents independently selected from R⁴⁰.


141. A pharmaceutical composition according to claim 139 wherein M is indolyne optionally substituted with one or more substituents independently selected from R⁴⁰.

142. A pharmaceutical composition according to claim 141 wherein M is



25 143. A pharmaceutical composition according to claim 139 wherein M is carbazolyne optionally substituted with one or more substituents independently selected from R⁴⁰.

144. A pharmaceutical composition according to claim 143 wherein M is



145. A pharmaceutical composition according to any one of the claims 127 to 144 wherein

30 R⁴⁰ is selected from

• hydrogen, halogen, -CN, -CF₃, -OCF₃, -NO₂, -OR⁴¹, -NR⁴¹R⁴², -SR⁴¹, -S(O)₂R⁴¹, -NR⁴¹C(O)R⁴², -OC₁-C₆-alkyl-C(O)NR⁴¹R⁴², -C₂-C₆-alkenyl-C(=O)OR⁴¹, -C(O)OR⁴¹, =O, -NH-C(=O)-O-C₁-C₆-alkyl, or -NH-C(=O)-C(=O)-O-C₁-C₆-alkyl,

5 C₁-C₆-alkyl or C₂-C₆-alkenyl which may each optionally be substituted with one or more substituents independently selected from R⁴³,

• aryl, aryloxy, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, or heteroaryl-C₂-C₆-alkenyl, wherein the cyclic moieties
10 optionally may be substituted with one or more substituents selected from R⁴⁴.

146. A pharmaceutical composition according to claim 145 wherein R⁴⁰ is selected from

• hydrogen, halogen, -CN, -CF₃, -OCF₃, -NO₂, -OR⁴¹, -NR⁴¹R⁴², -SR⁴¹, -S(O)₂R⁴¹, -NR⁴¹C(O)R⁴², -OC₁-C₆-alkyl-C(O)NR⁴¹R⁴², -C₂-C₆-alkenyl-C(=O)OR⁴¹, -C(O)OR⁴¹, =O, -NH-C(=O)-O-C₁-C₆-alkyl, or -NH-C(=O)-C(=O)-O-C₁-C₆-alkyl,

15 C₁-C₆-alkyl or C₂-C₆-alkenyl which may each optionally be substituted with one or more substituents independently selected from R⁴³,

• ArG1, ArG1-O-, ArG1-C₁-C₆-alkoxy, ArG1-C₁-C₆-alkyl, ArG1-C₂-C₆-alkenyl, Het3, Het3-C₁-C₆-alkyl, or Het3-C₂-C₆-alkenyl, wherein the cyclic moieties optionally may be
20 substituted with one or more substituents selected from R⁴⁴.

147. A pharmaceutical composition according to claim 146 wherein R⁴⁰ is selected from

• hydrogen, halogen, -CF₃, -NO₂, -OR⁴¹, -NR⁴¹R⁴², -C(O)OR⁴¹, =O, or -NR⁴¹C(O)R⁴²,
• C₁-C₆-alkyl,
25 • ArG1.

148. A pharmaceutical composition according to claim 147 wherein R⁴⁰ is hydrogen.

149. A pharmaceutical composition according to claim 147 wherein R⁴⁰ is selected from

• Halogen, -NO₂, -OR⁴¹, -NR⁴¹R⁴², -C(O)OR⁴¹, or -NR⁴¹C(O)R⁴²,
30 • Methyl,
• Phenyl.

150. A pharmaceutical composition according to any one of the claims 127 to 149 wherein R⁴¹ and R⁴² are independently selected from hydrogen, C₁-C₆-alkyl, or aryl, wherein the aryl moieties may optionally be substituted with halogen or -COOH.

151. A pharmaceutical composition according to claim 150 wherein R⁴¹ and R⁴² are independently selected from hydrogen, methyl, ethyl, or phenyl, wherein the phenyl moieties may optionally be substituted with halogen or -COOH.

5 152. A pharmaceutical composition according to any one of the claims 127 to 151 wherein Q is a valence bond, C₁-C₆-alkylene, -C₁-C₆-alkyl-O-, -C₁-C₆-alkyl-NH-, -NH-C₁-C₆-alkyl, -NH-C(=O)-, -C(=O)-NH-, -O-C₁-C₆-alkyl, -C(=O)-, or -C₁-C₆-alkyl-C(=O)-N(R⁴⁷)- wherein the alkyl moieties are optionally substituted with one or more substituents independently selected from R⁴⁸.

10 153. A pharmaceutical composition according to claim 152 wherein Q is a valence bond, -CH₂-, -CH₂-CH₂-, -CH₂-O-, -CH₂-CH₂-O-, -CH₂-NH-, -CH₂-CH₂-NH-, -NH-CH₂-, -NH-CH₂-CH₂-, -NH-C(=O)-, -C(=O)-NH-, -O-CH₂-, -O-CH₂-CH₂-, or -C(=O)-.

154. A pharmaceutical composition according to any one of the claims 127 to 153 wherein R⁴⁷ and R⁴⁸ are independently selected from hydrogen, methyl and phenyl.

15 155. A pharmaceutical composition according to any one of the claims 127 to 154 wherein T is

- Hydrogen,
- C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R⁵⁰,
- aryl, aryl-C₁-C₆-alkyl, heteroaryl, wherein the alkyl, aryl and heteroaryl moieties are
- 20 optionally substituted with one or more substituents independently selected from R⁵⁰.

156. A pharmaceutical composition according to claim 155 wherein T is

- hydrogen,
- C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R⁵⁰,
- 25 • ArG1, ArG1-C₁-C₆-alkyl, Het3, wherein the alkyl, aryl and heteroaryl moieties are optionally substituted with one or more substituents independently selected from R⁵⁰.

157. A pharmaceutical composition according to claim 156 wherein T is

- hydrogen,
- 30 • C₁-C₆-alkyl, optionally substituted with one or more substituents independently selected from R⁵⁰,
- phenyl, phenyl-C₁-C₆-alkyl, wherein the alkyl and phenyl moieties are optionally substituted with one or more substituents independently selected from R⁵⁰.

35 158. A pharmaceutical composition according to claim 157 wherein T is phenyl substituted with R⁵⁰.

159. A pharmaceutical composition according to any one of the claims 127 to 158 wherein R^{50} is C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, aryl, aryloxy, aryl- C_1 - C_6 -alkoxy, $-C(=O)-NH-C_1$ - C_6 -alkyl-aryl, $-C(=O)-NR^{50A}-C_1$ - C_6 -alkyl, $-C(=O)-NH-(CH_2CH_2O)_mC_1$ - C_6 -alkyl-COOH, heteroaryl, $-C_1$ - C_6 -alkyl-COOH, $-O-C_1$ - C_6 -alkyl-COOH, $-S(O)_2R^{51}$, $-C_2$ - C_6 -alkenyl-COOH, $-OR^{51}$, $-NO_2$, halogen, $-COOH$, $-CF_3$, $-CN$, $=O$, $-N(R^{51}R^{52})$, wherein the aryl or heteroaryl moieties are optionally substituted with one or more R^{53} .

160. A pharmaceutical composition according to claim 159 wherein R^{50} is C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, aryl, aryloxy, $-C(=O)-NR^{50A}-C_1$ - C_6 -alkyl, $-C(=O)-NH-(CH_2CH_2O)_mC_1$ - C_6 -alkyl-COOH, aryl- C_1 - C_6 -alkoxy, $-OR^{51}$, $-NO_2$, halogen, $-COOH$, $-CF_3$, wherein any aryl moiety is optionally substituted with one or more R^{53} .

161. A pharmaceutical composition according to claim 160 wherein R^{50} is C_1 - C_6 -alkyl, aryloxy, $-C(=O)-NR^{50A}-C_1$ - C_6 -alkyl, $-C(=O)-NH-(CH_2CH_2O)_mC_1$ - C_6 -alkyl-COOH, aryl- C_1 - C_6 -alkoxy, $-OR^{51}$, halogen, $-COOH$, $-CF_3$, wherein any aryl moiety is optionally substituted with one or more R^{53} .

162. A pharmaceutical composition according to claim 161 wherein R^{50} is C_1 - C_6 -alkyl, ArG1-O-, $-C(=O)-NR^{50A}-C_1$ - C_6 -alkyl, $-C(=O)-NH-(CH_2CH_2O)_mC_1$ - C_6 -alkyl-COOH, ArG1- C_1 - C_6 -alkoxy, $-OR^{51}$, halogen, $-COOH$, $-CF_3$, wherein any aryl moiety is optionally substituted with one or more R^{53} .

163. A pharmaceutical composition according to claim 162 wherein R^{50} is $-C(=O)-NR^{50A}CH_2$, $-C(=O)-NH-(CH_2CH_2O)_2CH_2$ -COOH, or $-C(=O)-NR^{50A}CH_2CH_2$.

164. A pharmaceutical composition according to claim 162 wherein R^{50} is phenyl, methyl or ethyl.

165. A pharmaceutical composition according to claim 164 wherein R^{50} is methyl or ethyl.

166. A pharmaceutical composition according to any one of the claims 127 to 165 wherein m is 1 or 2.

167. A pharmaceutical composition according to any one of the claims 127 to 166 wherein R^{51} is methyl.


168. A pharmaceutical composition according to any one of the claims 127 to 167 wherein R^{53} is C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, $-OR^{51}$, halogen, or $-CF_3$.

169. A pharmaceutical composition according to any one of the claims 127 to 168 wherein R^{50A} is $-C(O)OCH_3$, $-C(O)OCH_2CH_3$, $-COOH$, $-CH_2C(O)OCH_3$, $-CH_2C(O)OCH_2CH_3$, $-CH_2CH_2C(O)OCH_3$, $-CH_2CH_2C(O)OCH_2CH_3$, $-CH_2COOH$, methyl, or ethyl.

170. A pharmaceutical composition according to any one of the claims 127 to 169 wherein R^{50B} is $-C(O)OCH_3$, $-C(O)OCH_2CH_3$, $-COOH$, $-CH_2C(O)OCH_3$, $-CH_2C(O)OCH_2CH_3$, $-CH_2CH_2C(O)OCH_3$, $-CH_2CH_2C(O)OCH_2CH_3$, $-CH_2COOH$, methyl, or ethyl.

171. A pharmaceutical composition according to claim 1 wherein the zinc-binding ligand is

5



wherein V is C_1 - C_6 -alkyl, aryl, heteroaryl, aryl- C_{1-6} -alkyl- or aryl- C_{2-6} -alkenyl-, wherein the alkyl or alkenyl is optionally substituted with one or more substituents independently selected from R^{54} , and the aryl or heteroaryl is optionally substituted with one or more substituents independently selected from R^{55} .

R^{54} is independently selected from halogen, $-CN$, $-CF_3$, $-OCF_3$, aryl, $-COOH$ and $-NH_2$,

R^{55} is independently selected from

15 • hydrogen, halogen, $-CN$, $-CH_2CN$, $-CHF_2$, $-CF_3$, $-OCF_3$, $-OCHF_2$, $-OCH_2CF_3$, $-OCF_2CHF_2$, $-S(O)_2CF_3$, $-OS(O)_2CF_3$, $-SCF_3$, $-NO_2$, $-OR^{56}$, $-NR^{56}R^{57}$, $-SR^{56}$, $-NR^{56}S(O)_2R^{57}$, $-S(O)_2NR^{56}R^{57}$, $-S(O)NR^{56}R^{57}$, $-S(O)R^{56}$, $-S(O)_2R^{56}$, $-OS(O)_2R^{56}$, $-C(O)NR^{56}R^{57}$, $-OC(O)NR^{56}R^{57}$, $-NR^{56}C(O)R^{57}$, $-CH_2C(O)NR^{56}R^{57}$, $-OC_1-C_6$ -alkyl- $C(O)NR^{56}R^{57}$, $-CH_2OR^{56}$, $-CH_2OC(O)R^{56}$, $-CH_2NR^{56}R^{57}$, $-OC(O)R^{56}$, $-OC_1-C_8$ -alkyl- $C(O)OR^{56}$, $-OC_1-C_6$ -alkyl- OR^{56} , $-SC_1-C_6$ -alkyl- $C(O)OR^{56}$, $-C_2-C_6$ -alkenyl- $C(=O)OR^{56}$, $-NR^{56}-C(=O)-C_1-C_6$ -alkyl- $C(=O)OR^{56}$, $-NR^{56}-C(=O)-C_1-C_6$ -alkenyl- $C(=O)OR^{56}$, $-C(O)OR^{56}$, or $-C_2-C_6$ -alkenyl- $C(=O)R^{56}$,

20 • C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl or C_2 - C_6 -alkynyl,

25

which may optionally be substituted with one or more substituents selected from R^{58} ,

30 • aryl, aryloxy, aryloxycarbonyl, aroyl, arylsulfanyl, aryl- C_1 - C_6 -alkoxy, aryl- C_1 - C_6 -alkyl, aryl- C_2 - C_6 -alkenyl, aroyl- C_2 - C_6 -alkenyl, aryl- C_2 - C_6 -alkynyl, heteroaryl, heteroaryl- C_1 - C_6 -alkyl, heteroaryl- C_2 - C_6 -alkenyl or heteroaryl- C_2 - C_6 -alkynyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R^{59} ,

R^{56} and R^{57} are independently selected from hydrogen, OH, CF_3 , C_1 - C_{12} -alkyl, aryl- C_1 - C_6 -alkyl, $-C(=O)$ - C_1 - C_6 -alkyl or aryl, wherein the alkyl groups may optionally be substituted with one or more substituents independently selected from R^{60} , and the aryl groups may optionally be substituted with one or more substituents independently selected from R^{61} ; R^{56} and R^{57} when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom, the heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds,

R^{58} is independently selected from halogen, $-CN$, $-CF_3$, $-OCF_3$, $-OR^{56}$, and $-NR^{56}R^{57}$,

R^{59} is independently selected from halogen, $-C(O)OR^{58}$, $-CH_2C(O)OR^{58}$, $-CH_2OR^{58}$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OR^{58}$, $-NR^{56}R^{57}$ and C_1 - C_6 -alkyl,

R^{60} is independently selected from halogen, $-CN$, $-CF_3$, $-OCF_3$, $-OC_1$ - C_6 -alkyl, $-C(O)OC_1$ - C_6 -alkyl, $-C(=O)R^{62}$, $-COOH$ and $-NH_2$,

R^{61} is independently selected from halogen, $-C(O)OC_1$ - C_6 -alkyl, $-COOH$, $-CN$, $-CF_3$, $-OCF_3$, $-NO_2$, $-OH$, $-OC_1$ - C_6 -alkyl, $-NH_2$, $C(=O)$ or C_1 - C_6 -alkyl,

R^{62} is C_1 - C_6 -alkyl, aryl optionally substituted with one or more substituents independently selected from halogen, or heteroaryl optionally substituted with one or more C_1 - C_6 -alkyl independently,

or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.

172. A pharmaceutical composition according to claim 171 wherein V is aryl, heteroaryl, or aryl- C_{1-6} -alkyl-, wherein the alkyl is optionally substituted with one or more substituents independently selected R^{54} , and the aryl or heteroaryl is optionally substituted with one or more substituents independently selected from R^{55} .

173. A pharmaceutical composition according to claim 172 wherein V is aryl, Het1, or aryl- C_{1-6} -alkyl-, wherein the alkyl is optionally substituted with one or more substituents independently selected from R^{54} , and the aryl or heteroaryl moiety is optionally substituted with one or more substituents independently selected from R^{55} .

174. A pharmaceutical composition according to claim 173 wherein V is aryl, Het2, or aryl-C₁₋₆-alkyl-, wherein the alkyl is optionally substituted with one or more substituents independently selected from R⁵⁴, and the aryl or heteroaryl moiety is optionally substituted with one or more substituents independently selected from R⁵⁵.

5 175. A pharmaceutical composition according to claim 174 wherein V is aryl, Het3, or aryl-C₁₋₆-alkyl-, wherein the alkyl is optionally substituted with one or more substituents independently selected from R⁵⁴, and the aryl or heteroaryl moiety is optionally substituted with one or more substituents independently selected from R⁵⁵.

176. A pharmaceutical composition according to claim 175 wherein V is aryl optionally substituted with one or more substituents independently selected from R⁵⁵.

177. A pharmaceutical composition according to claim 176 wherein V is ArG1 optionally substituted with one or more substituents independently selected from R⁵⁵.

178. A pharmaceutical composition according to claim 177 wherein V is phenyl, naphthyl or anthranlyl optionally substituted with one or more substituents independently selected from R⁵⁵.

179. A pharmaceutical composition according to claim 178 wherein V is phenyl optionally substituted with one or more substituents independently selected from R⁵⁵.

180. A pharmaceutical composition according to any one of the claims 171 to 179 wherein R⁵⁵ is independently selected from

20 • halogen, C₁-C₆-alkyl, -CN, -OCF₃, -CF₃, -NO₂, -OR⁵⁶, -NR⁵⁶R⁵⁷, -NR⁵⁶C(O)R⁵⁷-SR⁵⁶, -OC₁-C₈-alkyl-C(O)OR⁵⁶, or -C(O)OR⁵⁶,
• C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R⁵⁸
• aryl, aryl-C₁-C₆-alkyl, heteroaryl, or heteroaryl-C₁-C₆-alkyl
25 of which the cyclic moieties optionally may be substituted with one or more substituents independently selected from R⁵⁹.

181. A pharmaceutical composition according to claim 180 wherein R⁵⁵ is independently selected from

30 • halogen, C₁-C₆-alkyl, -CN, -OCF₃, -CF₃, -NO₂, -OR⁵⁶, -NR⁵⁶R⁵⁷, -NR⁵⁶C(O)R⁵⁷-SR⁵⁶, -OC₁-C₈-alkyl-C(O)OR⁵⁶, or -C(O)OR⁵⁶
• C₁-C₆-alkyl optionally substituted with one or more substituents independently selected from R⁵⁸
• ArG1, ArG1-C₁-C₆-alkyl, Het3, or Het3-C₁-C₆-alkyl
35 of which the cyclic moieties optionally may be substituted with one or more substituents independently selected from R⁵⁹.

182. A pharmaceutical composition according to claim 181 wherein R^{55} is independently selected from halogen, $-OR^{56}$, $-NR^{56}R^{57}$, $-C(O)OR^{56}$, $-OC_1-C_8\text{-alkyl-C}(O)OR^{56}$, $-NR^{56}C(O)R^{57}$ or $C_1-C_6\text{-alkyl}$.


5 183. A pharmaceutical composition according to claim 182 wherein R^{55} is independently selected from halogen, $-OR^{56}$, $-NR^{56}R^{57}$, $-C(O)OR^{56}$, $-OC_1-C_8\text{-alkyl-C}(O)OR^{56}$, $-NR^{56}C(O)R^{57}$, methyl or ethyl.

184. A pharmaceutical composition according to any one of the claims 171 to 183 wherein R^{56} and R^{57} are independently selected from hydrogen, CF_3 , $C_1-C_{12}\text{-alkyl}$, or
10 $-C(=O)-C_1-C_6\text{-alkyl}$; R^{56} and R^{57} when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom.

185. A pharmaceutical composition according to claim 184 wherein R^{56} and R^{57} are independently selected from hydrogen or $C_1-C_{12}\text{-alkyl}$, R^{56} and R^{57} when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen
15 atom.

186. A pharmaceutical composition according to claim 185 wherein R^{56} and R^{57} are independently selected from hydrogen or methyl, ethyl, propyl butyl, R^{56} and R^{57} when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic ring with the said nitrogen atom.

20 187. A pharmaceutical composition according to claim 1 wherein the zinc-binding ligand is



wherein AA is $C_1-C_6\text{-alkyl}$, aryl, heteroaryl, aryl- $C_{1-6}\text{-alkyl}$ - or aryl- $C_{2-6}\text{-alkenyl}$ -, wherein the
25 alkyl or alkenyl is optionally substituted with one or more substituents independently selected from R^{63} , and the aryl or heteroaryl is optionally substituted with one or more substituents independently selected from R^{64} ,

R^{63} is independently selected from halogen, $-CN$, $-CF_3$, $-OCF_3$, aryl, $-COOH$ and $-NH_2$,

30

R^{64} is independently selected from

- 5 • hydrogen, halogen, -CN, -CH₂CN, -CHF₂, -CF₃, -OCF₃, -OCHF₂, -OCH₂CF₃, -OCF₂CHF₂, -S(O)₂CF₃, -OS(O)₂CF₃, -SCF₃, -NO₂, -OR⁶⁵, -NR⁶⁵R⁶⁶, -SR⁶⁵, -NR⁶⁵S(O)₂R⁶⁶, -S(O)₂NR⁶⁵R⁶⁶, -S(O)NR⁶⁵R⁶⁶, -S(O)R⁶⁵, -S(O)₂R⁶⁵, -OS(O)₂R⁶⁵, -C(O)NR⁶⁵R⁶⁶, -OC(O)NR⁶⁵R⁶⁶, -NR⁶⁵C(O)R⁶⁶, -CH₂C(O)NR⁶⁵R⁶⁶, -OC₁-C₆-alkyl-C(O)NR⁶⁵R⁶⁶, -CH₂OR⁶⁵, -CH₂OC(O)R⁶⁵, -CH₂NR⁶⁵R⁶⁶, -OC(O)R⁶⁵, -OC₁-C₆-alkyl-C(O)OR⁶⁵, -OC₁-C₆-alkyl-OR⁶⁵, -SC₁-C₆-alkyl-C(O)OR⁶⁵, -C₂-C₆-alkenyl-C(=O)OR⁶⁵, -NR⁶⁵-C(=O)-C₁-C₆-alkyl-C(=O)OR⁶⁵, -NR⁶⁵-C(=O)-C₁-C₆-alkenyl-C(=O)OR⁶⁵, -C(O)OR⁶⁵, or -C₂-C₆-alkenyl-C(=O)R⁶⁵,
- 10 • C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl, each of which may optionally be substituted with one or more substituents selected from R⁶⁷,
- 15 • aryl, aryloxy, aryloxycarbonyl, aroyl, arylsulfanyl, aryl-C₁-C₆-alkoxy, aryl-C₁-C₆-alkyl, aryl-C₂-C₆-alkenyl, aroyl-C₂-C₆-alkenyl, aryl-C₂-C₆-alkynyl, heteroaryl, heteroaryl-C₁-C₆-alkyl, heteroaryl-C₂-C₆-alkenyl or heteroaryl-C₂-C₆-alkynyl,

of which the cyclic moieties optionally may be substituted with one or more substituents selected from R⁶⁸,

- 20 R⁶⁵ and R⁶⁶ are independently selected from hydrogen, OH, CF₃, C₁-C₁₂-alkyl, aryl-C₁-C₆-alkyl, -C(=O)-R⁶⁹, aryl or heteroaryl, wherein the alkyl groups may optionally be substituted with one or more substituents selected from R⁷⁰, and the aryl and heteroaryl groups may optionally be substituted with one or more substituents independently selected from R⁷¹; R⁶⁵ and R⁶⁶ when attached to the same nitrogen atom may form a 3 to 8 membered heterocyclic
- 25 ring with the said nitrogen atom, the heterocyclic ring optionally containing one or two further heteroatoms selected from nitrogen, oxygen and sulphur, and optionally containing one or two double bonds,

R⁶⁷ is independently selected from halogen, -CN, -CF₃, -OCF₃, -OR⁶⁵, and -NR⁶⁵R⁶⁶,

30

R⁶⁸ is independently selected from halogen, -C(O)OR⁶⁵, -CH₂C(O)OR⁶⁵, -CH₂OR⁶⁵, -CN, -CF₃, -OCF₃, -NO₂, -OR⁶⁵, -NR⁶⁵R⁶⁶ and C₁-C₆-alkyl,

- 35 R⁶⁹ is independently selected from C₁-C₆-alkyl, aryl optionally substituted with one or more halogen, or heteroaryl optionally substituted with one or more C₁-C₆-alkyl,

R⁷⁰ is independently selected from halogen, -CN, -CF₃, -OCF₃, -OC₁-C₆-alkyl, -C(O)OC₁-C₆-alkyl, -COOH and -NH₂,

- 5 R⁷¹ is independently selected from halogen, -C(O)OC₁-C₆-alkyl, -COOH, -CN, -CF₃, -OCF₃, -NO₂, -OH, -OC₁-C₆-alkyl, -NH₂, C(=O) or C₁-C₆-alkyl,

or any enantiomer, diastereomer, including a racemic mixture, tautomer as well as a salt thereof with a pharmaceutically acceptable acid or base.

- 10 188. A pharmaceutical composition according to claim 187 wherein AA is aryl, heteroaryl or aryl-C₁₋₆-alkyl-, wherein the alkyl is optionally substituted with one or more R⁶³, and the aryl or heteroaryl is optionally substituted with one or more substituents independently selected from R⁶⁴.

189. A pharmaceutical composition according to claim 188 wherein AA is aryl or heteroaryl optionally substituted with one or more substituents independently selected from R⁶⁴.

190. A pharmaceutical composition according to claim 189 wherein AA is ArG1 or Het1 optionally substituted with one or more substituents independently selected from R⁶⁴.

191. A pharmaceutical composition according to claim 190 wherein AA is ArG1 or Het2 optionally substituted with one or more substituents independently selected from R⁶⁴.

- 20 192. A pharmaceutical composition according to claim 191 wherein AA is ArG1 or Het3 optionally substituted with one or more substituents independently selected from R⁶⁴.

193. A pharmaceutical composition according to claim 192 wherein AA is phenyl, naphthyl, anthryl, carbazolyl, thienyl, pyridyl, or benzodioxyl optionally substituted with one or more substituents independently selected from R⁶⁴.

- 25 194. A pharmaceutical composition according to claim 193 wherein AA is phenyl or naphthyl optionally substituted with one or more substituents independently selected from R⁶⁴.

195. A pharmaceutical composition according to any one of the claims 187 to 194 wherein R⁶⁴ is independently selected from hydrogen, halogen, -CF₃, -OCF₃, -OR⁶⁵, -NR⁶⁵R⁶⁶, C₁-C₆-alkyl, -OC(O)R⁶⁵, -OC₁-C₆-alkyl-C(O)OR⁶⁵, aryl-C₂-C₆-alkenyl, aryloxy or aryl, wherein C₁-

- 30 C₆-alkyl is optionally substituted with one or more substituents independently selected from R⁶⁷, and the cyclic moieties optionally are substituted with one or more substituents independently selected from R⁶⁸.

196. A pharmaceutical composition according to claim 195 wherein R⁶⁴ is independently selected from halogen, -CF₃, -OCF₃, -OR⁶⁵, -NR⁶⁵R⁶⁶, methyl, ethyl, propyl, -OC(O)R⁶⁵,

-OCH₂-C(O)OR⁶⁵, -OCH₂-CH₂-C(O)OR⁶⁵, phenoxy optionally substituted with one or more substituents independently selected from R⁶⁸.

197. A pharmaceutical composition according to any one of the claims 187 to 196 wherein R⁶⁵ and R⁶⁶ are independently selected from hydrogen, CF₃, C₁-C₁₂-alkyl, aryl, or heteroaryl optionally substituted with one or more substituents independently selected from R⁷¹.

198. A pharmaceutical composition according to claim 197 wherein R⁶⁵ and R⁶⁶ are independently hydrogen, C₁-C₁₂-alkyl, aryl, or heteroaryl optionally substituted with one or more substituents independently selected from R⁷¹.

199. A pharmaceutical composition according to claim 198 wherein R⁶⁵ and R⁶⁶ are independently hydrogen, methyl, ethyl, propyl, butyl, 2,2-dimethyl-propyl, ArG1 or Het1 optionally substituted with one or more substituents independently selected from R⁷¹.

200. A pharmaceutical composition according to claim 199 wherein R⁶⁵ and R⁶⁶ are independently hydrogen, methyl, ethyl, propyl, butyl, 2,2-dimethyl-propyl, ArG1 or Het2 optionally substituted with one or more substituents independently selected from R⁷¹.

201. A pharmaceutical composition according to claim 200 wherein R⁶⁵ and R⁶⁶ are independently hydrogen, methyl, ethyl, propyl, butyl, 2,2-dimethyl-propyl, ArG1 or Het3 optionally substituted with one or more substituents independently selected from R⁷¹.

202. A pharmaceutical composition according to claim 201 wherein R⁶⁵ and R⁶⁶ are independently hydrogen, methyl, ethyl, propyl, butyl, 2,2-dimethyl-propyl, phenyl, naphthyl, thiadiazolyl optionally substituted with one or more R⁷¹ independently; or isoxazolyl optionally substituted with one or more substituents independently selected from R⁷¹.

203. A pharmaceutical composition according to any one of the claims 187 to 202 wherein R⁷¹ is halogen or C₁-C₆-alkyl.

204. A pharmaceutical composition according to claim 203 wherein R⁷¹ is halogen or methyl.

205. A pharmaceutical composition according to any one of the claims 1 to 204 wherein the insulin is rapid acting insulin.

206. A pharmaceutical composition according to any one of the claims 1 to 205 wherein the insulin is selected from the group consisting of human insulin, an analogue thereof, a derivative thereof, and combinations of any of these.

207. A pharmaceutical composition according to claim 206 wherein the insulin is an analogue of human insulin selected from the group consisting of

iii. An analogue wherein position B28 is Asp, Lys, Leu, Val, or Ala and position B29 is Lys or Pro; and

iv. des(B28-B30), des(B27) or des(B30) human insulin.

208. A pharmaceutical composition according to claim 207, wherein the insulin is an analogue of human insulin wherein position B28 is Asp or Lys, and position B29 is Lys or Pro.

209. A pharmaceutical composition according to claim 207 wherein the insulin is des(B30) human insulin.

210. A pharmaceutical composition according to claim 207 wherein the insulin is an analogue of human insulin wherein position B3 is Lys and position B29 is Glu or Asp.

211. A pharmaceutical composition according to claim 206 wherein the insulin is a derivative of human insulin having one or more lipophilic substituents.

212. A pharmaceutical composition according to claim 211 wherein the insulin derivative is selected from the group consisting of B29-N^E-myristoyl-des(B30) human insulin, B29-N^E-palmitoyl-des(B30) human insulin, B29-N^E-myristoyl human insulin, B29-N^E-palmitoyl human insulin, B28-N^E-myristoyl Lys^{B28} Pro^{B29} human insulin, B28-N^E-palmitoyl Lys^{B28} Pro^{B29} human insulin, B30-N^E-myristoyl-Thr^{B29}Lys^{B30} human insulin, B30-N^E-palmitoyl-Thr^{B29}Lys^{B30} human insulin, B29-N^E-(N-palmitoyl-γ-glutamyl)-des(B30) human insulin, B29-N^E-(N-lithocholyl-γ-glutamyl)-des(B30) human insulin, B29-N^E-(ω-carboxyheptadecanoyl)-des(B30) human insulin and B29-N^E-(ω-carboxyheptadecanoyl) human insulin.

213. A pharmaceutical composition according to claim 212 wherein the insulin derivative is B29-N^E-myristoyl-des(B30) human insulin.

214. A pharmaceutical composition according to any one of the claims 1 to 213 comprising 2-6 moles zinc²⁺ ions per mole insulin.

215. A pharmaceutical composition according to claim 214 comprising 2-3 moles zinc²⁺ ions per mole insulin.

216. A pharmaceutical composition according to any one of the claims 1 to 215 further comprising at least 3 molecules of a phenolic compound per insulin hexamer.

217. A pharmaceutical composition according to any one of the claims 1 to 216 further comprising an isotonicity agent.

218. A pharmaceutical composition according to any one of the claims 1 to 217 further comprising a buffer substance.

219. A method of stabilising an insulin composition comprising adding a zinc-binding ligand according to any one of the claims 1 to 204 to the insulin composition.

220. A method of treating type 1 or type 2 diabetes comprising administering to a patient in need thereof a pharmaceutically effective dose of an insulin composition according to any one of the claims 1 to 204.

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
8 July 2004 (08.07.2004)

PCT

(10) International Publication Number
WO 2004/056347 A3

(51) International Patent Classification⁷: **A61K 38/28**,
C07D 249/18, 277/34, A61K 31/4192, 31/416, C07D
417/02, 413/02, C07K 5/00, 7/00, 14/62

[DK/DK]; Nærum Hovedgade 64G, DK-2850 Nærum
(DK). **SCHLUCKEBIER, Gerd** [DE/DK]; Vævergade 5,
4th, DK-2200 København N (DK).

(21) International Application Number:
PCT/DK2003/000931

(74) Common Representative: **NOVO NORDISK A/S**; Nove
Allé, DK-2880 Bagsværd (DK).

(22) International Filing Date:
22 December 2003 (22.12.2003)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR,
CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,
MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU,
SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PA 2002 01991 20 December 2002 (20.12.2002) DK
60/439,382 10 January 2003 (10.01.2003) US

(84) Designated States (*regional*): ARIPO patent (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **NOVO
NORDISK A/S** [DK/DK]; Nove Allé, DK-2880 Bagsværd
(DK).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **KAARSHOLM**,
Niels, Christian [DK/DK]; 38, Clausholmvej, DK-2720
Vanløse (DK). **MADSEN, Peter** [DK/DK]; 7 Ulveh-
jerg, DK-2880 Bagsværd (DK). **SCHLEIN, Morten**
[DK/DK]; Under Elmene 16, 4. tv., DK-2300 Copenhagen
S (DK). **OLSEN, Helle, Birk** [DK/DK]; Skoleloddén 23,
DK-3450 Allerød (DK). **HAVELUND, Svend** [DK/DK];
Kurvej 24, DK-2880 Bagsværd (DK). **STEENSGAARD**,
Dorte, Bjerre [DK/DK]; Stockholmsgade 3, st. tv.,
DK- 2100_København Ø (DK). **LUDVIGSEN, Svend**
[DK/DK]; 13, Baunedalen, DK-3540 Lynge (DK).
JAKOBSEN, Palle [DK/DK]; 14 Langkær Vænge,
DK-3500 Værløse (DK). **PETERSEN, Anders, Klarskov**

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(88) Date of publication of the international search report:
12 August 2004

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: PHARMACEUTICAL COMPOSITIONS COMPRISING INSULIN AND LEGENDS OF INSULIN HEXAMER

(57) Abstract: The present invention provides pharmaceutical compositions comprising insulin and novel ligands for the His^{B10}
Zn²⁺ sites of the R-state insulin hexamer. The resulting preparations have improved physical and chemical stability.



WO 2004/056347 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/DK 03/00931

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K38/28 C07D249/18 C07D277/34 A61K31/4192 A61K31/416
 C07D417/02 C07D413/02 C07K5/00 C07K7/00 C07K14/62

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, MEDLINE, CHEM ABS Data, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|-----------------------|
| P,X | WO 03 027081 A (KAARSHOLM NIELS C ;JAKOBSEN PALLE (DK); LUDVIGSEN SVEND (DK); MADSEN) 3 April 2003 (2003-04-03) claims 1-204 --- | 1-220 |
| P,X | US 2003/229120 A1 (KAARSHOLM NIELS C ET AL) 11 December 2003 (2003-12-11) claims 1-204 --- | 1-220 |
| X | US 5 830 999 A (DUNN MICHAEL F) 3 November 1998 (1998-11-03) claims 1-6 --- -/- | 1-220 |



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

13 May 2004

Date of mailing of the international search report

07. 06. 2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

FERNANDO FARIETA /EÖ

INTERNATIONAL SEARCH REPORT

International Application No

PCT/DK 03/00931

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|---|-----------------------|
| A | FERRARI D ET AL: "Raman signatures of ligand binding and allosteric conformation change in hexameric insulin." BIOPOLYMERS. UNITED STATES 2001, vol. 62, no. 5, 2001, pages 249-260, XP002280143 ISSN: 0006-3525 Discussion --- | 1-220 |
| A | HUANG S T ET AL: "Carboxylate ions are strong allosteric ligands for the HisB10 sites of the R-state insulin hexamer." BIOCHEMISTRY. UNITED STATES 12 AUG 1997, vol. 36, no. 32, 12 August 1997 (1997-08-12), pages 9878-9888, XP002280144 ISSN: 0006-2960 scheme 1 table 1 --- | 1-220 |
| A | BRADER M L ET AL: "Characterization of the R-state insulin hexamer and its derivatives. The hexamer is stabilized by heterotropic ligand binding interactions." BIOCHEMISTRY. UNITED STATES 9 JUL 1991, vol. 30, no. 27, 9 July 1991 (1991-07-09), pages 6636-6645, XP002280145 ISSN: 0006-2960 table 1 --- | 1-220 |
| A | US 4 476 118 A (BRANGE JENS J V ET AL) 9 October 1984 (1984-10-09) claims 1-16 --- | 1-220 |
| E | EP 1 396 272 A (LILLY CO ELI) 10 March 2004 (2004-03-10) claims 1-38 ----- | 1-220 |

INTERNATIONAL SEARCH REPORT

International application No.
PCT/DK 03/00931

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 220
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☒ Claims Nos.: 1-218
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☒ Claims Nos.: 219
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Claims Nos.: 220

Claims 220 relate to method of treatment of the human or animal body by surgery or by therapy or diagnostic methods practised on the human or animal body (PCT Rule 39.1(iv)). Nevertheless, a search has been executed for these claim. The search has been based on the alleged effects of the compounds or compositions.

Continuation of Box I.2

Claims Nos.: 1-218

Present claims 1-219 relate to a pharmaceutical composition (insulin and zinc-ligand) defined by reference to a desirable characteristic or property, namely a zinc ligand which reversibly binds to a HisB10Zn²⁺ site of an insulin hexamer.

The claims cover all possible compounds (benzotriazoles, tetrazoles, racemic mixtures etc) having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and / or disclosure within the meaning of Article 5 PCT for only a very limited number of such compounds. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the compositions by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible.

Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the benzotriazoles, tetrazoles and thiazolidinediones prepared in examples 1-1009.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/DK 03/00931

| Patent document cited in search report | | Publication date | Patent family member(s) | Publication date |
|---|----|---------------------|----------------------------|---------------------|
| WO 03027081 | A | 03-04-2003 | WO 03027081 A2 | 03-04-2003 |
| | | | US 2003229120 A1 | 11-12-2003 |
| US 2003229120 | A1 | 11-12-2003 | WO 03027081 A2 | 03-04-2003 |
| US 5830999 | A | 03-11-1998 | NONE | |
| US 4476118 | A | 09-10-1984 | AT 382780 B | 10-04-1987 |
| | | | AT 38983 A | 15-09-1986 |
| | | | AU 549981 B2 | 20-02-1986 |
| | | | AU 1113683 A | 11-08-1983 |
| | | | BE 895832 A1 | 04-08-1983 |
| | | | CA 1203476 A1 | 22-04-1986 |
| | | | CH 657530 A5 | 15-09-1986 |
| | | | DE 3303860 A1 | 18-08-1983 |
| | | | DK 42483 A ,B, | 06-08-1983 |
| | | | ES 8500060 A1 | 01-01-1985 |
| | | | FI 830375 A ,B, | 06-08-1983 |
| | | | FR 2521009 A1 | 12-08-1983 |
| | | | GB 2114573 A ,B | 24-08-1983 |
| | | | GR 77388 A1 | 11-09-1984 |
| | | | IE 54548 B1 | 08-11-1989 |
| | | | IT 1178455 B | 09-09-1987 |
| | | | JP 1707986 C | 27-10-1992 |
| | | | JP 3067056 B | 21-10-1991 |
| | | | JP 58192818 A | 10-11-1983 |
| | | | LU 84632 A1 | 08-09-1983 |
| | | | NL 8300440 A ,C | 01-09-1983 |
| | | | NO 830385 A ,B, | 08-08-1983 |
| | | | NZ 203189 A | 31-07-1985 |
| | | | PT 76195 A ,B | 01-03-1983 |
| | | | SE 460948 B | 11-12-1989 |
| | | | SE 8300598 A | 06-08-1983 |
| | | | ZA 8300742 A | 30-11-1983 |
| EP 1396272 | A | 10-03-2004 | EP 1396272 A1 | 10-03-2004 |
| | | | AU 2008899 A | 12-07-1999 |
| | | | BR 9814471 A | 10-10-2000 |
| | | | CA 2315300 A1 | 01-07-1999 |
| | | | CN 1284876 T | 21-02-2001 |
| | | | EA 3394 B1 | 24-04-2003 |
| | | | EP 0925792 A2 | 30-06-1999 |
| | | | HR 20000427 A1 | 30-06-2001 |
| | | | HU 0100243 A2 | 28-08-2001 |
| | | | ID 25476 A | 05-10-2000 |
| | | | JP 2001526225 T | 18-12-2001 |
| | | | NO 20003269 A | 23-08-2000 |
| | | | NZ 505091 A | 26-11-2002 |
| | | | PL 341300 A1 | 09-04-2001 |
| | | | TR 200001935 T2 | 21-12-2000 |
| | | | WO 9932116 A1 | 01-07-1999 |
| | | | US 6531448 B1 | 11-03-2003 |
| | | | US 2003144181 A1 | 31-07-2003 |
| | | | ZA 9811728 A | 27-06-2000 |